

Datasheet: MCA1369SBV610

BATCH NUMBER 100008143

Description:	HAMSTER ANTI MOUSE CD11c:StarBright Violet 610
Specificity:	CD11c
Other names:	INTEGRIN ALPHA X CHAIN
Format:	StarBright Violet 610
Product Type:	Monoclonal Antibody
Clone:	N418
Isotype:	IgG
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjugat	ed to StarBright Violet	610 - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	StarBright Violet 610	403	607
reparation	supernatant	d by affinity chromatog	raphy on Protein A f
ffer Solution	Phosphate buffered s	saline	
eservative	0.09% Sodium Azide	(NaN ₃)	
abilisers	1% Bovine Serum All	bumin	
	0.1% Pluronic F68		
	0.1% PEG 3350		

Immunoge	n
----------	---

Mouse spleen dendritic cells.

External Database Links

UniProt:

Q9QXH4 Related reagents

Entrez Gene:

16411 Itgax Related reagents

Fusion Partners

Spleen cells from immunised Armenian Hamster were fused with cells of the Sp2/0 myeloma cell line.

Specificity

Hamster anti Mouse CD11c antibody, clone N418 recognizes the murine homolog of human CD11c, also known as Integrin Alpha X, a 150/90 kDa member of the beta 2 integrin family. In mice, CD11c is primarily expressed by dendritic cells.

Hamster anti Mouse CD11c antibody, clone N418 has been reported to enhance antigen specific responses when used to target dendritic cells *in vivo* (Wang *et al.* 2000).

Flow Cytometry

Use 5μ I of the suggested working dilution to label 10^6 cells in 100μ I. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Crowley, M.T. *et al.* (1990) Use of the fluorescence activated cell sorter to enrich dendritic cells from mouse spleen. <u>J Immunol Methods</u>. 133 (1): 55-66.
- 2. Dahlen, E. *et al.* (1998) Dendritic cells and macrophages are the first and major producers of TNF-alpha in pancreatic islets in the nonobese diabetic mouse. <u>J Immunol.</u> 160: 3585-93.
- 3. Wang, H. *et al.* (2000) Rapid antibody responses by low-dose, single-step, dendritic cell-targeted immunization. Proc Natl Acad Sci U S A. 97 (2): 847-52.
- 4. Lundqvist, J. *et al.* (2010) Concomitant infection decreases the malaria burden but escalates relapsing fever borreliosis. Infect Immun. 78 (5): 1924-30.
- 5. Beyer, M. *et al.* (2005) The beta2 integrin CD11c distinguishes a subset of cytotoxic pulmonary T cells with potent antiviral effects in vitro and in vivo. Respir Res. 6: 70.
- 6. Goupil, M. *et al.* (2009) Macrophage-mediated responses to Candida albicans in mice expressing the human immunodeficiency virus type 1 transgene. <u>Infect Immun. 77:</u> 4136-49.
- 7. Linehan, S.A.. (2005) The mannose receptor is expressed by subsets of APC in non-lymphoid organs. BMC Immunol. 6:4.
- 8. Bjorck, P. (2004) Dendritic cells exposed to herpes simplex virus *in vivo* do not produce IFN-alpha after rechallenge with virus *in vitro* and exhibit decreased T cell alloreactivity. J Immunol. 172: 5396-404.
- 9. Mercier, S. *et al.* (2002) Distinct roles of adenovirus vector-transduced dendritic cells, myoblasts, and endothelial cells in mediating an immune response against a transgene product. J Virol. 76: 2899-911.
- 10. de Jersey, J. *et al.* (2002) Activation of CD8 T cells by antigen expressed in the pituitary gland. <u>J Immunol. 169: 6753-9.</u>

- 11. Dimier-Poisson, I. *et al.* (2003) Protective mucosal Th2 immune response against Toxoplasma gondii by murine mesenteric lymph node dendritic cells. <u>Infect Immun. 71:</u> 5254-65.
- 12. Gonzalez-Juarrero, M. and Orme, I.M. (2001) Characterization of murine lung dendritic cells infected with *Mycobacterium tuberculosis*. <u>Infect Immun. 69: 1127-33.</u>
- 13. Hamada, H. *et al.* (2002) Identification of multiple isolated lymphoid follicles on the antimesenteric wall of the mouse small intestine. J Immunol. 168: 57-64.
- 14. Meng, Q. *et al.* (2007) Phenotypes, distribution, and morphological features of antigen-presenting cells in the murine cornea following intravitreal injection. <u>Mol Vis. 13: 475-86.</u>
- 15. Moos, M.P. *et al.* (2005) The lamina adventitia is the major site of immune cell accumulation in standard chow-fed apolipoprotein E-deficient mice <u>Arterioscler Thromb</u> Vasc Biol. 25: 2386-91.
- 16. Nunez, R. *et al.* (1999) Immortalized cell lines derived from mice lacking both type I and type II IFN receptors unify some functions of immature and mature dendritic cells. <u>Immunol Cell Biol. 77: 153-63.</u>
- 17. Ponce, L.V. *et al.* (2005) Adoptive transfer of dendritic cells modulates immunogenesis and tolerogenesis in a neonatal model of murine cutaneous leishmaniasis. <u>Kinetoplastid Biol Dis. 4: 2.</u>
- 18. Zhang, L. *et al.* (2011) The inflammatory changes of adipose tissue in late pregnant mice. <u>J Mol Endocrinol</u>. 47 (2): 157-65.
- 19. Donaldson, D.S. *et al.* (2012) M cell-depletion blocks oral prion disease pathogenesis. Mucosal Immunol. 5: 216-25.
- 20. Wada, T. *et al.* (2013) Eplerenone ameliorates the phenotypes of metabolic syndrome with NASH in liver-specific SREBP-1c Tg mice fed high-fat and high-fructose diet. <u>Am J Physiol Endocrinol Metab.</u> 305 (11): E1415-25.
- 21. Kan, M.J. *et al.* (2015) Arginine deprivation and immune suppression in a mouse model of Alzheimer's disease. J Neurosci. 35 (15): 5969-82.
- 22. Kayser, B.D. *et al.* (2015) Perinatal Overnutrition Exacerbates Adipose Tissue Inflammation Caused by High-Fat Feeding in C57BL/6J Mice. <u>PLoS One. 10 (3)</u>: e0121954.
- 23. Sehgal, A. *et al.* (2017) c-Rel is dispensable for the differentiation and functional maturation of M cells in the follicle-associated epithelium. <u>Immunobiology. 222 (2): 316-26.</u>
- 24. Bender, L.H. *et al.* (2020) Intratumoral Administration of a Novel Cytotoxic Formulation with Strong Tissue Dispersive Properties Regresses Tumor Growth and Elicits Systemic Adaptive Immunity in *In Vivo* Models. <u>Int J Mol Sci. 21 (12) Jun 24 [Epub ahead of print].</u>
- 25. lida, Y. *et al.* (2020) Local injection of CCL19-expressing mesenchymal stem cells augments the therapeutic efficacy of anti-PD-L1 antibody by promoting infiltration of immune cells. <u>J Immunother Cancer. 8 (2) [Epub ahead of print].</u>
- 26. Kouwenberg, M. *et al.* (2021) Reduced CXCL1 production by endogenous IL-37 expressing dendritic cells does not affect T cell activation. <u>PLoS One. 16 (5): e0251809.</u>
- 27. Curina, G. *et al.* (2018) Evaluation of immune responses in mice and sheep inoculated with a live attenuated Brucella melitensis REV1 vaccine produced in bioreactor. <u>Vet Immunol Immunopathol.</u> 198: 44-53.
- 28. Mohanta, S.K. *et al.* (2022) Neuroimmune cardiovascular interfaces control atherosclerosis. Nature. 605 (7908): 152-159.
- 29. Mihindukulasooriya, P.S. *et al.* (2022) Polyphenol-rich Sargassum horneri alleviates atopic dermatitis-like skin lesions in NC/Nga mice by suppressing Th2-mediated cytokine

IL-13 Algae. 37 (4): 331-347.

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.	
Guarantee	12 months from date of despatch	
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and relaced counterparts	ated U.S. and foreign
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1369SBV610 20471	
Regulatory	For research purposes only	

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M421780:230726'

Fax: +44 (0)1865 852 739

Printed on 15 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint