

## Datasheet: MCA1370Z

**BATCH NUMBER 156607**

<b>Description:</b>	HAMSTER ANTI MOUSE CD31:Preservative Free
<b>Specificity:</b>	CD31
<b>Other names:</b>	PECAM-1
<b>Format:</b>	Preservative Free
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	2H8
<b>Isotype:</b>	IgG
<b>Quantity:</b>	0.5 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			0.1ug/ml
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			
Functional Assays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from ascites
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative</b>	None present.

<b>Stabilisers</b>	Sterile filtered.
<b>Approx. Protein Concentrations</b>	IgG concentration 0.5 mg/ml
<b>Immunogen</b>	D10.G4.1 cells ( <a href="#">Kaye et al. 1984</a> ).
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">Q08481</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">18613</a> Pecam1    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Pecam, Pecam-1
<b>RRID</b>	AB_321653
<b>Fusion Partners</b>	Splenic lymphocytes from an immunized Armenian hamster were fused with cells from the SP2/0 murine myeloma.
<b>Specificity</b>	<p><b>Hamster anti Mouse CD31 monoclonal antibody, clone 2H8</b> recognizes murine CD31, also known as Platelet endothelial cell adhesion molecule or PECAM-1. CD31 is a ~130kDa single pass type-1 membrane glycoprotein bearing six C2 Ig-like domains, expressed on all continuous endothelium including arteries, veins and non-sinusoidal capillaries. CD31 is also expressed on all haemopoietic lineages with the exception of the erythroid line (<a href="#">Bogen et al. 1992</a>) .</p> <p>Hamster anti mouse CD31, clone 2H8 effectively inhibits transmigration of activated polymorphonuclear cells and monocytes across the endothelium. In a mouse model for acute peritonitis clone 2H8 blocks acute inflammation (<a href="#">Bogen et al. 1994</a>). CD31 has also been shown to be critically involved in the sensing of changes in shear stress associated with atherosclerotic lesions and in the associated atherogenesis (<a href="#">Stevens et al. 2008</a>).</p>
<b>Flow Cytometry</b>	Use 5ul of the suggested working dilution to label 100ul of whole blood.
<b>References</b>	<ol style="list-style-type: none"> <li>1. Bogen, S.A. <i>et al.</i> (1992) Association of murine CD31 with transmigrating lymphocytes following antigenic stimulation. <a href="#">Am J Pathol. 141 (4): 843-54.</a></li> <li>2. Xie, Y. &amp; Muller, W.A. (1993) Molecular cloning and adhesive properties of murine platelet/endothelial cell adhesion molecule 1. <a href="#">Proc Natl Acad Sci U S A. 90 (12): 5569-73.</a></li> <li>3. Bogen, S. <i>et al.</i> (1994) Monoclonal antibody to murine PECAM-1 (CD31) blocks acute inflammation <i>in vivo</i>. <a href="#">J Exp Med. 179 (3): 1059-64.</a></li> <li>4. Bixel, M.G. <i>et al.</i> (2010) CD99 and CD99L2 act at the same site as, but independently of, PECAM-1 during leukocyte diapedesis. <a href="#">Blood. 116: 1172-84.</a></li> <li>5. Ishikawa, J. <i>et al.</i> (2002) Use of anti-platelet-endothelial cell adhesion molecule-1 antibody in the control of disease progression in established collagen-induced arthritis in DBA/1J mice. <a href="#">Jpn J Pharmacol. 88: 332-40.</a></li> <li>6. Thurston, G. <i>et al</i> (2005) Angiopoietin 1 causes vessel enlargement, without angiogenic</li> </ol>

- sprouting, during a critical developmental period [Development. 132: 3317-26.](#)
7. Lonsdorf, A.S. *et al.* (2012) Engagement of αIIbβ3 (GPIIb/IIIa) with αvβ3 Integrin Mediates Interaction of Melanoma Cells with Platelets: A CONNECTION TO HEMATOGENOUS METASTASIS. [J Biol Chem. 287: 2168-78.](#)
  8. Rijcken, E. *et al.* (2007) PECAM-1 (CD 31) mediates transendothelial leukocyte migration in experimental colitis. [Am J Physiol Gastrointest Liver Physiol. 293: G446-52.](#)
  9. Vielhauer, V. *et al.* (2005) Renal cell-expressed TNF receptor 2, not receptor 1, is essential for the development of glomerulonephritis. [J Clin Invest. 115: 1199-209.](#)
  10. Brackett, C.M. *et al.* (2013) IL-17 promotes neutrophil entry into tumor-draining lymph nodes following induction of sterile inflammation. [J Immunol. 191 \(8\): 4348-57.](#)
  11. Bixel, M.G. *et al.* (2010) CD99 and CD99L2 act at the same site as, but independently of, PECAM-1 during leukocyte diapedesis. [Blood. 116: 1172-84.](#)
  12. Wu, Y. *et al.* (2010) Therapeutic antibody targeting of individual Notch receptors. [Nature. 464: 1052-7.](#)
  13. Wilhelm, A. *et al.* (2015) CD248/Endosialin critically regulates hepatic stellate cell proliferation during chronic liver injury via a PDGF-regulated mechanism. [Gut. pii: gutjnl-2014-308325.](#)
  14. Smith, S.W. *et al.* (2015) Genetic Deletion of the Stromal Cell Marker CD248 (Endosialin) Protects against the Development of Renal Fibrosis. [Nephron. 131 \(4\): 265-77.](#)

---

**Storage**

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10162 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1370Z>  
10162

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight@550](#), [DyLight@650](#), [DyLight@800](#),  
[FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

'M365229:200529'

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

**Printed on 25 Mar 2023**

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)