

Datasheet: MCA1370Z

| | |
|----------------------|---|
| Description: | HAMSTER ANTI MOUSE CD31:Preservative Free |
| Specificity: | CD31 |
| Other names: | PECAM-1 |
| Format: | Preservative Free |
| Product Type: | Monoclonal Antibody |
| Clone: | 2H8 |
| Isotype: | IgG |
| Quantity: | 0.5 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 0.1ug/ml |
| Immunohistology - Frozen | | | ▪ | |
| Immunohistology - Paraffin | | | ▪ | |
| ELISA | | | ▪ | |
| Immunoprecipitation | | | ▪ | |
| Western Blotting | | | ▪ | |
| Immunofluorescence | ▪ | | | |
| Functional Assays | ▪ | | | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | |
|---------------------------------|--|
| Target Species | Mouse |
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein G from ascites |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | None present. |

Sterile filtered.

| | |
|---------------------------------------|-----------------------------|
| Approx. Protein Concentrations | IgG concentration 0.5 mg/ml |
|---------------------------------------|-----------------------------|

| | |
|------------------|--|
| Immunogen | D10.G4.1 cells (Kaye et al. 1984). |
|------------------|--|

| | |
|--------------------------------|--|
| External Database Links | UniProt: Q08481 Related reagents Entrez Gene: 18613 Pecam1 Related reagents |
|--------------------------------|--|

| | |
|-----------------|----------------|
| Synonyms | Pecam, Pecam-1 |
|-----------------|----------------|

| | |
|-------------|-----------|
| RRID | AB_321653 |
|-------------|-----------|

| | |
|------------------------|---|
| Fusion Partners | Splenic lymphocytes from an immunized Armenian hamster were fused with cells from the SP2/0 murine myeloma. |
|------------------------|---|

| | |
|--------------------|--|
| Specificity | <p>Hamster anti Mouse CD31 monoclonal antibody, clone 2H8 recognizes murine CD31, also known as Platelet endothelial cell adhesion molecule or PECAM-1. CD31 is a ~130kDa single pass type-1 membrane glycoprotein bearing six C2 Ig-like domains, expressed on all continuous endothelium including arteries, veins and non-sinusoidal capillaries. CD31 is also expressed on all haemopoietic lineages with the exception of the erythroid line (Bogen et al. 1992).</p> <p>Hamster anti mouse CD31, clone 2H8 effectively inhibits transmigration of activated polymorphonuclear cells and monocytes across the endothelium. In a mouse model for acute peritonitis clone 2H8 blocks acute inflammation (Bogen et al. 1994). CD31 has also been shown to be critically involved in the sensing of changes in shear stress associated with atherosclerotic lesions and in the associated atherogenesis (Stevens et al. 2008).</p> |
|--------------------|--|

| | |
|-----------------------|--|
| Flow Cytometry | Use 5ul of the suggested working dilution to label 100ul of whole blood. |
|-----------------------|--|

| | |
|-------------------|---|
| References | <ol style="list-style-type: none">1. Bogen, S.A. <i>et al.</i> (1992) Association of murine CD31 with transmigrating lymphocytes following antigenic stimulation. Am J Pathol. 141 (4): 843-54.2. Xie, Y. & Muller, W.A. (1993) Molecular cloning and adhesive properties of murine platelet/endothelial cell adhesion molecule 1. Proc Natl Acad Sci U S A. 90 (12): 5569-73.3. Bogen, S. <i>et al.</i> (1994) Monoclonal antibody to murine PECAM-1 (CD31) blocks acute inflammation <i>in vivo</i>. J Exp Med. 179 (3): 1059-64.4. Bixel, M.G. <i>et al.</i> (2010) CD99 and CD99L2 act at the same site as, but independently of, PECAM-1 during leukocyte diapedesis. Blood. 116: 1172-84.5. Ishikawa, J. <i>et al.</i> (2002) Use of anti-platelet-endothelial cell adhesion molecule-1 antibody in the control of disease progression in established collagen-induced arthritis in DBA/1J mice. Jpn J Pharmacol. 88: 332-40.6. Thurston, G. <i>et al</i> (2005) Angiopoietin 1 causes vessel enlargement, without angiogenic sprouting, during a critical developmental period Development. 132: 3317-26. |
|-------------------|---|

7. Lonsdorf, A.S. *et al.* (2012) Engagement of $\alpha\text{IIb}\beta\text{3}$ (GPIIb/IIIa) with $\alpha\text{v}\beta\text{3}$ Integrin Mediates Interaction of Melanoma Cells with Platelets: A CONNECTION TO HEMATOGENOUS METASTASIS. [J Biol Chem. 287: 2168-78.](#)
8. Rijcken, E. *et al.* (2007) PECAM-1 (CD 31) mediates transendothelial leukocyte migration in experimental colitis. [Am J Physiol Gastrointest Liver Physiol. 293: G446-52.](#)
9. Vielhauer, V. *et al.* (2005) Renal cell-expressed TNF receptor 2, not receptor 1, is essential for the development of glomerulonephritis. [J Clin Invest. 115: 1199-209.](#)
10. Brackett, C.M. *et al.* (2013) IL-17 promotes neutrophil entry into tumor-draining lymph nodes following induction of sterile inflammation. [J Immunol. 191 \(8\): 4348-57.](#)
11. Bixel, M.G. *et al.* (2010) CD99 and CD99L2 act at the same site as, but independently of, PECAM-1 during leukocyte diapedesis. [Blood. 116: 1172-84.](#)
12. Wu, Y. *et al.* (2010) Therapeutic antibody targeting of individual Notch receptors. [Nature. 464: 1052-7.](#)
13. Wilhelm, A. *et al.* (2015) CD248/Endosialin critically regulates hepatic stellate cell proliferation during chronic liver injury via a PDGF-regulated mechanism. [Gut. pii: gutjnl-2014-308325.](#)
14. Smith, S.W. *et al.* (2015) Genetic Deletion of the Stromal Cell Marker CD248 (Endosialin) Protects against the Development of Renal Fibrosis. [Nephron. 131 \(4\): 265-77.](#)

Storage Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10162 available at: 10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight@800](#), [FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

 Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

Printed on 17 Sep 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)