

Datasheet: MCA1371Z

Description:	HAMSTER ANTI MOUSE CD54:Preservative Free
Specificity:	CD54
Other names:	ICAM-1
Format:	Preservative Free
Product Type:	Monoclonal Antibody
Clone:	3E2B
Isotype:	IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			neat
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting			▪	
Functional Assays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present. Sterile filtered.

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	<p>UniProt: P13597 Related reagents</p> <p>Entrez Gene: 15894 Icam1 Related reagents</p>
Synonyms	Icam-1
RRID	AB_2121770
Specificity	<p>Hamster anti Mouse CD54 antibody, clone 3E2B recognizes murine Intercellular adhesion molecule 1, also known as ICAM-1, CD54, MALA-2 or MyD10. CD54 is a 537 amino acid ~89kDa single pass type I transmembrane glycoprotein bearing five Ig-like C2 type domains. Two isoforms, differing at the N-terminal signal peptide sequence and flanking end terminal domain of the processed molecule are generated by alternative splicing (UniProt P13577). CD45 is a counter-receptor for LFA-1 (Diamond et al. 1990).</p> <p>Murine CD54 is expressed at low levels on T-cells, B-cells, monocytes and endothelial cells but can be induced strongly by a number of stimuli, including UV radiation (Tang and Udey 1991).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Scheynius, A. <i>et al.</i> (1993) Reduced contact sensitivity reactions in mice treated with monoclonal antibodies to leukocyte function-associated molecule-1 and intercellular adhesion molecule-1. J Immunol. 150 (2): 655-63. Fabry, Z. <i>et al.</i> (1992) Adhesion molecules on murine brain microvascular endothelial cells: expression and regulation of ICAM-1 and Lgp 55. J Neuroimmunol. 36 (1): 1-11. Greter, M. <i>et al.</i> (2009) Neo-lymphoid aggregates in the adult liver can initiate potent cell-mediated immunity. PLoS Biol. 7 (5): e1000109. Brochard, V. <i>et al.</i> (2009) Infiltration of CD4+ lymphocytes into the brain contributes to neurodegeneration in a mouse model of Parkinson disease. J Clin Invest. 119: 182-92. Stein, S. <i>et al.</i> (2010) SIRT1 reduces endothelial activation without affecting vascular function in ApoE^{-/-} mice. Aging (Albany NY). 2: 353-60. Impellizzeri, D. <i>et al.</i> (2015) The anti-inflammatory and antioxidant effects of bergamot juice extract (BJe) in an experimental model of inflammatory bowel disease. Clin Nutr. 34 (6): 1146-54. Cherry, J.D. <i>et al.</i> (2013) Thermal injury lowers the threshold for radiation-induced neuroinflammation and cognitive dysfunction. Radiat Res. 180 (4): 398-406. Fusco, R. <i>et al.</i> (2017) A flavonoid-rich extract of orange juice reduced oxidative stress in an experimental model of inflammatory bowel disease J Funct Foods. 30: 168-78. Xuemei, H. <i>et al.</i> (2020) Mesenchymal stem cell-derived exosomal microRNA-34c-5p ameliorates renal pericyte activation by inhibiting core fucosylation[Preprint]. Nov 13. [Epub ahead of print].

Further Reading 1. Johnston, S.C. *et al.* (1990) On the species specificity of the interaction of LFA-1 with intercellular adhesion molecules. [J Immunol. 145 \(4\): 1181-7.](#)

Storage Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee Guaranteed until date of expiry. Please see product label.

Health And Safety Information Material Safety Datasheet documentation #10162 available at: 10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight®800](#), [FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

Recommended Negative Controls

[HAMSTER \(ARMENIAN\) IgG NEGATIVE CONTROL \(MCA2356\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M354361:190704'

Printed on 11 Jul 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)