

## Datasheet: MCA1379

**BATCH NUMBER 160549**

<b>Description:</b>	MOUSE ANTI HUMAN CD24
<b>Specificity:</b>	CD24
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	SN3
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Chimpanzee</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.</p>

<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Glycoproteins purified from human NALM-1 cell line
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P25063</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">100133941</a>    CD24    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	CD24A
<b>RRID</b>	AB_321526
<b>Fusion Partners</b>	Spleen cells from immunized BALB/c mice were fused with cells of a mouse myeloma cell line
<b>Specificity</b>	<b>Mouse anti Human CD24 antibody, clone SN3</b> recognizes the human CD24 cell surface antigen, a heavily glycosylated molecule of ~35-45 kDa. CD24 is expressed by granulocytes and B lymphocytes, and by some activated T cells and T cell malignancies.
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood
<b>Histology Positive Control Tissue</b>	Tonsil
<b>References</b>	<ol style="list-style-type: none"> <li>1. Barcos, M. <i>et al.</i> (1986) Follicular mantle zone cell subpopulations detected by monoclonal antibody SN3. <a href="#">Hematol Oncol. 4 (4): 251-9.</a></li> <li>2. Fukukawa, T. <i>et al.</i> (1986) New monoclonal antibodies SN3, SN3a, and SN3b directed to sialic acid of glycoprotein on human non-T leukemia cells. <a href="#">Exp Hematol. 14 (9): 850-5.</a></li> <li>3. Sallustio, F. <i>et al.</i> (2010) TLR2 plays a role in the activation of human resident renal stem/progenitor cells. <a href="#">FASEB J. 24 (2): 514-25.</a></li> <li>4. Ohyama, M. <i>et al.</i> (2006) Characterization and isolation of stem cell-enriched human hair follicle bulge cells. <a href="#">J Clin Invest. 116:249-60.</a></li> <li>5. Kanderová, V. <i>et al.</i> (2010) Aberrantly expressed CEACAM6 is involved in the signaling</li> </ol>

leading to apoptosis of acute lymphoblastic leukemia cells. [Exp Hematol. 38: 653-660.e1.](#)

6. Kap, Y.S. *et al.* (2009) A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. [J Histochem Cytochem. 57 \(12\): 1159-67.](#)

7. Franz, C. *et al.* (2016) Extracellular vesicles in human follicular fluid do not promote coagulation. [Reprod Biomed Online. Aug 22. pii: S1472-6483\(16\)30453-9. \[Epub ahead of print\]](#)

<b>Storage</b>	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
----------------	---

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1379">https://www.bio-rad-antibodies.com/SDS/MCA1379</a> 10040
--------------------------------------	---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M383442:210513'

Printed on 18 Jul 2024

