

Datasheet: MCA1379F BATCH NUMBER 159429

| Description: | MOUSE ANTI HUMAN CD24:FITC |
|---------------|----------------------------|
| Specificity: | CD24 |
| Format: | FITC |
| Product Type: | Monoclonal Antibody |
| Clone: | SN3 |
| Isotype: | lgG1 |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | • | | | Neat - 1/2 |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| Target Species | Human | | | |
|-----------------------------|--|-----------------------------|---------------------|---------------------|
| Species Cross Reactivity | Reacts with: Chimpanzee N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. | | | |
| Product Form | Purified IgG conju | ugated to Fluorescein Isotl | niocyanate Isomer 1 | l (FITC) - liquid |
| Max Ex/Em | Fluorophore FITC | Excitation Max (nm) 490 | Emission Max (nm | <u>)</u> |
| Preparation | Purified IgG preparent | ared by affinity chromatog | raphy on Protein A | from tissue culture |
| Buffer Solution | Phosphate buffer | ed saline | | |

| Preservative Stabilisers | 0.09% Sodium Azide 1% Bovine Serum Albumin | |
|--------------------------------|--|--|
| Approx. Protein Concentrations | IgG concentration 0.1 mg/ml | |
| Immunogen | Glycoproteins purified from human NALM-1 cell line | |
| External Database Links | UniProt: P25063 Related reagents Entrez Gene: 100133941 CD24 Related reagents | |
| Synonyms | CD24A | |
| RRID | AB_321527 | |
| Fusion Partners | Spleen cells from immunized BALB/c mice were fuzed with cells of a mouse myeloma cell line | |
| Specificity | Mouse anti Human CD24 antibody, clone SN3 recognizes the human CD24 cell surface antigen, a heavily glycosylated molecule of ~35-45 kDa. CD24 is expressed by granulocytes and B lymphocytes, and by some activated T cells and T cell malignancies. | |
| Flow Cytometry | Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood | |
| References | 1. Barcos, M. <i>et al.</i> (1986) Follicular mantle zone cell subpopulations detected by monoclonal antibody SN3. Hematol Oncol. 4 (4): 251-9. 2. Fukukawa, T. <i>et al.</i> (1986) New monoclonal antibodies SN3, SN3a, and SN3b directed to sialic acid of glycoprotein on human non-T leukemia cells. Exp Hematol. 14 (9): 850-5. 3. Sallustio, F. <i>et al.</i> (2010) TLR2 plays a role in the activation of human resident renal stem/progenitor cells. FASEB J. 24 (2): 514-25. 4. Ohyama, M. <i>et al.</i> (2006) Characterization and isolation of stem cell-enriched human hair follicle bulge cells. J Clin Invest. 116:249-60. 5. Kanderová, V. <i>et al.</i> (2010) Aberrantly expressed CEACAM6 is involved in the signaling leading to apoptosis of acute lymphoblastic leukemia cells. Exp Hematol. 38: 653-660.e1. 6. Kap, Y.S. <i>et al.</i> (2009) A monoclonal antibody selection for immunohistochemical examination of lymphoid tissues from non-human primates. J Histochem Cytochem. 57 (12): 1159-67. 7. Franz, C. <i>et al.</i> (2016) Extracellular vesicles in human follicular fluid do not promote coagulation. Reprod Biomed Online. Aug 22. pii: S1472-6483(16)30453-9. [Epub ahead of print] | |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. | |

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

| Guarantee | 12 months from date of despatch |
|----------------------------------|--|
| Health And Safety Information | Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1379F 10041 |
| Regulatory | For research purposes only |

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M391395:211013'

Printed on 09 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint