

Datasheet: MCA1388

Description:	RAT ANTI MOUSE CD45
Specificity:	CD45
Other names:	LCA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	IBL-3/16
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen (1)	▪			1/50 - 1/100
Immunohistology - Paraffin		▪		
Immunohistology - Resin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting		▪		
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Mouse
Species Cross Reactivity	Does not react with:Rat, Human, Chicken
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline.
Preservative	<0.1% sodium azide (NaN ₃)
Stabilisers	1% bovine serum albumin
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml
Immunogen	Purified B cells from mouse lymph nodes.
External Database Links	<p>UniProt: P06800 Related reagents</p> <p>Entrez Gene: 19264 Ptprc Related reagents</p>
Synonyms	Ly-5
RRID	AB_321729
Fusion Partners	Spleen cells from an immunized Wistar rat were fused with cells of the SP2/0-Ag14 mouse myeloma cell line.
Specificity	<p>Rat anti Mouse CD45 antibody, clone IBL-3/16 recognizes murine Receptor-type tyrosine-protein phosphatase C, also known as CD45, Leukocyte common antigen, T200 or Lymphocyte antigen 5 (Ly5). CD45 is a 1291 amino acid ~175kDa single pass type I transmembrane glycoprotein belonging to the protein -tyrosing phosphatase family. CD45 has two fibronectin type-III domains and two tyrosine protein phosphatase domains (UniProt:: P06800).</p> <p>Multiple isoforms are generated by alternative splicing with isoforms having differing deletions in the N-terminal region (Saga et al. 1987). Rat anti Mouse CD45 antibody, clone IBL-3/16 is expected to recognize all isoforms of murine CD45.</p> <p>Rat anti Mouse CD45 antibody, clone IBL-3/16 has been used successfully for the identification of CD45 in murine samples using Immunohistochemical (both cryo and FFPE), immunofluorescence and western blotting techniques (Kondo et al. 2011, Cuadros et al. 2006).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Jiang, H.R. <i>et al.</i> (2001) Total dose and frequency of administration critically affect success of nasal mucosal tolerance induction. Br J Ophthalmol. 85: 739-44. Richards, J.G. <i>et al.</i> (2003) PS2APP transgenic mice, coexpressing hPS2mut and hAPPswe, show age-related cognitive deficits associated with discrete brain amyloid

- deposition and inflammation. [J Neurosci. 23: 8989-9003.](#)
3. Yan, Q. *et al.* (2003) Anti-inflammatory drug therapy alters beta-amyloid processing and deposition in an animal model of Alzheimer's disease. [J Neurosci. 23: 7504-9.](#)
 4. Paris, D. *et al.* (2010) Reduction of beta-amyloid pathology by celastrol in a transgenic mouse model of Alzheimer's disease. [J Neuroinflammation. 7: 17.](#)
 5. Laporte, V. *et al.* (2006) CD40 deficiency mitigates Alzheimer's disease pathology in transgenic mouse models. [J Neuroinflammation. 3: 3.](#)
 6. Cuadros, M.A. *et al.* (2006) Specific immunolabeling of brain macrophages and microglial cells in the developing and mature chick central nervous system. [J Histochem Cytochem. 54: 727-38.](#)
 7. Neutzner, M. *et al.* (2007) MFG-E8/lactadherin promotes tumor growth in an angiogenesis-dependent transgenic mouse model of multistage carcinogenesis. [Cancer Res. 67: 6777-85.](#)
 8. Zehntner, S.P. *et al.* (2007) X-linked inhibitor of apoptosis regulates T cell effector function. [J Immunol. 2007 Dec 1;179\(11\):7553-60.](#)
 9. Boger, H.A. *et al.* (2007) Long-term consequences of methamphetamine exposure in young adults are exacerbated in glial cell line-derived neurotrophic factor heterozygous mice. [J Neurosci. 27: 8816-25.](#)
 10. Santos, A.M. *et al.* (2008) Embryonic and postnatal development of microglial cells in the mouse retina. [J Comp Neurol. 506: 224-39.](#)
 11. Clausen, B.H. *et al.* (2008) Interleukin-1beta and tumor necrosis factor-alpha are expressed by different subsets of microglia and macrophages after ischemic stroke in mice. [J Neuroinflammation. 5: 46.](#)
 12. Li, L. *et al.* (2008) GRK5 deficiency exaggerates inflammatory changes in TgAPPsw mice. [J Neuroinflammation. 5: 24.](#)
 13. Santos, A.M. *et al.* (2010) Microglial response to light-induced photoreceptor degeneration in the mouse retina. [J Comp Neurol. 518: 477-92.](#)
 14. Boger, H.A. *et al.* (2009) Minocycline restores striatal tyrosine hydroxylase in GDNF heterozygous mice but not in methamphetamine-treated mice. [Neurobiol Dis. 33: 459-66.](#)
 15. Rahman, A. *et al.* (2011) Chronic colitis induces expression of β -defensins in murine intestinal epithelial cells. [Clin Exp Immunol. 163: 123-30.](#)
 16. Peng, Y. *et al.* (2010) L-3-n-butylphthalide improves cognitive impairment and reduces amyloid-beta in a transgenic model of Alzheimer's disease. [J Neurosci. 30: 8180-9.](#)
 17. Ricciardelli, C. *et al.* (2011) The ADAMTS1 protease gene is required for mammary tumor growth and metastasis. [Am J Pathol. 179: 3075-85.](#)
 18. Medeiros, R. *et al.* (2011) Loss of muscarinic M1 receptor exacerbates Alzheimer's disease-like pathology and cognitive decline. [Am J Pathol. 179: 980-91.](#)
 19. Kondo, Y. *et al.* (2011) Macrophages counteract demyelination in a mouse model of globoid cell leukodystrophy. [J Neurosci. 31: 3610-24.](#)
 20. Wang, P. *et al.* (2012) IL-22 signaling contributes to West Nile encephalitis pathogenesis. [PLoS One. 7: e44153.](#)
 21. Klose, A. *et al.* (2013) Monocyte/macrophage MMP-14 modulates cell infiltration and T-cell attraction in contact dermatitis but not in murine wound healing. [Am J Pathol. 182: 755-64.](#)
 22. Passos, G.F. *et al.* (2013) The bradykinin B1 receptor regulates A β deposition and neuroinflammation in Tg-SwDI mice. [Am J Pathol. 182: 1740-9.](#)
 23. Ferrer-Martín, R.M. *et al.* (2015) Microglial Activation Promotes Cell Survival in

- Organotypic Cultures of Postnatal Mouse Retinal Explants. [PLoS One. 10 \(8\): e0135238.](#)
24. Tzekov, R. *et al.* (2016) Sub-Chronic Neuropathological and Biochemical Changes in Mouse Visual System after Repetitive Mild Traumatic Brain Injury. [PLoS One. 11 \(4\): e0153608.](#)
25. Da Ros, F. *et al.* (2017) Targeting Interleukin-1 β Protects from Aortic Aneurysms Induced by Disrupted Transforming Growth Factor β Signaling. [Immunity. 47 \(5\): 959-973.e9.](#)
26. Witcher, K.G. *et al.* (2018) Traumatic brain injury-induced neuronal damage in the somatosensory cortex causes formation of rod-shaped microglia that promote astrogliosis and persistent neuroinflammation. [Glia. 66 \(12\): 2719-36.](#)
27. Singh, T. *et al.* (2019) Loss of MafA and MafB expression promotes islet inflammation. [Sci Rep. 9 \(1\): 9074.](#)
28. Boza-Serrano, A. *et al.* (2019) Galectin-3, a novel endogenous TREM2 ligand, detrimentally regulates inflammatory response in Alzheimer's disease. [Acta Neuropathol. 138 \(2\): 251-73.](#)
29. Liu, X. *et al.* (2019) Cell-Type-Specific Interleukin 1 Receptor 1 Signaling in the Brain Regulates Distinct Neuroimmune Activities. [Immunity. 50 \(2\): 317-33.e6.](#)
30. Tóth, K. *et al.* (2022) The NKCC1 ion transporter modulates microglial phenotype and inflammatory response to brain injury in a cell-autonomous manner. [PLoS Biol. 20 \(1\): e3001526.](#)
31. Cheng-Hathaway, P.J. *et al.* (2018) The Trem2 R47H variant confers loss-of-function-like phenotypes in Alzheimer's disease [Molecular Neurodegeneration. 13 \(1\), 29.](#)
32. Jürgenson, M. *et al.* (2019) Effects of the drug combination memantine and melatonin on impaired memory and brain neuronal deficits in an amyloid-predominant mouse model of Alzheimer's disease. [J Pharm Pharmacol. 71 \(11\): 1695-705.](#)
33. Joly-Amado, A. *et al.* (2020) CCL2 Overexpression in the Brain Promotes Glial Activation and Accelerates Tau Pathology in a Mouse Model of Tauopathy. [Front Immunol. 11: 997.](#)
34. Li, L. *et al.* (2021) Systematic review and meta-analysis of ureteral stent for risk factors of restenosis after laparoscopic pyeloplasty. [Ann Palliat Med. 10 \(10\): 10527-34.](#)
35. Takemura, S. *et al.* (2023) Juvenile social isolation affects the structure of the tanycyte-vascular interface in the hypophyseal portal system of the adult mice. [Neurochem Int. 162: 105439.](#)

Storage	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
Guarantee	12 months from date of despatch
Health And Safety Information	<p>Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1388</p> <p>10041</p>
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR72...)	HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M433687:241119'

Printed on 19 Nov 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)