

## Datasheet: MCA1396D550 BATCH NUMBER 165781

Description:	MOUSE ANTI HISTIDINE TAG:DyLight®550
Specificity:	HISTIDINE TAG
Format:	DyLight®550
Product Type:	Monoclonal Antibody
Clone:	AD1.1.10
Isotype:	lgG1
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further							
	information. For gener	information. For general protocol recommendations, please visit <u>www.bio-</u>						
	rad-antibodies.com/pr	otocols.		. –				
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	-						
	Western Blotting				1/1,000 - 1/10,000			
	Immunofluorescence	-			1/50 - 1/200			
	Where this product has not been tested for use in a particular technique this does not							
	necessarily exclude its	s use in such	n procedu	res. Suggested workin	g dilutions are given as			
	a quide only. It is reco	mmended th	at the us	er titrates the product f	for use in their own			
	system using appropri							
Target Species	Synthetic Peptide							
Product Form	Purified IgG conjugated to DyLight® 550 - liquid							
Max Ex/Em	Fluorophore	Excitation I	Max (nm)	Emission Max (nm)				
	Dylight®550	562	2	576				
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant				n tissue culture			
Preservative Stabilisers	0.09% sodium azide (NaN <sub>3</sub> )							
Approx. Protein Concentrations	IgG concentration 1.0	mg/ml						

Immunogen	PAX6 transcription factor linked to histidine tag.			
Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells of the mouse NS1 myeloma cell line.			
Specificity	<b>Mouse anti Histidine tag antibody, clone AD1.1.10</b> recognizes proteins and peptides containing the motif H-H-H-H-H and is therefore of value in detecting proteins containing histidine tags. Clone AD1.1.10 has been used to detect and purify histidine-tagged proteins expressed in mammalian (Hoffmann <i>et al.</i> 2007) and Hwang <i>et al.</i> 2008) and non-mammalian (Zheng <i>et al.</i> 2007; Gunnarsen <i>et al.</i> 2010; and <u>de Vooght <i>et al.</i></u> 2012) cell lines.			
	In Western blotting of bacterial extracts the antibody has been shown not to cross-react with any endogenous products, although some cross-reactivity may be seen with extracts of insect or mammalian cells.			
	This antibody is routinely tested in Western blotting on histidine tagged recombinant proteins and reacts against all histidine-tagged proteins so far tested.			
References	<ol> <li>Els Conrath, K. <i>et al.</i> (2001) Camel single-domain antibodies as modular building units in bispecific and bivalent antibody constructs. J Biol Chem. 276 (10): 7346-50.</li> <li>Suen, J.L. <i>et al.</i> (2001) Characterization of self-T-cell response and antigenic determinant of U1A protein with bone marrow-derived dendritic cells in NZB x NZW F1 mice. Immunol. 103: 301-309.</li> <li>Hoffmann, S.C. <i>et al.</i> (2007) Identification of CLEC12B, an inhibitory receptor on myeloid cells. J Biol Chem. 282 (31): 22370-5.</li> <li>Zheng, J. <i>et al.</i> (2007) Serum from mice immunized in the context of Treg inhibition identifies DEK as a neuroblastoma tumor antigen. BMC Immunol. 8: 4.</li> <li>Bahi, A. &amp; Dreyer, J.L. (2008) Overexpression of plasminogen activators in the nucleus accumbens enhances cocaine-, amphetamine- and morphine-induced reward and behavioral sensitization. Genes Brain Behav. 7 (2): 244-56.</li> <li>Wrighton, K.H. <i>et al.</i> (2001) Transforming Growth Factor {beta} Can Stimulate Smad1 Phosphorylation Independently of Bone Morphogenic Protein Receptors. J Biol Chem. 284 (15): 9755-63.</li> <li>Diefenbacher, M. <i>et al.</i> (2011) The Dsl1 Tethering Complex Actively Participates in Soluble NSF (N-Ethylmaleimide-sensitive Factor) Attachment Protein Receptor (SNARE) Complex Assembly at the Endoplasmic Reticulum in Saccharomyces cerevisiae. J Biol Chem. 286: 25027-38.</li> <li>Alvarez, M.M. <i>et al.</i> (2010) Specific recognition of influenza A/H1N1/2009 antibodies in human serum: a simple virus-free ELISA method. PLoS One. 5: e10176.</li> <li>Bahi, A. <i>et al.</i> (2008) The role of tissue-type plasminogen activator system in amphetamine-induced conditional place preference extinction and reinstatement. Neuropsychopharmacology. 33: 2726-34.</li> <li>Gunnarsen, K.S. <i>et al.</i> (2010) Periplasmic expression of soluble single chain T cell receptors is rescued by the chaperone FkpA. <u>BMC Biotechnol. 10: 8.</u></li> <li>Hwang, H.Y. <i>et al.</i> (2008) Highly specific inhibition of C1q gl</li></ol>			

12. De Vooght, L. *et al.* (2012) Expression and extracellular release of a functional anti-trypanosome Nanobody® in Sodalis glossinidius, a bacterial symbiont of the tsetse fly. <u>Microb Cell Fact. 11: 23.</u>

13. Saerens, D. *et al.* (2004) Single domain antibodies derived from dromedary lymph node and peripheral blood lymphocytes sensing conformational variants of prostate-specific antigen. <u>J Biol Chem. 279 (50): 51965-72.</u>

14. Than, N.G. *et al.* (2014) Evolutionary origins of the placental expression of chromosome 19 cluster galectins and their complex dysregulation in preeclampsia. <u>Placenta. 35 (11): 855-65.</u>

 15. Elders RC *et al.* (2014) Recombinant canine IgE Fc and an IgE Fc-TRAIL fusion protein bind to neoplastic canine mast cells. <u>Vet Immunol Immunopathol. 159 (1-2): 29-40.</u>
 16. Chin, S.E. *et al.* (2015) Isolation of high-affinity, neutralizing anti-idiotype antibodies by phage and ribosome display for application in immunogenicity and pharmacokinetic analyses. <u>J Immunol Methods. 416: 49-58.</u>

17. Peyrassol, X. *et al.* (2016) Development by Genetic Immunization of Monovalent Antibodies (Nanobodies) Behaving as Antagonists of the Human ChemR23 Receptor. J Immunol. 196 (6): 2893-901.

18. Kim H & Loparo JJ (2016) Multistep assembly of DNA condensation clusters by SMC. <u>Nat Commun. 7: 10200.</u>

Borg M *et al.* (2014) A novel interaction between Rab7b and actomyosin reveals a dual role in intracellular transport and cell migration. <u>J Cell Sci. 127 (Pt 22): 4927-39.</u>
 De Meyer, T. *et al.* (2015) Comparison of VHH-Fc antibody production in Arabidopsis

thaliana, Nicotiana benthamiana and Pichia pastoris. <u>Plant Biotechnol J. 13 (7): 938-47.</u>

21. Siddiqui AA *et al.* (2015) Humoral immune responses to a recombinant Plasmodium vivax tryptophan-rich antigen among *Plasmodium vivax*-infected patients and its localization in the parasite. <u>Appl Biochem Biotechnol. 175 (4): 2166-77.</u>

22. Warnecke, A. *et al.* (2017) Nitration of MOG diminishes its encephalitogenicity depending on MHC haplotype. <u>J Neuroimmunol. 303: 1-12.</u>

23. Bertucci, A. *et al.* (2011) A new coral carbonic anhydrase in *Stylophora pistillata*. <u>Mar</u> <u>Biotechnol (NY). 13 (5): 992-1002.</u>

24. Boujon, C.L. *et al.* (2017) Development and validation of an immunohistochemistry procedure for the detection of a neurotropic bovine astrovirus. <u>J Virol Methods. 239:</u> <u>26-33.</u>

25. Cartwright, S.P. *et al.* (2017) Rapid expression and purification of the hepatitis delta virus antigen using the methylotropic yeast *Pichia pastoris*. <u>BMC Res Notes. 10 (1): 340.</u>
 26. Thanongsaksrikul, J. *et al.* (2018) Identification and production of mouse scFv to specific epitope of enterovirus-71 virion protein-2 (VP2). <u>Arch Virol. 163 (5): 1141-1152.</u>
 27. Gunnarsen, K.S. *et al.* (2018) Soluble T-cell receptor design influences functional yield in an E. coli chaperone-assisted expression system. <u>PLoS One. 13 (4): e0195868.</u>
 28. Ascione, A. *et al.* (2019) Development of a novel human phage display-derived anti-LAG3 scFv antibody targeting CD8<sup>+</sup> T lymphocyte exhaustion. <u>BMC Biotechnol. 19 (1): 67.</u>

29. Zoccola, D. *et al.* (2017) Structural and functional analysis of coral Hypoxia Inducible Factor. <u>PLoS One. 12 (11): e0186262.</u>

30. Kimura, K. *et al.* (2021) Overexpression of human BAG3<sup>P209L</sup> in mice causes restrictive cardiomyopathy. <u>Nat Commun. 12 (1): 3575</u>.

31. Dongdem, J.T. et al. (2021) Modification of small ubiquitin-related modifier 2 (SUMO2)

	<ul> <li>by phosphoubiquitin in HEK293T cells. <u>Proteomics. 21 (15): e2000234.</u></li> <li>32. Chuang, H.C. <i>et al.</i> (2021) Effect of cell-permeable grouper Manganese Superoxide Dismutase on environmental stress in fish. <u>Protein Expr Purif. 187: 105951.</u></li> <li>33. Cheng, C.M. <i>et al.</i> (2021) Heterologous expression of bacterial CotA-laccase, characterization and its application for biodegradation of malachite green. <u>Bioresour Technol. 340: 125708.</u></li> <li>34. De Vooght, L. <i>et al.</i> (2022) Targeting the tsetse-trypanosome interplay using genetically engineered <i>Sodalis glossinidius.</i>. <u>PLoS Pathog. 18 (3): e1010376.</u></li> <li>35. Minami, S.A. <i>et al.</i> (2022) Production of novel SARS-CoV-2 Spike truncations in Chinese hamster ovary cells leads to high expression and binding to antibodies. <u>Biotechnol J. 17 (9): e2100678.</u></li> <li>36. Chen, Y.J. <i>et al.</i> (2023) A non-genetic engineering platform for rapidly generating and expanding cancer-specific armed T cells. <u>J Biomed Sci. 30 (1): 35.</u></li> <li>37. Boudkkazi, S. <i>et al.</i> (2023) Heterologous expression and characterization of a MoAA16 polysaccharide monoxygenase from the rice blast fungus Magnaporthe oryzae <u>Electronic Journal of Biotechnology. 66: 1-16.</u></li> <li>39. Rossey, I. <i>et al.</i> (2021) A vulnerable, membrane-proximal site in human respiratory syncytial virus F revealed by a prefusion-specific single-domain antibody. J Virol. 95 (11): <u>e02279-20.</u></li> <li>40. Khosravi, M. <i>et al.</i> (2024) Shiga toxin type 2 B subunit protects mice against toxin challenge when leashed and bundled by a stable pentameric coiled-coil molecule. <u>Vaccine. Feb 15 S0264-410X(24)00129-4. [Epub ahead of print].</u></li> <li>42. Kimura, T. <i>et al.</i> (2024) Quantification of lipoprotein lipase in mouse plasma with a sandwich enzyme-linked immunosorbent assay. <u>J Lipid Res. 65 (4): 100532.</u></li> </ul>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Acknowledgements	DyLight is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. His-tag is a registered trademark of EMD Biosciences.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1396D550 10040
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-ra	Worldwide ad.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-ra	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-ra	To <sup>d.</sup> ფეუკე	
batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419191:230606'							

## Printed on 15 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint