

Datasheet: MCA1396D550

Description:	MOUSE ANTI HISTIDINE TAG:DyLight®550
Specificity:	HISTIDINE TAG
Format:	DyLight®550
Product Type:	Monoclonal Antibody
Clone:	AD1.1.10
Isotype:	lgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Western Blotting	•			1/1,000 - 1/10,000
Immunofluorescence	•			1/50 - 1/200

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Synthetic Peptide			
Product Form	Purified IgG conju			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nn	n)
	Dylight®550	562	576	
Preparation	Purified IgG preparent	ared by affinity chromatog	raphy on Protein A	from tissue culture
Preservative Stabilisers	0.09% sodium az			
Approx. Protein Concentrations	IgG concentration	n 1.0 mg/ml		
Immunogen	PAX6 transcriptio	n factor linked to histidine	tag.	

Fusion Partners

Spleen cells from immunized Balb/c mice were fused with cells of the mouse NS1 myeloma cell line.

Specificity

Mouse anti Histidine tag antibody, clone AD1.1.10 recognizes proteins and peptides containing the motif H-H-H-H-H and is therefore of value in detecting proteins containing histidine tags. Clone AD1.1.10 has been used to detect and purify histidine-tagged proteins expressed in mammalian (Hoffmann *et al.* 2007) and Hwang *et al.* 2008) and non-mammalian (Zheng *et al.* 2007; Gunnarsen *et al.* 2010; and de Vooght *et al.* 2012) cell lines.

In Western blotting of bacterial extracts the antibody has been shown not to cross-react with any endogenous products, although some cross-reactivity may be seen with extracts of insect or mammalian cells.

This antibody is routinely tested in Western blotting on histidine tagged recombinant proteins and reacts against all histidine-tagged proteins so far tested.

References

- 1. Els Conrath, K. *et al.* (2001) Camel single-domain antibodies as modular building units in bispecific and bivalent antibody constructs. J Biol Chem. 276 (10): 7346-50.
- 2. Suen, J.L. *et al.* (2001) Characterization of self-T-cell response and antigenic determinant of U1A protein with bone marrow-derived dendritic cells in NZB x NZW F₁ mice. Immunol. 103: 301-309.
- 3. Hoffmann, S.C. *et al.* (2007) Identification of CLEC12B, an inhibitory receptor on myeloid cells. <u>J Biol Chem. 282 (31): 22370-5.</u>
- 4. Zheng, J. *et al.* (2007) Serum from mice immunized in the context of Treg inhibition identifies DEK as a neuroblastoma tumor antigen. BMC Immunol. 8: 4.
- 5. Bahi, A. & Dreyer, J.L. (2008) Overexpression of plasminogen activators in the nucleus accumbens enhances cocaine-, amphetamine- and morphine-induced reward and behavioral sensitization. Genes Brain Behav. 7 (2): 244-56.
- 6. Wrighton, K.H. *et al.* (2009) Transforming Growth Factor {beta} Can Stimulate Smad1 Phosphorylation Independently of Bone Morphogenic Protein Receptors. <u>J Biol Chem. 284</u> (15): 9755-63.
- 7. Diefenbacher, M. *et al.* (2011) The Dsl1 Tethering Complex Actively Participates in Soluble NSF (N-Ethylmaleimide-sensitive Factor) Attachment Protein Receptor (SNARE) Complex Assembly at the Endoplasmic Reticulum in Saccharomyces cerevisiae. <u>J Biol</u> Chem. 286: 25027-38.
- 8. Alvarez, M.M. *et al.* (2010) Specific recognition of influenza A/H1N1/2009 antibodies in human serum: a simple virus-free ELISA method. <u>PLoS One. 5: e10176.</u>
- 9. Bahi, A. *et al.* (2008) The role of tissue-type plasminogen activator system in amphetamine-induced conditional place preference extinction and reinstatement. Neuropsychopharmacology. 33: 2726-34.
- 10. Gunnarsen, K.S. *et al.* (2010) Periplasmic expression of soluble single chain T cell receptors is rescued by the chaperone FkpA. <u>BMC Biotechnol. 10: 8.</u>
- 11. Hwang, H.Y. *et al.* (2008) Highly specific inhibition of C1q globular-head binding to human IgG: a novel approach to control and regulate the classical complement pathway using an engineered single chain antibody variable fragment. <u>Mol Immunol. 45: 2570-80.</u>
- 12. De Vooght, L. et al. (2012) Expression and extracellular release of a functional

- anti-trypanosome Nanobody® in Sodalis glossinidius, a bacterial symbiont of the tsetse fly. Microb Cell Fact. 11: 23.
- 13. Saerens, D. *et al.* (2004) Single domain antibodies derived from dromedary lymph node and peripheral blood lymphocytes sensing conformational variants of prostate-specific antigen. <u>J Biol Chem. 279 (50): 51965-72.</u>
- 14. Than, N.G. *et al.* (2014) Evolutionary origins of the placental expression of chromosome 19 cluster galectins and their complex dysregulation in preeclampsia. Placenta. 35 (11): 855-65.
- 15. Elders RC *et al.* (2014) Recombinant canine IgE Fc and an IgE Fc-TRAIL fusion protein bind to neoplastic canine mast cells. <u>Vet Immunol Immunopathol</u>. 159 (1-2): 29-40.
- 16. Chin, S.E. *et al.* (2015) Isolation of high-affinity, neutralizing anti-idiotype antibodies by phage and ribosome display for application in immunogenicity and pharmacokinetic analyses. J Immunol Methods. 416: 49-58.
- 17. Peyrassol, X. *et al.* (2016) Development by Genetic Immunization of Monovalent Antibodies (Nanobodies) Behaving as Antagonists of the Human ChemR23 Receptor. <u>J. Immunol.</u> 196 (6): 2893-901.
- 18. Kim H & Loparo JJ (2016) Multistep assembly of DNA condensation clusters by SMC. Nat Commun. 7: 10200.
- 19. Borg M *et al.* (2014) A novel interaction between Rab7b and actomyosin reveals a dual role in intracellular transport and cell migration. <u>J Cell Sci. 127 (Pt 22): 4927-39.</u>
- 20. De Meyer, T. *et al.* (2015) Comparison of VHH-Fc antibody production in Arabidopsis thaliana, Nicotiana benthamiana and Pichia pastoris. Plant Biotechnol J. 13 (7): 938-47.
- 21. Siddiqui AA *et al.* (2015) Humoral immune responses to a recombinant Plasmodium vivax tryptophan-rich antigen among *Plasmodium vivax*-infected patients and its localization in the parasite. Appl Biochem Biotechnol. 175 (4): 2166-77.
- 22. Warnecke, A. *et al.* (2017) Nitration of MOG diminishes its encephalitogenicity depending on MHC haplotype. J Neuroimmunol. 303: 1-12.
- 23. Bertucci, A. *et al.* (2011) A new coral carbonic anhydrase in *Stylophora pistillata*. Mar Biotechnol (NY). 13 (5): 992-1002.
- 24. Boujon, C.L. *et al.* (2017) Development and validation of an immunohistochemistry procedure for the detection of a neurotropic bovine astrovirus. <u>J Virol Methods. 239:</u> 26-33.
- 25. Cartwright, S.P. *et al.* (2017) Rapid expression and purification of the hepatitis delta virus antigen using the methylotropic yeast *Pichia pastoris*. BMC Res Notes. 10 (1): 340.
- 26. Thanongsaksrikul, J. *et al.* (2018) Identification and production of mouse scFv to specific epitope of enterovirus-71 virion protein-2 (VP2). <u>Arch Virol. 163 (5): 1141-1152</u>.
- 27. Gunnarsen, K.S. *et al.* (2018) Soluble T-cell receptor design influences functional yield in an E. coli chaperone-assisted expression system. <u>PLoS One. 13 (4): e0195868.</u>
- 28. Ascione, A. *et al.* (2019) Development of a novel human phage display-derived anti-LAG3 scFv antibody targeting CD8⁺ T lymphocyte exhaustion. <u>BMC Biotechnol. 19</u> (1): 67.
- 29. Zoccola, D. *et al.* (2017) Structural and functional analysis of coral Hypoxia Inducible Factor. PLoS One. 12 (11): e0186262.
- 30. Kimura, K. *et al.* (2021) Overexpression of human BAG3^{P209L} in mice causes restrictive cardiomyopathy. Nat Commun. 12 (1): 3575.
- 31. Dongdem, J.T. *et al.* (2021) Modification of small ubiquitin-related modifier 2 (SUMO2) by phosphoubiquitin in HEK293T cells. <u>Proteomics. 21 (15): e2000234.</u>

- 32. Chuang, H.C. *et al.* (2021) Effect of cell-permeable grouper Manganese Superoxide Dismutase on environmental stress in fish. Protein Expr Purif. 187: 105951.
- 33. Cheng, C.M. *et al.* (2021) Heterologous expression of bacterial CotA-laccase, characterization and its application for biodegradation of malachite green. <u>Bioresour Technol.</u> 340: 125708.
- 34. De Vooght, L. *et al.* (2022) Targeting the tsetse-trypanosome interplay using genetically engineered *Sodalis glossinidius*.. <u>PLoS Pathog. 18 (3): e1010376.</u>
- 35. Minami, S.A. *et al.* (2022) Production of novel SARS-CoV-2 Spike truncations in Chinese hamster ovary cells leads to high expression and binding to antibodies. Biotechnol J. 17 (9): e2100678.
- 36. Chen, Y.J. *et al.* (2023) A non-genetic engineering platform for rapidly generating and expanding cancer-specific armed T cells. <u>J Biomed Sci. 30 (1): 35.</u>
- 37. Boudkkazi, S. *et al.* (2023) A Noelin-organized extracellular network of proteins required for constitutive and context-dependent anchoring of AMPA-receptors. <u>Neuron. 111 (16): 2544-56.e9.</u>
- 38. Nguyen, H.M. *et al.* (2023) Heterologous expression and characterization of a MoAA16 polysaccharide monooxygenase from the rice blast fungus Magnaporthe oryzae <u>Electronic Journal of Biotechnology.</u> 66: 1-16.
- 39. Rossey, I. *et al.* (2021) A vulnerable, membrane-proximal site in human respiratory syncytial virus F revealed by a prefusion-specific single-domain antibody. <u>J Virol. 95 (11):</u> e02279-20.
- 40. Khosravi, M. *et al.* (2016) Canine Distemper Virus Fusion Activation: Critical Role of Residue E123 of CD150/SLAM. J Virol. 90 (3): 1622-37.
- 41. Tamaki, Y. *et al.* (2024) Shiga toxin type 2 B subunit protects mice against toxin challenge when leashed and bundled by a stable pentameric coiled-coil molecule. Vaccine. Feb 15 S0264-410X(24)00129-4. [Epub ahead of print].
- 42. Kimura, T. *et al.* (2024) Quantification of lipoprotein lipase in mouse plasma with a sandwich enzyme-linked immunosorbent assay. <u>J Lipid Res. 65 (4): 100532.</u>
- 43. Wu, M.C. *et al.* (2022) Recombinant suilysin of Streptococcus suis enhances the protective efficacy of an engineered Pasteurella multocida toxin protein. <u>Res Vet Sci. 151:</u> 175-183.
- 44. Ramou, I. *et al.* (2025) Expression and purification of an activated Orexin receptor 1-G-protein complex <u>Protein Expression and Purification.</u>: 106660.
- 45. Paiva, S.C. *et al.* (2025) Exploring the Structure and Nucleic Acid Interactions of the *Leishmania* sp. Telomerase Reverse Transcriptase N-Terminal Region. <u>Arch Biochem Biophys.</u>: 110289.
- 46. Hayes JWP, Edwards JC, Ramirez Valdez K *et al.* (0) Evaluation of the contribution of the minor envelope complex glycoprotein 3 to the porcine reproductive and respiratory syndrome virus 1 neutralizing antibody response [version 1]. <u>VeriXiv 2025, 2:139</u>

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch	
Acknowledgements	DyLight is a trademark of Thermo Fisher Scientific Inc. and its registered trademark of EMD Biosciences.	subsidiaries. His-tag is a
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1396D550	
Regulatory	For research purposes only	

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419191:230606'

Printed on 30 Jun 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint