

## Datasheet: MCA1396D800GA

<b>Description:</b>	MOUSE ANTI HISTIDINE TAG:DyLight®800
<b>Specificity:</b>	HISTIDINE TAG
<b>Format:</b>	DyLight®800
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	AD1.1.10
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

### Product Details

**RRID** AB\_11152596

**Applications** This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000 - 1/5000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls

**Target Species** Synthetic Peptide

**Product Form** Purified IgG conjugated to DyLight®800 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	DyLight®800	777	794

**Preparation** Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

**Buffer Solution** Phosphate buffered saline

**Preservative Stabilisers** 0.09% Sodium Azide (NaN<sub>3</sub>)

**Approx. Protein Concentrations** IgG concentration 1.0mg/ml

**Immunogen** PAX6 transcription factor linked to histidine tag.

**Fusion Partners** Spleen cells from immunised Balb/c mice were fused with cells of the mouse NS1 myeloma cell line.

**Specificity** **Mouse anti Histidine tag antibody, clone AD1.1.10**, recognizes proteins and peptides containing

the motif H-H-H-H-H-H and is therefore of value in detecting proteins containing histidine tags. Clone AD1.1.10 has been used to detect and purify histidine-tagged proteins expressed in mammalian ([Hoffmann et al. 2007](#)) and [Hwang et al. 2008](#)) and non-mammalian ([Zheng et al. 2007](#); [Gunnarsen et al. 2010](#); and [de Vooght et al. 2012](#)) cell lines.

In Western blotting of bacterial extracts the antibody has been shown not to cross-react with any endogenous products, although some cross-reactivity may be seen with extracts of insect or mammalian cells.

This antibody is routinely tested in Western blotting on histidine tagged recombinant proteins and reacts against all histidine-tagged proteins so far tested.

---

## References

1. Els Conrath, K. *et al.* (2001) Camel single-domain antibodies as modular building units in bispecific and bivalent antibody constructs. [J Biol Chem. 276 \(10\): 7346-50.](#)
2. Suen, J.L. *et al.* (2001) Characterization of self-T-cell response and antigenic determinant of U1A protein with bone marrow-derived dendritic cells in NZB x NZW F<sub>1</sub> mice. [Immunol. 103: 301-309.](#)
3. Hoffmann, S.C. *et al.* (2007) Identification of CLEC12B, an inhibitory receptor on myeloid cells. [J Biol Chem. 282 \(31\): 22370-5.](#)
4. Zheng, J. *et al.* (2007) Serum from mice immunized in the context of Treg inhibition identifies DEK as a neuroblastoma tumor antigen. [BMC Immunol. 8: 4.](#)
5. Bahi, A. & Dreyer, J.L. (2008) Overexpression of plasminogen activators in the nucleus accumbens enhances cocaine-, amphetamine- and morphine-induced reward and behavioral sensitization. [Genes Brain Behav. 7 \(2\): 244-56.](#)
6. Wrighton, K.H. *et al.* (2009) Transforming Growth Factor {beta} Can Stimulate Smad1 Phosphorylation Independently of Bone Morphogenic Protein Receptors. [J Biol Chem. 284 \(15\): 9755-63.](#)
7. Diefenbacher, M. *et al.* (2011) The Dsl1 Tethering Complex Actively Participates in Soluble NSF (N-Ethylmaleimide-sensitive Factor) Attachment Protein Receptor (SNARE) Complex Assembly at the Endoplasmic Reticulum in *Saccharomyces cerevisiae*. [J Biol Chem. 286: 25027-38.](#)
8. Alvarez, M.M. *et al.* (2010) Specific recognition of influenza A/H1N1/2009 antibodies in human serum: a simple virus-free ELISA method. [PLoS One. 5: e10176.](#)
9. Bahi, A. *et al.* (2008) The role of tissue-type plasminogen activator system in amphetamine-induced conditional place preference extinction and reinstatement. [Neuropsychopharmacology. 33: 2726-34.](#)
10. Gunnarsen, K.S. *et al.* (2010) Periplasmic expression of soluble single chain T cell receptors is rescued by the chaperone FkpA. [BMC Biotechnol. 10: 8.](#)
11. Hwang, H.Y. *et al.* (2008) Highly specific inhibition of C1q globular-head binding to human IgG: a novel approach to control and regulate the classical complement pathway using an engineered single chain antibody variable fragment. [Mol Immunol. 45: 2570-80.](#)
12. De Vooght, L. *et al.* (2012) Expression and extracellular release of a functional anti-trypanosome Nanobody® in *Sodalis glossinidius*, a bacterial symbiont of the tsetse fly. [Microb Cell Fact. 11: 23.](#)
13. Saerens, D. *et al.* (2004) Single domain antibodies derived from dromedary lymph node and peripheral blood lymphocytes sensing conformational variants of prostate-specific antigen. [J Biol Chem. 279 \(50\): 51965-72.](#)
14. Than, N.G. *et al.* (2014) Evolutionary origins of the placental expression of chromosome 19 cluster galectins and their complex dysregulation in preeclampsia. [Placenta. 35: 855-65.](#)
15. Elders RC *et al.* (2014) Recombinant canine IgE Fc and an IgE Fc-TRAIL fusion protein bind to neoplastic canine mast cells. [Vet Immunol Immunopathol. 159 \(1-2\): 29-40.](#)
16. Chin, S.E. *et al.* (2015) Isolation of high-affinity, neutralizing anti-idiotypic antibodies by phage and ribosome display for application in immunogenicity and pharmacokinetic analyses. [J Immunol Methods. 416: 49-58.](#)
17. Peyrassol, X. *et al.* (2016) Development by Genetic Immunization of Monovalent Antibodies

- (Nanobodies) Behaving as Antagonists of the Human ChemR23 Receptor. [J Immunol. 196 \(6\): 2893-901.](#)
18. Kim H & Loparo JJ (2016) Multistep assembly of DNA condensation clusters by SMC. [Nat Commun. 7: 10200.](#)
19. Borg M *et al.* (2014) A novel interaction between Rab7b and actomyosin reveals a dual role in intracellular transport and cell migration. [J Cell Sci. 127 \(Pt 22\): 4927-39.](#)
20. De Meyer, T. *et al.* (2015) Comparison of VHH-Fc antibody production in *Arabidopsis thaliana*, *Nicotiana benthamiana* and *Pichia pastoris*. [Plant Biotechnol J. 13 \(7\): 938-47.](#)
21. Siddiqui AA *et al.* (2015) Humoral immune responses to a recombinant *Plasmodium vivax* tryptophan-rich antigen among *Plasmodium vivax*-infected patients and its localization in the parasite. [Appl Biochem Biotechnol. 175 \(4\): 2166-77.](#)
22. Warnecke, A. *et al.* (2016) Nitration of MOG diminishes its encephalitogenicity depending on MHC haplotype. [J Neuroimmunol. pii: S0165-5728\(16\)30422-2. \[Epub ahead of print\]](#)
23. Bertucci, A. *et al.* (2011) A new coral carbonic anhydrase in *Stylophora pistillata*. [Mar Biotechnol \(NY\). 13 \(5\): 992-1002.](#)
24. Liu, C.C. *et al.* (2016) The Fimbrial Protein is a Virulence Factor and Potential Vaccine Antigen of *Avibacterium paragallinarum*. [Avian Dis. 60 \(3\): 649-55.](#)
25. Boujon, C.L. *et al.* (2017) Development and validation of an immunohistochemistry procedure for the detection of a neurotropic bovine astrovirus. [J Virol Methods. 239: 26-33.](#)
26. Cartwright, S.P. *et al.* (2017) Rapid expression and purification of the hepatitis delta virus antigen using the methylotropic yeast *Pichia pastoris*. [BMC Res Notes. 10 \(1\): 340.](#)
27. Thanongsaksrikul, J. *et al.* (2018) Identification and production of mouse scFv to specific epitope of enterovirus-71 virion protein-2 (VP2). [Arch Virol. Jan 22 \[Epub ahead of print\].](#)
28. Gunnarsen, K.S. *et al.* (2018) Soluble T-cell receptor design influences functional yield in an *E. coli* chaperone-assisted expression system. [PLoS One. 13 \(4\): e0195868.](#)

---

#### Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use

---

#### Guarantee

18 months from date of despatch

---

#### Acknowledgements

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. His-tag is a registered trademark of EMD Biosciences.

---

#### Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

#### Regulatory

For research purposes only

**North & South America** Tel: +1 800 265 7376  
 Fax: +1 919 878 3751  
 Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide** Tel: +44 (0)1865 852 700  
 Fax: +44 (0)1865 852 739  
 Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe** Tel: +49 (0) 89 8090 95 21  
 Fax: +49 (0) 89 8090 95 50  
 Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M345449:190125'

