

Datasheet: MCA1424F

BATCH NUMBER 159935

Description:	MOUSE ANTI BOVINE CD21:FITC
Specificity:	CD21
Other names:	CR2
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	CC21
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Bovine		
Species Cross Reactivity	Reacts with: Goat, Sheep, Red deer, Mule deer N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant		

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
RRID	AB_323511
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NSI myeloma cell line.
Specificity	<p>Mouse anti Bovine CD21 monoclonal antibody, clone CC21 recognizes the bovine CD21 cell surface antigen, a ~145 kDa single pass type I membrane glycoprotein containing multiple sushi domains. CD21 is also known as complement receptor type 2. In cattle CD21 expression is restricted to B lymphocytes (Naessens et al. 1990). CD21 may be expressed on B cells as either a long or a short form (Pringle et al. 2012)</p> <p>Mouse anti bovine CD21, clone CC21 has been used to demonstrate the co-expression of CD21 with PrP^C on B cells of scrapie infected sheep (Halliday et al. 2005).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Howard, C.J. <i>et al.</i> (1991) Summary of workshop findings for leukocyte antigens of cattle. Vet Immunol Immunopathol. 27 (1-3): 21-7. Naessens, J. <i>et al.</i> (1990) Characterization of a bovine leucocyte differentiation antigen of 145,000 Mw restricted to B lymphocytes. Immunology 69: 525-30. Sopp, P. & Howard, C.J. (2001) IFN gamma and IL-4 production by CD4, CD8 and WC1 gamma delta TCR(+) T cells from cattle lymph nodes and blood. Vet Immunol Immunopathol. 81 (1-2): 85-96. Lwin, S. <i>et al.</i> (2009) Immune cell types involved in early uptake and transport of recombinant mouse prion protein in Peyer's patches of calves. Cell Tissue Res. 338: 343-54. Breugelmans, S. <i>et al.</i> (2011) Immunoassay of lymphocyte subsets in ovine palatine tonsils. Acta Histochem. 113: 416-22. Halliday, S. <i>et al.</i> (2005) Expression of PrPC on cellular components of sheep blood. J Gen Virol. 86 (Pt 5): 1571-9. Brackenbury, L.S. <i>et al.</i> (2005) Identification of a cell population that produces alpha/beta interferon <i>in vitro</i> and <i>in vivo</i> in response to noncytopathic bovine viral diarrhea virus. J Virol. 79: 7738-44. Breugelmans, S. <i>et al.</i> (2011) Differences between the ovine tonsils based on an immunohistochemical quantification of the lymphocyte subpopulations. Comp Immunol Microbiol Infect Dis. 34: 217-25. Richt, J.A. <i>et al.</i> (2007) Production of cattle lacking prion protein. Nat Biotechnol. 25: 132-8. Brujeni, G.N. <i>et al.</i> (2010) Bovine immunodeficiency virus and bovine leukemia virus and their mixed infection in Iranian Holstein cattle. J Infect Dev Ctries. 4 (9): 576-9.

11. Kiku, Y. *et al.* (2010) Decrease in bovine CD14 positive cells in colostrum is associated with the incidence of mastitis after calving. [Vet Res Commun. 34: 197-203.](#)
12. Pilla, R. *et al.* (2012) Long-term study of MRSA ST1, t127 mastitis in a dairy cow. [Vet Rec. 170: 312.](#)
13. Chattha, K.S. *et al.* (2010) Immunohistochemical investigation of cells expressing CD21, membrane IgM, CD32 and a follicular dendritic cell marker in the lymphoid tissues of neonatal calves. [Vet Immunol Immunopathol. 137: 284-90.](#)
14. Weiss, D.J. *et al.* (2006) Mucosal immune response in cattle with subclinical Johne's disease. [Vet Pathol. 43: 127-35.](#)
15. Summers, C. *et al.* (2012) The distribution of immune cells in the lungs of classical and atypical ovine pulmonary adenocarcinoma. [Vet Immunol Immunopathol. 146: 1-7.](#)
16. Sigurdson, C.J. *et al.* (2002) PrP(CWD) lymphoid cell targets in early and advanced chronic wasting disease of mule deer. [J Gen Virol. 83: 2617-28.](#)
17. Dagleish, M.P. *et al.* (2012) Immunophenotype of cells within cervine rectoanal mucosa-associated lymphoid tissue and mesenteric lymph nodes. [J Comp Pathol. 146: 365-71.](#)
18. Edwards, J.C. *et al.* (2010) PrP(Sc) is associated with B cells in the blood of scrapie-infected sheep. [Virology. 405: 110-9.](#)
19. Brodzki, P. *et al.* (2014) Phenotyping of leukocytes and granulocyte and monocyte phagocytic activity in the peripheral blood and uterus of cows with endometritis. [Theriogenology. 82: 403-10.](#)
20. Meganck, V. *et al.* (2014) Development of a method for isolating bovine colostrum mononuclear leukocytes for phenotyping and functional studies. [Vet J. 200: 294-8.](#)
21. Booth, J.S. *et al.* (2010) Co-stimulation with TLR7/8 and TLR9 agonists induce down-regulation of innate immune responses in sheep blood mononuclear and B cells. [Dev Comp Immunol. 34 \(5\): 572-8.](#)
22. Nikbakht Brujeni, G. *et al.* (2016) Association of BoLA-DRB3.2 Alleles with BLV Infection Profiles (Persistent Lymphocytosis/Lymphosarcoma) and Lymphocyte Subsets in Iranian Holstein Cattle. [Biochem Genet. 54 \(2\): 194-207.](#)
23. Kruger, E.F. *et al.* (2003) Bovine monocytes induce immunoglobulin production in peripheral blood B lymphocytes. [Dev Comp Immunol. 27 \(10\): 889-97.](#)
24. De Matteis G *et al.* (2016) Evaluation of leptin receptor expression on buffalo leukocytes. [Vet Immunol Immunopathol. 177: 16-23.](#)
25. Ekman, A. *et al.* (2010) B-cell development in bovine fetuses proceeds via a pre-B like cell in bone marrow and lymph nodes. [Dev Comp Immunol. 34 \(8\): 896-903.](#)
26. Silva, A.P. *et al.* (2015) Encapsulated *Brucella ovis* Lacking a Putative ATP-Binding Cassette Transporter (Δ abcBA) Protects against Wild Type *Brucella ovis* in Rams. [PLoS One. 10 \(8\): e0136865.](#)
27. Jimbo, S. *et al.* (2019) Natural and inducible regulatory B cells are widely distributed in ovine lymphoid tissues. [Vet Immunol Immunopathol. 211: 44-8.](#)
28. Brodzki, P. *et al.* (2019) Selected leukocyte subpopulations in peripheral blood and uterine washings in cows before and after intrauterine administration of cefapirin and methisoprinol. [Anim Sci J. Nov 06 \[Epub ahead of print\].](#)
29. Souza, F.N. *et al.* (2020) Lymphocyte proliferative responses in dairy cows supplemented with an immunomodulatory feed additive and administered polyvalent vaccination. [Arquivo Brasileiro de Medicina Veterinária e Zootecnia. 72 \(6\): 2397-401.](#)
30. Gondaira, S. *et al.* (2020) Immunosuppression in Cows following Intramammary

- Infusion of *Mycoplasma bovis*. [Infect Immun. 88 \(3\) Feb 20 \[Epub ahead of print\]](#).
31. Khosa, S. *et al.* (2020) Bovine Adenovirus-3 Tropism for Bovine Leukocyte Sub-Populations. [Viruses. 12 \(12\) Dec 12 \[Epub ahead of print\]](#).
32. Ramm, R. *et al.* (2020) Decellularization combined with enzymatic removal of N-linked glycans and residual DNA reduces inflammatory response and improves performance of porcine xenogeneic pulmonary heart valves in an ovine *in vivo* model. [Xenotransplantation. 27 \(2\): e12571.](#)
33. Ramos, A. *et al.* (2018) Melatonin enhances responsiveness to *Dichelobacter nodosus* vaccine in sheep and increases peripheral blood CD4 T lymphocytes and IgG-expressing B lymphocytes. [Vet Immunol Immunopathol. 206: 1-8.](#)
34. Okino, C.H. *et al.* (2020) A polymorphic CD4 epitope related to increased susceptibility to *Babesia bovis*. in Canchim calves. [Vet Immunol Immunopathol. 230: 110132.](#)
35. Park, D.S. *et al.* (2021) Dynamic changes in blood immune cell composition and function in Holstein and Jersey steers in response to heat stress. [Cell Stress Chaperones. 26 \(4\): 705-20.](#)
36. Radley, G. *et al.* (2020) *In Vitro*. Benchmarking Study of Ventricular Assist Devices in Current Clinical Use. [J Card Fail. 26 \(1\): 70-79.](#)
37. Colombatti Olivieri, M.A. *et al.* (2021) Evaluation of a virulent strain of *Mycobacterium avium*. subsp. *paratuberculosis*. used as a heat-killed vaccine. [Vaccine. NNov 10;S0264-410X\(21\)01433-X.](#)
38. Casaro, S. *et al.* (2022) Flow cytometry panels for immunophenotyping dairy cattle peripheral blood leukocytes [Vet Immunol Immunopathol. 248: 110417.](#)
39. Okino, C.H. *et al.* (2022) CD4 bovine gene: Differential polymorphisms among cattle breeds and a new tool for rapid identification. [Vet Immunol Immunopathol. 251: 110462.](#)

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
----------------	---

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
------------------	---------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1424F 10041
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M385465:210513'

Printed on 21 Feb 2024