

Datasheet: MCA1427

BATCH NUMBER 0315

Description:	MOUSE ANTI RAT CD161
Specificity:	CD161
Other names:	NATURAL KILLER CELLS, NKR-P1A
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	10/78
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			
Radioimmunoassays	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Purified splenic NK cells from the LEW rat strain.

External Database Links

UniProt:

[P27471](#)

[Related reagents](#)

[A4KWA1](#)

[Related reagents](#)

Entrez Gene:

[362443](#)

Klrb1a

[Related reagents](#)

[25192](#)

Klrb1b

[Related reagents](#)

Synonyms Nkrp1a, Nkrp1b

RRID AB_2234351

Fusion Partners Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.Ag8653 myeloma cell line.

Specificity **Mouse anti Rat CD161 antibody, clone 10/78** recognizes the rat Killer cell lectin-like receptor subfamily B protein, also known as NKR-PI or CD161. CD161 is a 233 amino acid ~60 kDa type II single pass protein containing a single [C-type lectin](#) domain. CD161 is expressed on rat NK cells and T cell subpopulations. CD161 exists in 2 forms NKR-PIa and NKR-PIb, Mouse anti Rat CD161 antibody, clone 10/78 recognizes both forms of CD161 ([Li et al. 2003](#)). Clone 10/78 competes with another anti CD161 clone, 3.2.3 for binding to antigen.

Mouse anti Rat CD161 antibody, clone 10/78 has been successfully employed for the *in vivo* depletion of rat NK cells in an experimental obesity model ([Wrann et al. 2010](#)).

References

1. Dyugovskaya, L. *et al.* (2003) Phenotypic profile and functional characterization of rat lymph node-derived gammadelta T cells: implication in the immune response to cytomegalovirus. [Immunology. 108 \(2\): 129-36.](#)
2. Sedgwick, J.D. *et al.* (1998) Central nervous system microglial cell activation and proliferation follows direct interaction with tissue-infiltrating T cell blasts. [J Immunol. 160 \(11\): 5320-30.](#)
3. Schwartzkopff, J. *et al.* (2010) NK cell depletion delays corneal allograft rejection in baby rats. [Mol Vis. 16: 1928-35.](#)
4. Lyons, A. *et al.* (2011) Atorvastatin prevents age-related and amyloid-beta-induced microglial activation by blocking interferon-gamma release from natural killer cells in the brain. [J Neuroinflammation. 8: 27.](#)
5. Ali, S. *et al.* (2005) Combined immunostimulation and conditional cytotoxic gene

- therapy provide long-term survival in a large glioma model. [Cancer Res. 65: 7194-204.](#)
6. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. [Br J Ophthalmol. 87: 1515-22.](#)
 7. Latta, M. *et al.* (2007) CXCR6 is expressed on T cells in both T helper type 1 (Th1) inflammation and allergen-induced Th2 lung inflammation but is only a weak mediator of chemotaxis. [Immunology. 121: 555-64.](#)
 8. Tliba, O. *et al.* (2002) Evaluation of the hepatic NK cell response during the early phase of *Fasciola hepatica* infection in rats. [Vet Res. 33 \(3\): 327-32.](#)
 9. Blöcher, S. *et al.* (2007) Acute rejection of experimental lung allografts: characterization of intravascular mononuclear leukocytes. [Clin Immunol. 124 \(1\): 98-108.](#)
 10. Koch, M. *et al.* (2015) Extracellular Vesicles from MSC Modulate the Immune Response to Renal Allografts in a MHC Disparate Rat Model. [Stem Cells Int. 2015: 486141.](#)
 11. Trama, A.M. *et al.* (2012) Lymphocyte phenotypes in wild-caught rats suggest potential mechanisms underlying increased immune sensitivity in post-industrial environments. [Cell Mol Immunol. 9 \(2\): 163-74.](#)
 12. Wrann, C.D. *et al.* (2010) Obesity and NK cells affect the expression of the long form of the leptin receptor Ob-Rb in liver of F344 rats. [Exp Toxicol Pathol. 62 \(1\): 1-8.](#)
 13. Ikezumi, Y. *et al.* (2000) An anti-CD5 monoclonal antibody ameliorates proteinuria and glomerular lesions in rat mesangioproliferative glomerulonephritis. [Kidney Int. 58 \(1\): 100-14.](#)
 14. Obara, H. *et al.* (2005) IFN-gamma, produced by NK cells that infiltrate liver allografts early after transplantation, links the innate and adaptive immune responses. [Am J Transplant. 5 \(9\): 2094-103.](#)
 15. Beutel, G. *et al.* (2013) Effect of chronic elevated asymmetric dimethylarginine (ADMA) levels on granulopoiesis. [Ann Hematol. 92 \(4\): 505-8.](#)
 16. Lee, J.S. *et al.* (2011) Immunomodulatory effect of mushrooms on cytotoxic activity and cytokine production of intestinal lamina propria leukocytes does not necessarily depend on β -glucan contents. [Food Chem. 126 \(4\): 1521-6.](#)
 17. Williamson, L.L. *et al.* (2016) Got worms? Perinatal exposure to helminths prevents persistent immune sensitization and cognitive dysfunction induced by early-life infection. [Brain Behav Immun. 51: 14-28.](#)
 18. Arndt, T. *et al.* (2014) Variable immune cell frequencies in peripheral blood of LEW.1AR1-iddm rats over time compared to other congenic LEW strains. [Clin Exp Immunol. 177 \(1\): 168-78.](#)
 19. Kuper, C.F. *et al.* (2011) Oxazolone (OXA) is a respiratory allergen in Brown Norway rats. [Toxicology. 290 \(1\): 59-68.](#)
 20. Arsenović-Ranin, N. *et al.* (2013) Ovarian hormone withdrawal in prepubertal developmental stage does not prevent thymic involution in rats. [Exp Biol Med \(Maywood\). 238 \(6\): 641-57.](#)
 21. Djikić J *et al.* (2014) Age-associated changes in rat immune system: lessons learned from experimental autoimmune encephalomyelitis. [Exp Gerontol. 58: 179-97.](#)
 22. Lemke, A. *et al.* (2015) Rat renal transplant model for mixed acute humoral and cellular rejection: Weak correlation of serum cytokines/chemokines with intragraft changes. [Transpl Immunol. 33 \(2\): 95-102.](#)
 23. Bähr, I. *et al.* (2017) Diet-Induced Obesity Is Associated with an Impaired NK Cell Function and an Increased Colon Cancer Incidence. [J Nutr Metab. 2017: 4297025.](#)

24. Sun, C.K. *et al.* (2017) Melatonin treatment enhances therapeutic effects of exosomes against acute liver ischemia-reperfusion injury. [Am J Transl Res. 9 \(4\): 1543-60.](#)
25. Chang, J.C. *et al.* (2019) Early Immune Response to Acute Gastric Fluid Aspiration in a Rat Model of Lung Transplantation. [Exp Clin Transplant. 17 \(1\): 84-92.](#)

Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1427 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M365295:200529'

Printed on 18 Jan 2024