

# Datasheet: MCA1427GA

**BATCH NUMBER 148575**

<b>Description:</b>	MOUSE ANTI RAT CD161
<b>Specificity:</b>	CD161
<b>Other names:</b>	NATURAL KILLER CELLS, NKR-P1A
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	10/78
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Rat
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified splenic NK cells from the LEW rat strain.
External Database Links	<p><b>UniProt:</b></p> <p><a href="#">P27471</a>      <a href="#">Related reagents</a></p> <p><a href="#">A4KWA1</a>      <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">362443</a>    Klr1a    <a href="#">Related reagents</a></p> <p><a href="#">25192</a>    Klr1b    <a href="#">Related reagents</a></p>
Synonyms	Nkrp1a, Nkrp1b
RRID	AB_566557
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.Ag8653 myeloma cell line.
Specificity	<p><b>Mouse anti Rat CD161 antibody, clone 10/78</b> recognizes the rat Killer cell lectin-like receptor subfamily B protein, also known as NKR-PI or CD161. CD161 is a 233 amino acid ~60 kDa type II single pass protein containing a single <a href="#">C-type lectin</a> domain. CD161 is expressed on rat NK cells and T cell subpopulations. CD161 exists in 2 forms NKR-PIa and NKR-PIb, Mouse anti Rat CD161 antibody, clone 10/78 recognizes both forms of CD161 (<a href="#">Li et al. 2003</a>). Clone 10/78 competes with another anti CD161 clone, 3.2.3 for binding to antigen.</p> <p>Mouse anti Rat CD161 antibody, clone 10/78 has been successfully employed for the <i>in vivo</i> depletion of rat NK cells in an experimental obesity model (<a href="#">Wrann et al. 2010</a>).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	<ol style="list-style-type: none"> <li>1. Dyugovskaya, L. <i>et al.</i> (2003) Phenotypic profile and functional characterization of rat lymph node-derived gammadelta T cells: implication in the immune response to cytomegalovirus. <a href="#">Immunology. 108 (2): 129-36.</a></li> <li>2. Sedgwick, J.D. <i>et al.</i> (1998) Central nervous system microglial cell activation and proliferation follows direct interaction with tissue-infiltrating T cell blasts. <a href="#">J Immunol. 160 (11): 5320-30.</a></li> <li>3. Schwartzkopff, J. <i>et al.</i> (2010) NK cell depletion delays corneal allograft rejection in baby rats. <a href="#">Mol Vis. 16: 1928-35.</a></li> <li>4. Lyons, A. <i>et al.</i> (2011) Atorvastatin prevents age-related and amyloid-beta-induced microglial activation by blocking interferon-gamma release from natural killer cells in the brain. <a href="#">J Neuroinflammation. 8: 27.</a></li> </ol>

5. Ali, S. *et al.* (2005) Combined immunostimulation and conditional cytotoxic gene therapy provide long-term survival in a large glioma model. [Cancer Res. 65: 7194-204.](#)
6. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. [Br J Ophthalmol. 87: 1515-22.](#)
7. Latta, M. *et al.* (2007) CXCR6 is expressed on T cells in both T helper type 1 (Th1) inflammation and allergen-induced Th2 lung inflammation but is only a weak mediator of chemotaxis. [Immunology. 121: 555-64.](#)
8. Tliba, O. *et al.* (2002) Evaluation of the hepatic NK cell response during the early phase of *Fasciola hepatica* infection in rats. [Vet Res. 33 \(3\): 327-32.](#)
9. Blöcher, S. *et al.* (2007) Acute rejection of experimental lung allografts: characterization of intravascular mononuclear leukocytes. [Clin Immunol. 124 \(1\): 98-108.](#)
10. Koch, M. *et al.* (2015) Extracellular Vesicles from MSC Modulate the Immune Response to Renal Allografts in a MHC Disparate Rat Model. [Stem Cells Int. 2015: 486141.](#)
11. Trama, A.M. *et al.* (2012) Lymphocyte phenotypes in wild-caught rats suggest potential mechanisms underlying increased immune sensitivity in post-industrial environments. [Cell Mol Immunol. 9 \(2\): 163-74.](#)
12. Wrann, C.D. *et al.* (2010) Obesity and NK cells affect the expression of the long form of the leptin receptor Ob-Rb in liver of F344 rats. [Exp Toxicol Pathol. 62 \(1\): 1-8.](#)
13. Ikezumi, Y. *et al.* (2000) An anti-CD5 monoclonal antibody ameliorates proteinuria and glomerular lesions in rat mesangioproliferative glomerulonephritis. [Kidney Int. 58 \(1\): 100-14.](#)
14. Obara, H. *et al.* (2005) IFN-gamma, produced by NK cells that infiltrate liver allografts early after transplantation, links the innate and adaptive immune responses. [Am J Transplant. 5 \(9\): 2094-103.](#)
15. Beutel, G. *et al.* (2013) Effect of chronic elevated asymmetric dimethylarginine (ADMA) levels on granulopoiesis. [Ann Hematol. 92 \(4\): 505-8.](#)
16. Lee, J.S. *et al.* (2011) Immunomodulatory effect of mushrooms on cytotoxic activity and cytokine production of intestinal lamina propria leukocytes does not necessarily depend on  $\beta$ -glucan contents. [Food Chem. 126 \(4\): 1521-6.](#)
17. Williamson, L.L. *et al.* (2016) Got worms? Perinatal exposure to helminths prevents persistent immune sensitization and cognitive dysfunction induced by early-life infection. [Brain Behav Immun. 51: 14-28.](#)
18. Arndt, T. *et al.* (2014) Variable immune cell frequencies in peripheral blood of LEW.1AR1-iddm rats over time compared to other congenic LEW strains. [Clin Exp Immunol. 177 \(1\): 168-78.](#)
19. Kuper, C.F. *et al.* (2011) Oxazolone (OXA) is a respiratory allergen in Brown Norway rats. [Toxicology. 290 \(1\): 59-68.](#)
20. Arsenović-Ranin, N. *et al.* (2013) Ovarian hormone withdrawal in prepubertal developmental stage does not prevent thymic involution in rats. [Exp Biol Med \(Maywood\). 238 \(6\): 641-57.](#)
21. Djikić J *et al.* (2014) Age-associated changes in rat immune system: lessons learned from experimental autoimmune encephalomyelitis. [Exp Gerontol. 58: 179-97.](#)
22. Lemke, A. *et al.* (2015) Rat renal transplant model for mixed acute humoral and cellular rejection: Weak correlation of serum cytokines/chemokines with intragraft changes. [Transpl Immunol. 33 \(2\): 95-102.](#)
23. Bähr, I. *et al.* (2017) Diet-Induced Obesity Is Associated with an Impaired NK Cell

Function and an Increased Colon Cancer Incidence. [J Nutr Metab. 2017: 4297025.](#)  
 24. Sun, C.K. *et al.* (2017) Melatonin treatment enhances therapeutic effects of exosomes against acute liver ischemia-reperfusion injury. [Am J Transl Res. 9 \(4\): 1543-60.](#)  
 25. Chang, J.C. *et al.* (2019) Early Immune Response to Acute Gastric Fluid Aspiration in a Rat Model of Lung Transplantation. [Exp Clin Transplant. 17 \(1\): 84-92.](#)

<b>Storage</b>	Store at +4°C or at -20°C if preferred.  This product should be stored undiluted.  Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1427GA">https://www.bio-rad-antibodies.com/SDS/MCA1427GA</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA1209\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
 'M365299:200529'

Printed on 18 Jan 2024

