## Datasheet: MCA1427GA BATCH NUMBER 162871

Description:	MOUSE ANTI RAT CD161
Specificity:	CD161
Other names:	NKR-P1A
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	10/78
Isotype:	lgG1
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	-			1/50 - 1/100
	Immunohistology - Frozen	-			
	Immunohistology - Paraffin				
	ELISA				
	Immunoprecipitation	-			
Target Species	necessarily exclude its us a guide only. It is recomn system using appropriate Rat	nended th	nat the us	er titrates the antibody	•
Product Form	Purified IgG - liquid				
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% Sodium Azide				

Carrier Free	Yes		
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml		
Immunogen	Purified splenic NK cells from the LEW rat strain.		
External Database Links	UniProt: $P27471$ Related reagents $A4KWA1$ Related reagentsEntrez Gene: $362443$ Klrb1aRelated reagents $25192$ Klrb1bRelated reagents		
Synonyms	Nkrp1a, Nkrp1b		
RRID	AB_566557		
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse X63.Ag8653 myeloma cell line.		
Specificity	<b>Mouse anti Rat CD161 antibody, clone 10/78</b> recognizes the rat Killer cell lectin-like receptor subfamily B protein, also known as NKR-PI or CD161. CD161 is a 233 amino acid ~60 kDa type II single pass protein containing a single <u>C-type lectin</u> domain. CD161 is expressed on rat NK cells and T cell subpopulations. CD161 exists in 2 forms NKR-PIa and NKR-PIb, Mouse anti Rat CD161 antibody, clone 10/78 recognizes both forms of CD161 ( <u>Li <i>et al.</i> 2003</u> ). Clone 10/78 competes with another anti CD161 clone, 3.2.3 for binding to antigen.		
	Mouse anti Rat CD161 antibody, clone 10/78 has been successfully employed for the <i>in vivo</i> depletion of rat NK cells in an experimental obesity model ( <u>Wrann <i>et al.</i> 2010</u> ).		
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.		
References	<ol> <li>Dyugovskaya, L. <i>et al.</i> (2003) Phenotypic profile and functional characterization of rat lymph node-derived gammadelta T cells: implication in the immune response to cytomegalovirus. <u>Immunology. 108 (2): 129-36.</u></li> <li>Sedgwick, J.D. <i>et al.</i> (1998) Central nervous system microglial cell activation and proliferation follows direct interaction with tissue-infiltrating T cell blasts. <u>J Immunol. 160</u> (<u>11): 5320-30.</u></li> <li>Schwartzkopff, J. <i>et al.</i> (2010) NK cell depletion delays corneal allograft rejection in baby rats. <u>Mol Vis. 16: 1928-35.</u></li> <li>Lyons, A. <i>et al.</i> (2011) Atorvastatin prevents age-related and amyloid-beta-induced microglial activation by blocking interferon-gamma release from natural killer cells in the brain. <u>J Neuroinflammation. 8: 27.</u></li> </ol>		

5. Ali, S. *et al.* (2005) Combined immunostimulation and conditional cytotoxic gene therapy provide long-term survival in a large glioma model. <u>Cancer Res. 65: 7194-204.</u>
6. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. <u>Br J Ophthalmol. 87: 1515-22.</u>

7. Latta, M. *et al.* (2007) CXCR6 is expressed on T cells in both T helper type 1 (Th1) inflammation and allergen-induced Th2 lung inflammation but is only a weak mediator of chemotaxis. <u>Immunology. 121: 555-64.</u>

8. Tliba, O. *et al.* (2002) Evaluation of the hepatic NK cell response during the early phase of *Fasciola hepatica* infection in rats. <u>Vet Res. 33 (3): 327-32.</u>

9. Blöcher, S. *et al.* (2007) Acute rejection of experimental lung allografts: characterization of intravascular mononuclear leukocytes. <u>Clin Immunol. 124 (1): 98-108.</u>

10. Koch, M. *et al.* (2015) Extracellular Vesicles from MSC Modulate the Immune Response to Renal Allografts in a MHC Disparate Rat Model. <u>Stem Cells Int. 2015:</u> <u>486141.</u>

11. Trama, A.M. *et al.* (2012) Lymphocyte phenotypes in wild-caught rats suggest potential mechanisms underlying increased immune sensitivity in post-industrial environments. <u>Cell</u> <u>Mol Immunol. 9 (2): 163-74.</u>

12. Wrann, C.D. *et al.* (2010) Obesity and NK cells affect the expression of the long form of the leptin receptor Ob-Rb in liver of F344 rats. <u>Exp Toxicol Pathol. 62 (1): 1-8.</u>

13. Ikezumi, Y. *et al.* (2000) An anti-CD5 monoclonal antibody ameliorates proteinuria and glomerular lesions in rat mesangioproliferative glomerulonephritis. <u>Kidney Int. 58 (1):</u> 100-14.

14. Obara, H. *et al.* (2005) IFN-gamma, produced by NK cells that infiltrate liver allografts early after transplantation, links the innate and adaptive immune responses. <u>Am J</u><u>Transplant. 5 (9): 2094-103.</u>

15. Beutel, G. *et al.* (2013) Effect of chronic elevated asymmetric dimethylarginine (ADMA) levels on granulopoiesis. <u>Ann Hematol. 92 (4): 505-8.</u>

16. Lee, J.S. *et al.* (2011) Immunomodulatory effect of mushrooms on cytotoxic activity and cytokine production of intestinal lamina propria leukocytes does not necessarily depend on  $\beta$ -glucan contents. <u>Food Chem. 126 (4): 1521-6.</u>

17. Williamson, L.L. *et al.* (2016) Got worms? Perinatal exposure to helminths prevents persistent immune sensitization and cognitive dysfunction induced by early-life infection. <u>Brain Behav Immun. 51: 14-28.</u>

18. Arndt, T. *et al.* (2014) Variable immune cell frequencies in peripheral blood of LEW.1AR1-iddm rats over time compared to other congenic LEW strains. <u>Clin Exp</u> <u>Immunol. 177 (1): 168-78.</u>

19. Kuper, C.F. *et al.* (2011) Oxazolone (OXA) is a respiratory allergen in Brown Norway rats. <u>Toxicology. 290 (1): 59-68.</u>

20. Arsenović-Ranin, N. *et al.* (2013) Ovarian hormone withdrawal in prepubertal developmental stage does not prevent thymic involution in rats. <u>Exp Biol Med (Maywood).</u> 238 (6): 641-57.

21. Djikić J *et al.* (2014) Age-associated changes in rat immune system: lessons learned from experimental autoimmune encephalomyelitis. <u>Exp Gerontol. 58: 179-97.</u>

22. Lemke, A. *et al.* (2015) Rat renal transplant model for mixed acute humoral and cellular rejection: Weak correlation of serum cytokines/chemokines with intragraft changes. <u>Transpl Immunol. 33 (2): 95-102.</u>

23. Bähr, I. et al. (2017) Diet-Induced Obesity Is Associated with an Impaired NK Cell

	Function and an Increased Colon Cancer Incidence. J Nutr Metab. 2017: 4297025.				
	24. Sun, C.K. et al. (2017) Melatonin treatment enhances therapeutic effects of exosomes				
	against acute liver ischemia-reperfusion injury. Am J Transl Res. 9 (4): 1543-60.				
	25. Chang, J.C. et al. (2019) Early Immune Response to Acute Gastric Fluid Aspira				
	a Rat Model of Lung Transplantation. Exp Clin Transplant. 17 (1): 84-92.				
	proves immune dysfunction				
	in the hippocampus after hypoxic-ischemic brain damage <u>Int J</u>	Med Sci. 19 (1): 142-51.			
Storage	StorageThis product is shipped at ambient temperature. It is recommended to aliquot and -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2 short term use (up to 4 weeks) and store the remaining aliquots at -20°C.				
	Avoid repeated freezing and thawing as this may denature the frost-free freezers is not recommended.	antibody. Storage in			
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1427GA 10040				
Regulatory	For research purposes only				

## **Related Products**

## **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12)	RPE		
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>			
Goat Anti Mouse IgG (STAR76)	<u>RPE</u>		
Rabbit Anti Mouse IgG (STAR13)	HRP		
Goat Anti Mouse IgG (STAR70)	FITC		
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,		
	DyLight®650, DyLight®680, DyLight®800,		
	<u>FITC</u> , <u>HRP</u>		
Rabbit Anti Mouse IgG (STAR9)	<u>FITC</u>		
Rabbit Anti Mouse IgG (STAR9) Goat Anti Mouse IgG (STAR77)	FITC HRP		
<b>5</b> ( )			

MOUSE IgG1 NEGATIVE CONTROL (MCA1209)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@b	io-rad.com	Email: antibody_sales_uk@bio	o-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M383053:210513'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint