

Datasheet: MCA1436

BATCH NUMBER 148130

Description:	tion: MOUSE ANTI hCG HOLO	
Specificity:	hCG HOLO	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	INN-hCG-53	
Isotype:	lgG1	
Quantity:	0.5 mg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA				
Immunoprecipitation			•	
Western Blotting				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	A from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)	
Carrier Free	Yes	

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
External Database Links	UniProt:
	P01233 Related reagents
	P01215 Related reagents
	Entrez Gene:
	1082 CGB Related reagents
	1081 CGA Related reagents
Synonyms	CGB3
RRID	AB_322005
Specificity	Mouse anti holo-chorilonic gonadotrophin anribody, clone INN-hCG-53 recognises epitope c2 on hCG. It does not recognize free hCG subunits.
Western Blotting	Clone INN-hCG-53 detects a band of about 47-50 kDa under non-reducing conditions.
References	1. Lei, Z.M. <i>et al.</i> (1993) Novel coexpression of human chorionic gonadotropin (hCG)/human luteinizing hormone receptors and their ligand hCG in human fallopian tubes. <u>J Clin Endocrinol Metab. 77 (3): 863-72.</u>
	2. Ben-Menahem, D. <i>et al.</i> (2001) The position of the alpha and beta subunits in a single chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the
	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u>
	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol</u>
	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol Endocrinol. 13 (12): 2175-88.</u>
	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol</u>
Storage	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol Endocrinol. 13 (12): 2175-88.</u> 4. Klonisch, T. <i>et al.</i> (1996) Relative location of epitopes involved in synergistic antibody
Storage	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol Endocrinol. 13 (12): 2175-88.</u> 4. Klonisch, T. <i>et al.</i> (1996) Relative location of epitopes involved in synergistic antibody binding using human chorionic gonadotropin as a model. <u>Eur J Immunol. 26: 1897-905.</u> Store at +4°C or at -20°C if preferred.
Storage	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol Endocrinol. 13 (12): 2175-88.</u> 4. Klonisch, T. <i>et al.</i> (1996) Relative location of epitopes involved in synergistic antibody binding using human chorionic gonadotropin as a model. <u>Eur J Immunol. 26: 1897-905.</u> Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this mandenature the antibody. Should this product contain a precipitate we recommend
Storage	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol Endocrinol. 13 (12): 2175-88.</u> 4. Klonisch, T. <i>et al.</i> (1996) Relative location of epitopes involved in synergistic antibody binding using human chorionic gonadotropin as a model. <u>Eur J Immunol. 26: 1897-905.</u> Store at +4°C or at -20°C if preferred.
Storage	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol Endocrinol. 13 (12): 2175-88.</u> 4. Klonisch, T. <i>et al.</i> (1996) Relative location of epitopes involved in synergistic antibody binding using human chorionic gonadotropin as a model. <u>Eur J Immunol. 26: 1897-905.</u> Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this mandenature the antibody. Should this product contain a precipitate we recommend
Guarantee Health And Safety	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol Endocrinol. 13 (12): 2175-88.</u> 4. Klonisch, T. <i>et al.</i> (1996) Relative location of epitopes involved in synergistic antibody binding using human chorionic gonadotropin as a model. <u>Eur J Immunol. 26: 1897-905.</u> Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at:
Guarantee Health And Safety	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. J Biol Chem. 276 (32): 29871-9. 3. Jackson, A.M. et al. (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. Mol Endocrinol. 13 (12): 2175-88. 4. Klonisch, T. et al. (1996) Relative location of epitopes involved in synergistic antibody binding using human chorionic gonadotropin as a model. Eur J Immunol. 26: 1897-905. Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1436
Guarantee	chain variant of human chorionic gonadotropin affects the heterodimeric interaction of the subunits and receptor-binding epitopes. <u>J Biol Chem. 276 (32): 29871-9.</u> 3. Jackson, A.M. <i>et al.</i> (1999) The biological action of choriogonadotropin is not dependent on the complete native quaternary interactions between the subunits. <u>Mol Endocrinol. 13 (12): 2175-88.</u> 4. Klonisch, T. <i>et al.</i> (1996) Relative location of epitopes involved in synergistic antibody binding using human chorionic gonadotropin as a model. <u>Eur J Immunol. 26: 1897-905.</u> Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. 12 months from date of despatch Material Safety Datasheet documentation #10040 available at:

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)

Goat Anti Mouse IgG IgA IgM (STAR87...)

Goat Anti Mouse IgG (STAR76...)

RPE

Goat Anti Mouse IgG (STAR70...)

FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR77...) HRP
Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365308:200529'

Printed on 21 Jun 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint