

Datasheet: MCA1439

BATCH NUMBER 1606

Description:	RAT ANTI MOUSE CD19
Specificity:	CD19
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	6D5
Isotype:	IgG2a
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human K562 cell line transfected with murine CD19.
External Database Links	<p>UniProt: P25918 Related reagents</p> <p>Entrez Gene: 12478 Cd19 Related reagents</p>
RRID	AB_321342
Fusion Partners	Spleen cells from immunised rats were fused with cells of the P3X63.Ag8.653 myeloma cell line.
Specificity	Rat anti Mouse CD19 antibody, clone 6D5 recognizes the murine CD19 cell surface antigen, a ~95 kDa glycoprotein expressed by B lymphocytes. Rat anti Mouse CD19 antibody, clone 6D5 recognizes the same, or a closely related epitope as clone Rat anti Mouse antibody, clone ID3 in cross-competition assays.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. Vernooij, J.H. <i>et al.</i> (2002) Long-term intratracheal lipopolysaccharide exposure in mice results in chronic lung inflammation and persistent pathology. Am J Respir Cell Mol Biol. 26 (1): 152-9. 2. Andrew, D. and Aspinall. R. (2001) Il-7 and not stem cell factor reverses both the increase in apoptosis and the decline in thymopoiesis seen in aged mice. J Immunol. 166: 1524-30. 3. Bermudez-Fajardo, A. <i>et al.</i> (2011) The effect of <i>Chlamydomytila pneumoniae</i> Major Outer Membrane Protein (MOMP) on macrophage and T cell-mediated immune responses. Immunobiology. 216: 152-63. 4. De Jesus, M. <i>et al.</i> (2009) Galactoxylomannan-mediated immunological paralysis results from specific B cell depletion in the context of widespread immune system damage. J Immunol. 183: 3885-94. 5. Jégou, J.F. <i>et al.</i> (2007) C3d binding to the myelin oligodendrocyte glycoprotein results in an exacerbated experimental autoimmune encephalomyelitis. J Immunol. 178: 3323-31. 6. Starck, J. <i>et al.</i> (2010) Inducible Fli-1 gene deletion in adult mice modifies several myeloid lineage commitment decisions and accelerates proliferation arrest and terminal erythrocytic differentiation. Blood. 116: 4795-805. 7. Scotland, R.S. <i>et al.</i> (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. Blood. 118 (22): 5918-27. 8. White, H.N. and Meng, Q.H. (2012) Recruitment of a Distinct but Related Set of VH Sequences into the Murine CD21hi/CD23- Marginal Zone B Cell Repertoire to That Seen in the Class-Switched Antibody Response. J Immunol. 188: 287-93. 9. Reynaud, J.M. <i>et al.</i> (2014) Human herpesvirus 6A infection in CD46 transgenic mice: viral persistence in the brain and increased production of proinflammatory chemokines via

- Toll-like receptor 9. [J Virol. 88: 5421-36.](#)
10. Candolfi, M. *et al.* (2011) B cells are critical to T-cell-mediated antitumor immunity induced by a combined immune-stimulatory/conditionally cytotoxic therapy for glioblastoma. [Neoplasia. 13: 947-60.](#)
11. Takabayashi, H. *et al.* (2014) Anti-inflammatory activity of bone morphogenetic protein signaling pathways in stomachs of mice. [Gastroenterology. 147: 396-406.e7.](#)
12. Weiss-Gayet, M. *et al.* (2016) Notch Stimulates Both Self-Renewal and Lineage Plasticity in a Subset of Murine CD9High Committed Megakaryocytic Progenitors. [PLoS One. 11 \(4\): e0153860.](#)
13. Meng, Q.H. & White, H.N. (2017) CD21^{int} CD23⁺ follicular B cells express antigen-specific secretory IgM mRNA as primary and memory responses. [Immunology. 151 \(2\): 211-8.](#)
14. Mccubbrey, A.L. *et al.* (2016) MicroRNA-34a Negatively Regulates Efferocytosis by Tissue Macrophages in Part via SIRT1. [J Immunol. 196 \(3\): 1366-75.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1439>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

'M365312:200529'

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 20 Nov 2023

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)