

## Datasheet: MCA1439APC

Description:	RAT ANTI MOUSE CD19:APC
Specificity:	CD19
Format:	APC
Product Type:	Monoclonal Antibody
Clone:	6D5
lsotype:	lgG2a
Quantity:	100 TESTS

## **Product Details**

RRID	AB_324804					
Applications	This product has been rep from testing within our labor the originators. Please refe recommendations, please Flow Cytometry Where this antibody has n exclude its use in such pro-	oratories, peer-re er to references i visit <u>www.bio-rac</u> Yes N • ot been tested fo ocedures. Sugge	viewed publications ndicated for further i <u>d-antibodies.com/pro</u> o Not Determ or use in a particular sted working dilution	or personal comm nformation. For ge tocols. nined Sug technique this doe as are given as a g	unications from eneral protocol gested Dilution Neat - 1/10 es not necessarily uide only. It is	
	recommended that the use negative/positive controls.		body for use in their		j appropriate	
Target Species	Mouse					
Product Form	Purified IgG conjugated to	Allophycocyanir	(APC) - lyophilised			
Reconstitution	Reconstitute with 1 ml dist	illed water				
Max Ex/Em	FluorophoreExcAPC	citation Max (nm) 650	Emission Max (nm) 661	_		
Preparation	Purified IgG prepared by a	ffinity chromatog	raphy on Protein G	from tissue culture	supernatant	
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin 5% Sucrose					
Immunogen	Human K562 cell line transfected with murine CD19.					
External Database Links	UniProt: P25918 Related rea	agents				

	Entrez Gene: <u>12478</u> Cd19 <u>Related reagents</u>				
Fusion Partners	Spleen cells from immunised rats were fused with cells of the P3X63.Ag8.653 myeloma cell line.				
Specificity	<b>Rat anti Mouse CD19 antibody, clone 6D5</b> recognizes the murine CD19 cell surface antigen, a 95kDa glycoprotein expressed by B lymphocytes. Clone 6D5 has been shown to recognize the same epitope as clone ID3 in cross-competition assays.				
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.				
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ( <u>BUF041A/B</u> ).				
References	<ol> <li>Vernooy, J.H. <i>et al.</i> (2002) Long-term intratracheal lipopolysaccharide exposure in mice results in chronic lung inflammation and persistent pathology. <u>Am J Respir Cell Mol Biol. 26 (1): 152-9</u>.</li> <li>Andrew, D. and Aspinall. R. (2001) II-7 and not stem cell factor reverses both the increase in apoptosis and the decline in thymopoiesis seen in aged mice. <u>J Immunol. 166: 1524-30</u>.</li> <li>Bermudez-Fajardo, A. <i>et al.</i> (2011) The effect of Chlamydophila pneumoniae Major Outer Membrane Protein (MOMP) on macrophage and T cell-mediated immune responses. <u>Immunobiology. 216: 152-63</u>.</li> <li>De Jesus, M. <i>et al.</i> (2009) Galactoxylomannan-mediated immune object paralysis results from specific B cell depletion in the context of widespread immune system damage. <u>J Immunol. 183: 3885-94</u>.</li> <li>Jegou, J.F. <i>et al.</i> (2007) C3d binding to the myelin oligodendrocyte glycoprotein results in an exacerbated experimental autoimmune encephalomyelitis. <u>J Immunol. 178: 3323-31</u>.</li> <li>Starck, J. <i>et al.</i> (2010) Inducible Fli-1 gene deletion in adult mice modifies several myeloid lineage commitment decisions and accelerates proliferation arrest and terminal erythrocytic differentiation. <u>Blood. 116: 4795-805</u>.</li> <li>Stotdand, R.S. <i>et al.</i> (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. <u>Blood. 118 (22): 5918-27</u>.</li> <li>White, H.N. and Meng, Q.H. (2012) Recruitment of a Distinct but Related Set of VH Sequences into the Murine CD21hi/CD23- Marginal Zone B Cell Repertoire to That Seen in the Class-Switched Antibody Response. JImmunol. 188: 287-93.</li> <li>Reynaud, J.M. <i>et al.</i> (2014) Human herpesvirus 6A infection in CD46 transgenic mice: viral persistence in the brain and increased production of proinflammatory chemokines via Toll-like receptor 9. J Virol. 88: 5421-36.</li> <li>Candolfi, M. <i>et al.</i> (2014) Anti-inflammatory activity of bone morphogenetic protein signaling pathways in s</li></ol>				
Storage	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.				

Storage

Prior to reconstitution store at +4 $^{\circ}$ C. Following reconstitution store at +4 $^{\circ}$ C.

## DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life12 months from date of reconstitution.						
Health And Safety Information		Material Safety Datasheet documentation #10075 available at: 10075: <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf</u>				df
Regulatory For research p		For research pu	rposes o	nly		
North & South America	Fax: +1 919 878 3	751	rldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
	Email: antibody_sa	ales_us@bio-rad.com	1	Email: antibody_sales_uk@bio 'M341508:190109'	-rad.com	Email: antibody_sales_de@bio-rad.com
			1	Printed on 11 Jan 2019		

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint