

Datasheet: MCA1439FA

Description:	RAT ANTI MOUSE CD19:FITC
Specificity:	CD19
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	6D5
Isotype:	lgG2a
Quantity:	50 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse				
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	FITC	490	525		
Preparation	Purified IgG prepa	red by affinity chromatog	raphy on Protein G fro	m tissue culture supernatant	
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azi	de			

Approx. Protein
Concentrations

IgG concentration 0.1 mg/ml

Human K562 cell line transfected with murine CD19.

Bovine Serum Albumin

External Database Links

Stabilisers

UniProt:

1%

P25918 Related reagents

Entrez Gene:

12478 Cd19 Related reagents

Fusion Partners Spleen cells from immunised rats were fused with cells of the P3X63.Ag8.653 myeloma cell line. **Specificity** Rat anti Mouse CD19 antibody, clone 6D5 recognizes the murine CD19 cell surface antigen, a 95kDa glycoprotein expressed by B lymphocytes. Clone 6D5 has been shown to recognize the same epitope as clone ID3 in cross-competition assays. Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul. The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity Fc receptors. This may be reduced by using SeroBlock FcR (BUF041A/B). References 1. Vernooy, J.H. et al. (2002) Long-term intratracheal lipopolysaccharide exposure in mice results in chronic lung inflammation and persistent pathology. Am J Respir Cell Mol Biol. 26 (1): 152-9. 2. Andrew, D. and Aspinall. R. (2001) II-7 and not stem cell factor reverses both the increase in apoptosis and the decline in thymopoiesis seen in aged mice. J Immunol. 166: 1524-30. 3. Bermudez-Fajardo, A. et al. (2011) The effect of Chlamydophila pneumoniae Major Outer Membrane Protein (MOMP) on macrophage and T cell-mediated immune responses. Immunobiology. 216: 152-63. 4. De Jesus, M. et al. (2009) Galactoxylomannan-mediated immunological paralysis results from specific B cell depletion in the context of widespread immune system damage. J Immunol. 183: 38<u>85-94.</u> 5. Jégou, J.F. et al. (2007) C3d binding to the myelin oligodendrocyte glycoprotein results in an exacerbated experimental autoimmune encephalomyelitis. J Immunol. 178: 3323-31. 6. Starck, J. et al. (2010) Inducible Fli-1 gene deletion in adult mice modifies several myeloid lineage commitment decisions and accelerates proliferation arrest and terminal erythrocytic differentiation. Blood. 116: 4795-805. 7. Scotland, R.S. et al. (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. Blood. 118 (22): 5918-27. 8. White, H.N. and Meng, Q.H. (2012) Recruitment of a Distinct but Related Set of VH Sequences into the Murine CD21hi/CD23- Marginal Zone B Cell Repertoire to That Seen in the Class-Switched Antibody Response. J Immunol. 188: 287-93. 9. Reynaud, J.M. et al. (2014) Human herpesvirus 6A infection in CD46 transgenic mice: viral persistence in the brain and increased production of proinflammatory chemokines via Toll-like receptor 9. J Virol. 88: 5421-36. 10. Candolfi, M. et al. (2011) B cells are critical to T-cell-mediated antitumor immunity induced by a combined immune-stimulatory/conditionally cytotoxic therapy for glioblastoma. Neoplasia. 13: 947-60. 11. Takabayashi, H. et al. (2014) Anti-inflammatory activity of bone morphogenetic protein signaling pathways in stomachs of mice. Gastroenterology. 147: 396-406.e7. 12. Weiss-Gayet, M. et al. (2016) Notch Stimulates Both Self-Renewal and Lineage Plasticity in a Subset of Murine CD9High Committed Megakaryocytic Progenitors. PLoS One. 11 (4): e0153860. 13. Meng, Q.H. & White, H.N. (2017) CD21^{int} CD23⁺ follicular B cells express antigen-specific secretory IgM mRNA as primary and memory responses. Immunology. 151 (2): 211-8. 14. Baer, A. et al. (2016) Protein Phosphatase-1 regulates Rift Valley fever virus replication. Antiviral Res. 127: 79-89. 15. Mccubbrey, A.L. et al. (2016) MicroRNA-34a Negatively Regulates Efferocytosis by Tissue

Storage

Store at +4°C or at -20°C if preferred.

Macrophages in Part via SIRT1. J Immunol. 196 (3): 1366-75.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

RAT IgG2a NEGATIVE CONTROL:FITC (MCA1212F)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 **Europe** Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

'M322762:180727'

Printed on 10 Oct 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint