

Datasheet: MCA1439GA

Description:	RAT ANTI MOUSE CD19
Specificity:	CD19
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	6D5
Isotype:	IgG2a
Quantity:	0.1 mg

Product Details

RRID AB_324453

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Mouse

Product Form Purified IgG - liquid

Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1 mg/ml

Immunogen Human K562 cell line transfected with murine CD19.

**External Database
Links**

UniProt:

[P25918](#) [Related reagents](#)

Entrez Gene:

[12478](#) Cd19 [Related reagents](#)

Fusion Partners Spleen cells from immunised rats were fused with cells of the P3X63.Ag8.653 myeloma cell line.

Specificity **Rat anti Mouse CD19 antibody, clone 6D5** recognizes the murine CD19 cell surface antigen, a 95kDa glycoprotein expressed by B lymphocytes. Clone 6D5 has been shown to recognize the same epitope as clone ID3 in cross-competition assays.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Vernooij, J.H. *et al.* (2002) Long-term intratracheal lipopolysaccharide exposure in mice results in chronic lung inflammation and persistent pathology. [Am J Respir Cell Mol Biol. 26 \(1\): 152-9.](#)
2. Andrew, D. and Aspinall, R. (2001) Il-7 and not stem cell factor reverses both the increase in apoptosis and the decline in thymopoiesis seen in aged mice. [J Immunol. 166: 1524-30.](#)
3. Bermudez-Fajardo, A. *et al.* (2011) The effect of Chlamydomydia pneumoniae Major Outer Membrane Protein (MOMP) on macrophage and T cell-mediated immune responses. [Immunobiology. 216: 152-63.](#)
4. De Jesus, M. *et al.* (2009) Galactoxylomannan-mediated immunological paralysis results from specific B cell depletion in the context of widespread immune system damage. [J Immunol. 183: 3885-94.](#)
5. Jégou, J.F. *et al.* (2007) C3d binding to the myelin oligodendrocyte glycoprotein results in an exacerbated experimental autoimmune encephalomyelitis. [J Immunol. 178: 3323-31.](#)
6. Starck, J. *et al.* (2010) Inducible Fli-1 gene deletion in adult mice modifies several myeloid lineage commitment decisions and accelerates proliferation arrest and terminal erythrocytic differentiation. [Blood. 116: 4795-805.](#)
7. Scotland, R.S. *et al.* (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. [Blood. 118 \(22\): 5918-27.](#)
8. White, H.N. and Meng, Q.H. (2012) Recruitment of a Distinct but Related Set of VH Sequences into the Murine CD21hi/CD23- Marginal Zone B Cell Repertoire to That Seen in the Class-Switched Antibody Response. [J Immunol. 188: 287-93.](#)
9. Reynaud, J.M. *et al.* (2014) Human herpesvirus 6A infection in CD46 transgenic mice: viral persistence in the brain and increased production of proinflammatory chemokines via Toll-like receptor 9. [J Virol. 88: 5421-36.](#)
10. Candolfi, M. *et al.* (2011) B cells are critical to T-cell-mediated antitumor immunity induced by a combined immune-stimulatory/conditionally cytotoxic therapy for glioblastoma. [Neoplasia. 13: 947-60.](#)
11. Takabayashi, H. *et al.* (2014) Anti-inflammatory activity of bone morphogenetic protein signaling pathways in stomachs of mice. [Gastroenterology. 147: 396-406.e7.](#)
12. Weiss-Gayet, M. *et al.* (2016) Notch Stimulates Both Self-Renewal and Lineage Plasticity in a Subset of Murine CD9High Committed Megakaryocytic Progenitors. [PLoS One. 11 \(4\): e0153860.](#)
13. Meng, Q.H. & White, H.N. (2017) CD21^{int} CD23⁺ follicular B cells express antigen-specific secretory IgM mRNA as primary and memory responses. [Immunology. 151 \(2\): 211-8.](#)
14. Baer, A. *et al.* (2016) Protein Phosphatase-1 regulates Rift Valley fever virus replication. [Antiviral Res. 127: 79-89.](#)
15. Mccubbrey, A.L. *et al.* (2016) MicroRNA-34a Negatively Regulates Efferocytosis by Tissue Macrophages in Part via SIRT1. [J Immunol. 196 \(3\): 1366-75.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	18 months from date of despatch.
------------------	----------------------------------

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (STAR72...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight@800
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@800

Recommended Negative Controls

[RAT IgG2a NEGATIVE CONTROL \(MCA1212\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M341182:190109'

Printed on 28 May 2019