

Datasheet: MCA1439PE

## **BATCH NUMBER 151853**

Description:	RAT ANTI MOUSE CD19:RPE	
Specificity:	CD19	
Format:	RPE	
<b>Product Type:</b>	Monoclonal Antibody	
Clone:	6D5	
Isotype:	lgG2a	
Quantity:	100 TESTS	

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified IgG conjuga	ted to R. Phycoerythrin	(RPE) - lyophilized
Reconstitution	Reconstitute with 1 n	nl distilled water	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	RPE 488nm laser	496	578
Preparation	Purified IgG prepared supernatant	d by affinity chromatog	raphy on Protein G from tissue cultu
Buffer Solution	Phosphate buffered s	saline	
Preservative Stabilisers	0.09% Sodium Azide		
	1% Bovine Serum Albumin		
	5% Sucrose		

Immunogen	Human K562 cell line transfected with murine CD19.
External Database Links	UniProt: P25918 Related reagents
	Entrez Gene:
	12478 Cd19 Related reagents
RRID	AB_321344
Fusion Partners	Spleen cells from immunised rats were fused with cells of the P3X63.Ag8.653 myeloma cell line.
Specificity	Rat anti Mouse CD19 antibody, clone 6D5 recognizes the murine CD19 cell surface antigen, a ~95 kDa glycoprotein expressed by B lymphocytes. Rat anti Mouse CD19 antibody, clone 6D5 recognizes the same, or a closely related epitope as clone Rat anti Mouse antibody, clone ID3 in cross-competition assays.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
	The Fc region of monoclonal antibodies may bind non-specifically to cells expressing low affinity fc receptors. This may be reduced by using SeroBlock FcR ( <u>BUF041A/B</u> ).
References	<ol> <li>Vernooy, J.H. <i>et al.</i> (2002) Long-term intratracheal lipopolysaccharide exposure in mice results in chronic lung inflammation and persistent pathology. <u>Am J Respir Cell Mol Biol.</u> 26 (1): 152-9.</li> <li>Andrew, D. and Aspinall. R. (2001) II-7 and not stem cell factor reverses both the</li> </ol>
	increase in apoptosis and the decline in thymopoiesis seen in aged mice. <u>J Immunol. 166:</u> 1524-30.
	3. Bermudez-Fajardo, A. et al. (2011) The effect of Chlamydophila pneumoniae Major
	Outer Membrane Protein (MOMP) on macrophage and T cell-mediated immune
	responses. Immunobiology. 216: 152-63.
	4. De Jesus, M. <i>et al.</i> (2009) Galactoxylomannan-mediated immunological paralysis

- 4. De Jesus, M. et al. (2009) Galactoxylomannan-mediated immunological paralysis results from specific B cell depletion in the context of widespread immune system
- damage. J Immunol. 183: 3885-94. 5. Jégou, J.F. et al. (2007) C3d binding to the myelin oligodendrocyte glycoprotein results
- in an exacerbated experimental autoimmune encephalomyelitis. J Immunol. 178: 3323-31.
- 6. Starck, J. et al. (2010) Inducible Fli-1 gene deletion in adult mice modifies several myeloid lineage commitment decisions and accelerates proliferation arrest and terminal erythrocytic differentiation. Blood. 116: 4795-805.
- 7. Scotland, R.S. et al. (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. <u>Blood. 118 (22): 5918-27.</u>
- 8. White, H.N. and Meng, Q.H. (2012) Recruitment of a Distinct but Related Set of VH Sequences into the Murine CD21hi/CD23- Marginal Zone B Cell Repertoire to That Seen in the Class-Switched Antibody Response. J Immunol. 188: 287-93.
- 9. Reynaud, J.M. et al. (2014) Human herpesvirus 6A infection in CD46 transgenic mice: viral persistence in the brain and increased production of proinflammatory chemokines via

Toll-like receptor 9. J Virol. 88: 5421-36.

- 10. Candolfi, M. *et al.* (2011) B cells are critical to T-cell-mediated antitumor immunity induced by a combined immune-stimulatory/conditionally cytotoxic therapy for glioblastoma. Neoplasia. 13: 947-60.
- 11. Takabayashi, H. *et al.* (2014) Anti-inflammatory activity of bone morphogenetic protein signaling pathways in stomachs of mice. <u>Gastroenterology. 147: 396-406.e7.</u>
- 12. Weiss-Gayet, M. *et al.* (2016) Notch Stimulates Both Self-Renewal and Lineage Plasticity in a Subset of Murine CD9High Committed Megakaryocytic Progenitors. <u>PLoS One. 11 (4): e0153860.</u>
- 13. Meng, Q.H. & White, H.N. (2017) CD21<sup>int</sup> CD23<sup>+</sup> follicular B cells express antigen-specific secretory IgM mRNA as primary and memory responses. <u>Immunology. 151 (2):</u> 211-8.
- 14. Mccubbrey, A.L. *et al.* (2016) MicroRNA-34a Negatively Regulates Efferocytosis by Tissue Macrophages in Part via SIRT1. <u>J Immunol. 196 (3): 1366-75.</u>

#### Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

#### DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #20487 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1439PE">https://www.bio-rad-antibodies.com/SDS/MCA1439PE</a> 20487	
Regulatory	For research purposes only	

# Related Products

### **Recommended Negative Controls**

RAT IgG2a NEGATIVE CONTROL:RPE (MCA1212PE)

#### **Recommended Useful Reagents**

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375326:210104'

### Printed on 20 Mar 2024