

Datasheet: MCA1439SBB675

Description:	RAT ANTI MOUSE CD19:StarBright Blue 675
Specificity:	CD19
Format:	StarBright Blue 675
Product Type:	Monoclonal Antibody
Clone:	6D5
Isotype:	IgG2a
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species Product Form	Mouse Purified IgG conjugate	ed to StarBright Blue 6	675 - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	StarBright Blue 675	476	675
eparation	Purified IgG prepared	by affinity chromatog	raphy on Protein G
	supernatant	by animaly contentations	,
fer Solution	•		
	supernatant	aline	
reservative	supernatant Phosphate buffered sa	aline NaN ₃)	
reservative	Phosphate buffered so	aline NaN ₃)	
Buffer Solution Preservative Stabilisers	Phosphate buffered sa 0.09% sodium azide (1% bovine serum albu	aline NaN ₃)	

Immunogen	lm	m	un	OQ	en
------------------	----	---	----	----	----

Human K562 cell line transfected with murine CD19.

External Database Links

UniProt:

P25918 Related reagents

Entrez Gene:

12478 Cd19 Related reagents

Fusion Partners

Spleen cells from immunised rats were fused with cells of the P3X63.Ag8.653 myeloma cell line.

Specificity

Rat anti Mouse CD19 antibody, clone 6D5 recognizes the murine CD19 cell surface antigen, a ~95 kDa glycoprotein expressed by B lymphocytes. Rat anti Mouse CD19 antibody, clone 6D5 recognizes the same, or a closely related epitope as clone Rat anti Mouse CD19 antibody, clone ID3 in cross-competition assays.

Flow Cytometry

Use 5μ I of the suggested working dilution to label 10^6 cells in 100ul. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Vernooy, J.H. *et al.* (2002) Long-term intratracheal lipopolysaccharide exposure in mice results in chronic lung inflammation and persistent pathology. <u>Am J Respir Cell Mol Biol.</u> 26 (1): 152-9.
- 2. Andrew, D. and Aspinall. R. (2001) II-7 and not stem cell factor reverses both the increase in apoptosis and the decline in thymopoiesis seen in aged mice. <u>J Immunol. 166:</u> 1524-30.
- 3. Bermudez-Fajardo, A. *et al.* (2011) The effect of *Chlamydophila pneumoniae* Major Outer Membrane Protein (MOMP) on macrophage and T cell-mediated immune responses. Immunobiology. 216: 152-63.
- 4. De Jesus, M. *et al.* (2009) Galactoxylomannan-mediated immunological paralysis results from specific B cell depletion in the context of widespread immune system damage. <u>J Immunol.</u> 183: 3885-94.
- 5. Jégou, J.F. *et al.* (2007) C3d binding to the myelin oligodendrocyte glycoprotein results in an exacerbated experimental autoimmune encephalomyelitis. <u>J Immunol</u>. 178: 3323-31.
- 6. Starck, J. *et al.* (2010) Inducible Fli-1 gene deletion in adult mice modifies several myeloid lineage commitment decisions and accelerates proliferation arrest and terminal erythrocytic differentiation. <u>Blood. 116: 4795-805.</u>
- 7. Scotland, R.S. *et al.* (2011) Sex differences in resident immune cell phenotype underlie more efficient acute inflammatory responses in female mice. <u>Blood.</u> 118 (22): 5918-27.
- 8. White, H.N. and Meng, Q.H. (2012) Recruitment of a Distinct but Related Set of VH Sequences into the Murine CD21hi/CD23- Marginal Zone B Cell Repertoire to That Seen in the Class-Switched Antibody Response. <u>J Immunol. 188: 287-93.</u>
- 9. Reynaud, J.M. *et al.* (2014) Human herpesvirus 6A infection in CD46 transgenic mice: viral persistence in the brain and increased production of proinflammatory chemokines via Toll-like receptor 9. J Virol. 88: 5421-36.
- 10. Candolfi, M. *et al.* (2011) B cells are critical to T-cell-mediated antitumor immunity induced by a combined immune-stimulatory/conditionally cytotoxic therapy for glioblastoma. Neoplasia. 13: 947-60.

- 11. Takabayashi, H. *et al.* (2014) Anti-inflammatory activity of bone morphogenetic protein signaling pathways in stomachs of mice. <u>Gastroenterology</u>. 147: 396-406.e7.
- 12. Weiss-Gayet, M. *et al.* (2016) Notch Stimulates Both Self-Renewal and Lineage Plasticity in a Subset of Murine CD9High Committed Megakaryocytic Progenitors. <u>PLoS One. 11 (4): e0153860.</u>
- 13. Meng, Q.H. & White, H.N. (2017) CD21^{int} CD23⁺ follicular B cells express antigen-specific secretory IgM mRNA as primary and memory responses. <u>Immunology. 151 (2):</u> 211-8.
- 14. Mccubbrey, A.L. *et al.* (2016) MicroRNA-34a Negatively Regulates Efferocytosis by Tissue Macrophages in Part via SIRT1. <u>J Immunol</u>. 196 (3): 1366-75.
- 15. Vila-Caballer, M. *et al.* (2019) Disruption of the CCL1-CCR8 axis inhibits vascular Treg recruitment and function and promotes atherosclerosis in mice. <u>J Mol Cell Cardiol. 132:</u> 154-63.
- 16. Domingues, C.S. *et al.* (2020) Host Genetics Background Influence in the Intragastric *Trypanosoma cruzi* Infection. <u>Front Immunol. 11: 566476.</u>

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.	
Guarantee	12 months from date of despatch	
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U. counterparts	S. and foreign
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1439SBB675 20471	
Regulatory	For research purposes only	

Related Products

Recommended Useful Reagents

MOUSE SEROBLOCK FcR (BUF041A)
MOUSE SEROBLOCK FcR (BUF041B)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M409709:221020'

Printed on 10 May 2024