

Datasheet: MCA1449F

Description:	RAT ANTI PIG CD44:FITC
Specificity:	CD44
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	MAC329
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Pig		
Species Cross Reactivity	Reacts with: Dog, Horse, Bovine, Goat, Sheep N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein A		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	Purified porcine CD44		
Fusion Partners	Spleen cells from immunised LOU Rats were fused with cells of the Y3.Ag1.2.3 myeloma cell line		

Specificity	Rat anti Pig CD44 antibody, clone MAC329 recognizes the porcine CD44 cell surface antigen which is involved in cell-cell interactions, cell adhesion and migration. CD44 is also a receptor for hyaluronic acid and can interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells or 100ul whole blood
References	<ol style="list-style-type: none"> 1. Yang, H. <i>et al.</i> (1993) Preparation and reactivities of anti-porcine CD44 monoclonal antibodies. Scand J Immunol. 37 (4): 490-8. 2. Kasper, M. <i>et al.</i> (1996) Expression of CD44 isoforms during bleomycin-or radiation-induced pulmonary fibrosis in rats and mini-pigs. Histochem Cell Biol. 105 (3): 221-30. 3. Gyongyosi, M. <i>et al.</i> (2010) Differential effect of ischaemic preconditioning on mobilisation and recruitment of haematopoietic and mesenchymal stem cells in porcine myocardial ischaemia-reperfusion. Thromb Haemost. 103: 1-9. 4. Smith, M.M. <i>et al.</i> (2008) Significant synovial pathology in a meniscectomy model of osteoarthritis: modification by intra-articular hyaluronan therapy. Rheumatology (Oxford). 47:1172-8. 5. Sanchez, M.A. <i>et al.</i> (2004) Organ-specific immunity in canine visceral leishmaniasis: analysis of symptomatic and asymptomatic dogs naturally infected with Leishmania chagasi. Am J Trop Med Hyg. 70: 618-24.
Further Reading	1. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	18 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: 10041: https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

[RAT IgG1 NEGATIVE CONTROL:FITC \(MCA6004F\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M333716:181119'

Printed on 20 May 2019