

Datasheet: MCA1456

**BATCH NUMBER 162873**

<b>Description:</b>	MOUSE ANTI HUMAN CD104
<b>Specificity:</b>	CD104
<b>Other names:</b>	INTEGRIN BETA 4
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	450-9D
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/100
Immunohistology - Frozen (1)	▪			1/500 - 1/1000
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	<p>Reacts with: Ferret, Mink</p> <p>Based on sequence similarity, is expected to react with: Mustelid</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or</p>

personal communications from the originators. Please refer to references indicated for further information.

<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Purified alpha6 beta4 integrin from A431 cells.
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P16144</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">3691</a> ITGB4 <a href="#">Related reagents</a>
<b>RRID</b>	AB_2129143
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/0 myeloma cell line.
<b>Specificity</b>	<b>Mouse anti Human CD104 antibody, clone 450-9D</b> recognizes the human beta4 integrin, also known as CD104. CD104 is a ~205 kDa glycoprotein which associates with the alpha6 integrin to form the alpha6/beta4 complex. CD104 is expressed on epithelial cells, Schwann cells and various tumor cell lines.  Mouse anti Human CD104 antibody, clone 450-9D recognizes an extracellular epitope on the CD104 molecule.
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>Histology Positive Control Tissue</b>	Human Tonsil
<b>References</b>	1. Kennel, S.J. <i>et al.</i> (1990) Second generation monoclonal antibodies to the human integrin alpha 6 beta 4. <a href="#">Hybridoma. 9 (3): 243-55.</a> 2. Barbaro, V. <i>et al.</i> (2009) Reconstruction of a human hemicornea through natural scaffolds compatible with the growth of corneal epithelial stem cells and stromal

keratocytes. [Mol Vis. 15: 2084-93.](#)

3. Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. [Vet Immunol Immunopathol. 132:109-15.](#)
4. Smith, K. *et al.* (2011) Mono- and tri-cationic porphyrin-monoclonal antibody conjugates: photodynamic activity and mechanism of action. [Immunology. 132: 256-65.](#)
5. Alam, H. *et al.* (2011) Loss of keratins 8 and 18 leads to alterations in  $\alpha 6\beta 4$ -integrin-mediated signalling and decreased neoplastic progression in an oral-tumour-derived cell line. [J Cell Sci. 124: 2096-106.](#)
6. Lai-cheong, J.E. *et al.* (2009) Loss-of-function FERMT1 mutations in kindler syndrome implicate a role for fermitin family homolog-1 in integrin activation. [Am J Pathol. 175 \(4\): 1431-41.](#)
7. Groves, R.W. *et al.* (2010) A homozygous nonsense mutation within the dystonin gene coding for the coiled-coil domain of the epithelial isoform of BPAG1 underlies a new subtype of autosomal recessive epidermolysis bullosa simplex. [J Invest Dermatol. 130 \(6\): 1551-7.](#)
8. Aasted, B. & Viuff, B. (2007) Reactivity of monoclonal antibodies to human CD antigens with cells from mink. [Vet Immunol Immunopathol. 119 \(1-2\): 27-37.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** 12 months from date of despatch

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1456>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

- |   |   |
|---|---|
| Rabbit Anti Mouse IgG (STAR12...)       | <a href="#">RPE</a>   |
| Goat Anti Mouse IgG IgA IgM (STAR87...) | <a href="#">HRP</a>   |
| Goat Anti Mouse IgG (STAR76...)         | <a href="#">RPE</a>   |
| Rabbit Anti Mouse IgG (STAR13...)       | <a href="#">HRP</a>   |
| Goat Anti Mouse IgG (STAR70...)         | <a href="#">FITC</a>  |
| Goat Anti Mouse IgG (H/L) (STAR117...)  | <a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> ,<br><a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> ,<br><a href="#">FITC</a> , <a href="#">HRP</a> |
| Rabbit Anti Mouse IgG (STAR9...)        | <a href="#">FITC</a>  |
| Goat Anti Mouse IgG (STAR77...)         | <a href="#">HRP</a>   |
| Goat Anti Mouse IgG (Fc) (STAR120...)   | <a href="#">FITC</a> , <a href="#">HRP</a>  |

## Recommended Negative Controls

### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M383051:210513'

**Printed on 18 Jan 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)