

## Datasheet: MCA1456

<b>Description:</b>	MOUSE ANTI HUMAN CD104
<b>Specificity:</b>	CD104
<b>Other names:</b>	INTEGRIN BETA 4
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	450-9D
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	0.2 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/100
Immunohistology - Frozen (1)	▪			1/500 - 1/1000
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Ferret, Mink Based on sequence similarity, is expected to react with:Mustelid <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide

<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Purified alpha6 beta4 integrin from A431 cells.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P16144</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3691</a>   ITGB4   <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_2129143
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse Sp2/0 myeloma cell line.
<b>Specificity</b>	<p><b>Mouse anti Human CD104 antibody, clone 450-9D</b> recognizes the human beta4 integrin, also known as CD104. CD104 is a ~205 kDa glycoprotein which associates with the alpha6 integrin to form the alpha6/beta4 complex. CD104 is expressed on epithelial cells, Schwann cells and various tumor cell lines.</p> <p>Mouse anti Human CD104 antibody, clone 450-9D recognizes an extracellular epitope on the CD104 molecule.</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>Histology Positive Control Tissue</b>	Human Tonsil
<b>References</b>	<ol style="list-style-type: none"> <li>1. Kennel, S.J. <i>et al.</i> (1990) Second generation monoclonal antibodies to the human integrin alpha 6 beta 4. <a href="#">Hybridoma. 9 (3): 243-55.</a></li> <li>2. Barbaro, V. <i>et al.</i> (2009) Reconstruction of a human hemiacornea through natural scaffolds compatible with the growth of corneal epithelial stem cells and stromal keratocytes. <a href="#">Mol Vis. 15: 2084-93.</a></li> <li>3. Martel, C.J. &amp; Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. <a href="#">Vet Immunol Immunopathol. 132:109-15.</a></li> <li>4. Smith, K. <i>et al.</i> (2011) Mono- and tri-cationic porphyrin-monoclonal antibody conjugates: photodynamic activity and mechanism of action. <a href="#">Immunology. 132: 256-65.</a></li> <li>5. Alam, H. <i>et al.</i> (2011) Loss of keratins 8 and 18 leads to alterations in α6β4-integrin-mediated signalling and decreased neoplastic progression in an oral-tumour-derived cell line. <a href="#">J Cell Sci. 124: 2096-106.</a></li> <li>6. Lai-cheong, J.E. <i>et al.</i> (2009) Loss-of-function FERMT1 mutations in kindler syndrome implicate a role for fermitin family homolog-1 in integrin activation. <a href="#">Am J Pathol. 175 (4): 1431-41.</a></li> <li>7. Groves, R.W. <i>et al.</i> (2010) A homozygous nonsense mutation within the dystonin gene coding for the coiled-coil domain of the epithelial isoform of BPAG1 underlies a new subtype of autosomal recessive epidermolysis bullosa simplex. <a href="#">J Invest Dermatol. 130 (6): 1551-7.</a></li> <li>8. Aasted, B. &amp; Viuff, B. (2007) Reactivity of monoclonal antibodies to human CD antigens with cells from mink. <a href="#">Vet Immunol Immunopathol. 119 (1-2): 27-37.</a></li> </ol>
<b>Storage</b>	Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">Alk. Phos.</a> , <a href="#">HRP</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Rabbit Anti Mouse IgG (STAR8...)	<a href="#">DyLight@800</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight@488</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M365339:200529'

Printed on 11 Aug 2020