

# Datasheet: MCA1457F BATCH NUMBER 151035

MOUSE ANTI HUMAN CD49f:FITC
CD49f
INTEGRIN ALPHA 6 CHAIN, VLA-6
FITC
Monoclonal Antibody
450-30A
lgG1
0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes N	o Not Dete	rmined	Suggested Dilution		
	Flow Cytometry	-			Neat		
	Immunohistology - Frozei	n	•				
	Immunohistology - Paraff	in	-	•			
	Where this antibody han necessarily exclude its a guide only. It is recon system using appropria	use in such pro nmended that th	cedures. Sugges e user titrates the	ted working	dilutions are given as		
Target Species	Human						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	Fluorophore	Excitation Max	nm) Emission N	/lax (nm)			
	FITC	490	525				
Preparation	Purified IgG prepared by affinity chromatography on Protein G supernatant				n tissue culture		
Buffer Solution	Phosphate buffered sa	line					
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum A	Albumin					

Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	alpha 6 beta 4 integrin purified from A431 cells.
External Database Links	UniProt: <u>P23229</u> <u>Related reagents</u> Entrez Gene: <u>3655</u> ITGA6 <u>Related reagents</u>
RRID	AB_321465
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
Specificity	<ul><li>Mouse anti Human CD49f antibody, clone 450-30A recognizes the human VLA-6 cell surface antigen, also known as the alpha 6 integrin and as CD49f.</li><li>CD49f is expressed by platelets, weakly by monocytes and by a subset of lymphocytes.</li><li>CD49f is also widely expressed on epithelial tissues.</li></ul>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
References	<ol> <li>Kennel, S.J. <i>et al.</i> (1990) Second generation monoclonal antibodies to the human integrin alpha 6 beta 4. <u>Hybridoma. 9 (3): 243-55.</u></li> <li>Maurice, S. <i>et al.</i> (2007) Isolation of progenitor cells from cord blood using adhesion matrices. <u>Cytotechnology. 54: 121-33.</u></li> <li>Cavers, M. <i>et al.</i> (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. <u>Clin Exp Immunol. 127: 60-5.</u></li> <li>López, J. <i>et al.</i> (2012) Cancer-initiating cells derived from established cervical cell lines exhibit stem-cell markers and increased radioresistance. <u>BMC Cancer. 12: 48.</u></li> <li>Keller, P.J. <i>et al.</i> (2012) Defining the cellular precursors to human breast cancer. <u>Proc Natl Acad Sci U S A. 109: 2772-7.</u></li> <li>Kaczmarek, M. <i>et al.</i> (2011) Evaluation of the phenotype pattern of macrophages isolated from malignant and non-malignant pleural effusions. <u>Tumour Biol. 32: 1123-32.</u></li> <li>Aldridge, V. <i>et al.</i> (2012) NX17 protects integrins from degradation by sorting between lysosomal and recycling pathways. <u>J Cell Biol. 197 (2): 219-30.</u></li> <li>Liu, L. <i>et al.</i> (2003) Priming of eosinophil migration across lung epithelial cell monolayers and upregulation of CD11b/CD18 are elicited by extracellular Ca<sup>2+.</sup> <u>Am J Respir Cell Mol Biol. 28: 713-21.</u></li> <li>Goyer, B. <i>et al.</i> (2017) Extracellular matrix and integrin expression profiles in Fuchs</li> </ol>

	endothelial corneal dystrophy cells and tissue model. <u>Tissue Eng Part A. Jul 20 [Epub</u> <u>ahead of print].</u>				
Storage	Store at +4°C or at -20°C if preferred.				
	This product should be stored undiluted.				
	Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.				
	Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.				
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1457F 10041				
Regulatory	For research purposes only				

## Related Products

### **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL: FITC (MCA928F)

### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rac	l.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365344:200529'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint