

Datasheet: MCA1457F

Description:	MOUSE ANTI HUMAN CD49f:FITC
Specificity:	CD49f
Other names:	INTEGRIN ALPHA 6 CHAIN, VLA-6
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	450-30A
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant		
Buffer Solution	Phosphate buffered saline		
Preservative	0.09% Sodium Azide		
Stabilisers	1% Bovine Serum Albumin		
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml		
Immunogen	alpha 6 beta 4 integrin purified from A431 cells.		
External Database Links	UniProt:		

Entrez Gene:

[3655](#) ITGA6 [Related reagents](#)

RRID	AB_321465
-------------	-----------

Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.
------------------------	---

Specificity	<p>Mouse anti Human CD49f antibody, clone 450-30A recognizes the human VLA-6 cell surface antigen, also known as the alpha 6 integrin and as CD49f.</p> <p>CD49f is expressed by platelets, weakly by monocytes and by a subset of lymphocytes.</p> <p>CD49f is also widely expressed on epithelial tissues.</p>
--------------------	---

Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
-----------------------	---

References	<ol style="list-style-type: none">1. Kennel, S.J. <i>et al.</i> (1990) Second generation monoclonal antibodies to the human integrin alpha 6 beta 4. Hybridoma. 9 (3): 243-55.2. Maurice, S. <i>et al.</i> (2007) Isolation of progenitor cells from cord blood using adhesion matrices. Cytotechnology. 54: 121-33.3. Cavers, M. <i>et al.</i> (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. Clin Exp Immunol. 127: 60-5.4. López, J. <i>et al.</i> (2012) Cancer-initiating cells derived from established cervical cell lines exhibit stem-cell markers and increased radioresistance. BMC Cancer. 12: 48.5. Keller, P.J. <i>et al.</i> (2012) Defining the cellular precursors to human breast cancer. Proc Natl Acad Sci U S A. 109: 2772-7.6. Kaczmarek, M. <i>et al.</i> (2011) Evaluation of the phenotype pattern of macrophages isolated from malignant and non-malignant pleural effusions. Tumour Biol. 32: 1123-32.7. Aldridge, V. <i>et al.</i> (2012) Human mesenchymal stem cells are recruited to injured liver in a β1-integrin and CD44 dependent manner. Hepatology. 56 (3): 1063-73.8. Steinberg, F. <i>et al.</i> (2012) SNX17 protects integrins from degradation by sorting between lysosomal and recycling pathways. J Cell Biol. 197 (2): 219-30.9. Liu, L. <i>et al.</i> (2003) Priming of eosinophil migration across lung epithelial cell monolayers and upregulation of CD11b/CD18 are elicited by extracellular Ca²⁺. Am J Respir Cell Mol Biol. 28: 713-21.10. Goyer, B. <i>et al.</i> (2017) Extracellular matrix and integrin expression profiles in Fuchs endothelial corneal dystrophy cells and tissue model. Tissue Eng Part A. Jul 20 [Epub ahead of print].
-------------------	---

Storage	<p>Store at +4°C or at -20°C if preferred.</p> <p>This product should be stored undiluted.</p> <p>Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
----------------	--

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M365344:200529'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)