

## Datasheet: MCA1457T

<b>Description:</b>	MOUSE ANTI HUMAN CD49f
<b>Specificity:</b>	CD49f
<b>Other names:</b>	INTEGRIN ALPHA 6 CHAIN, VLA-6
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	450-30A
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	25 µg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/10 - 1/50
Immunohistology - Frozen (1)	▪			1/50 - 1/100
Immunohistology - Paraffin		▪		
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.**

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml

**Immunogen** alpha 6 beta 4 integrin purified from A431 cells.

---

**External Database Links**

**UniProt:**  
[P23229](#) [Related reagents](#)

**Entrez Gene:**  
[3655](#) ITGA6 [Related reagents](#)

---

**RRID** AB\_2128304

---

**Fusion Partners** Spleen cells from immunized BALB/c mice were fused with cells of the mouse SP2/0 myeloma cell line.

---

**Specificity** **Mouse anti Human CD49f antibody, clone 450-30A** recognizes the human VLA-6 cell surface antigen, also known as the alpha 6 integrin and as CD49f.

CD49f is expressed by platelets, weakly by monocytes and by a subset of lymphocytes.

CD49f is also widely expressed on epithelial tissues.

---

**Flow Cytometry** Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

---

**Histology Positive Control Tissue** Tonsil

---

**References**

1. Kennel, S.J. *et al.* (1990) Second generation monoclonal antibodies to the human integrin alpha 6 beta 4. [Hybridoma. 9 \(3\): 243-55.](#)
  2. Maurice, S. *et al.* (2007) Isolation of progenitor cells from cord blood using adhesion matrices. [Cytotechnology. 54: 121-33.](#)
  3. Cavers, M. *et al.* (2002) Differential expression of beta1 and beta2 integrins and L-selectin on CD4+ and CD8+ T lymphocytes in human blood: comparative analysis between isolated cells, whole blood samples and cryopreserved preparations. [Clin Exp Immunol. 127: 60-5.](#)
  4. López, J. *et al.* (2012) Cancer-initiating cells derived from established cervical cell lines exhibit stem-cell markers and increased radioresistance. [BMC Cancer. 12: 48.](#)
  5. Keller, P.J. *et al.* (2012) Defining the cellular precursors to human breast cancer. [Proc Natl Acad Sci U S A. 109: 2772-7.](#)
  6. Kaczmarek, M. *et al.* (2011) Evaluation of the phenotype pattern of macrophages isolated from malignant and non-malignant pleural effusions. [Tumour Biol. 32: 1123-32.](#)
  7. Aldridge, V. *et al.* (2012) Human mesenchymal stem cells are recruited to injured liver in a  $\beta$ 1-integrin and CD44 dependent manner. [Hepatology. 56 \(3\): 1063-73.](#)
  8. Steinberg, F. *et al.* (2012) SNX17 protects integrins from degradation by sorting between lysosomal and recycling pathways. [J Cell Biol. 197 \(2\): 219-30.](#)
  9. Liu, L. *et al.* (2003) Priming of eosinophil migration across lung epithelial cell monolayers and upregulation of CD11b/CD18 are elicited by extracellular Ca<sup>2+</sup>. [Am J Respir Cell Mol Biol. 28: 713-21.](#)
  10. Goyer, B. *et al.* (2017) Extracellular matrix and integrin expression profiles in Fuchs endothelial corneal dystrophy cells and tissue model. [Tissue Eng Part A. Jul 20 \[Epub ahead of print\].](#)
- 

**Storage** Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
--------------------------------------	--

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)  
Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)  
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®680](#),  
[DyLight®800](#), [FITC](#), [HRP](#)

### Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M365348:200529'

Printed on 11 Aug 2020

---

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)