

Datasheet: MCA1477

Description:	RAT ANTI HUMAN CD3
Specificity:	CD3
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CD3-12
Isotype:	IgG1
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/50 - 1/100
Immunohistology - Frozen	▪			1/100
Immunohistology - Paraffin (2)	▪			1/100
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting	▪			
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)Membrane permeabilization is required for this application. Bio-Rad recommends the use of Leucoperm (BUF09) for this purpose.

(2)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. Tris/EDTA buffer pH 9.0 is recommended for this purpose.

Target Species

Human

Species Cross Reactivity

Reacts with: Bovine, Dog, Horse, Rhesus Monkey, Pig, Chicken, Mouse, Duck, Koala, Harbour Porpoise, Alpaca, Cynomolgus monkey, Spotted Hyena, Sea Lion, Cat, Amazon Parrot, Raccoon, Great horned owl (*Bubo virginianus*), Bullfrog, Xenopus, Rabbit
Based on sequence similarity, is expected to react with: Mammals, Birds, Amphibia
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Synthetic peptide sequence derived from cytoplasmic epitope of CD3 (Glu-Arg-Pro-Pro-Pro-Val-Pro-Asn-Pro-Asp-Tyr-Glu-Pro-Cys) (ERPPPVPNPDYEPC)
External Database Links	<p>UniProt: P07766 Related reagents</p> <p>Entrez Gene: 916 CD3E Related reagents</p>
Synonyms	T3E
RRID	AB_321245
Specificity	<p>Rat anti Human CD3, clone CD3-12 raised against a peptide representing an invariant cytoplasmic sequence within the CD3ϵ chain recognizes human CD3ϵ. CD3 is a multimeric protein complex composed of four distinct polypeptide chains (ϵ, γ, δ, ζ) that assemble and function as three pairs of dimers ($\epsilon\gamma$, $\epsilon\delta$, $\zeta\zeta$). The CD3 complex serves as a T cell co-receptor that associates non-covalently with the T cell receptor (TCR) (Malissen 2008; Guy and Vignali 2009; Smith-Garvin et al. 2009). CD3 is a defining feature of cells belonging to the T cell lineage and can therefore be used as T cell marker.</p> <p>As Rat anti Human CD3, clone CD3-12 has been specifically raised against an epitope within the epsilon peptide chain, highly conserved among species clone CD3-12 has a very broad species crossreactivity for the CD3 marker. (Jones et al. 1993; Kothlow et al. 2005).</p>
Histology Positive Control Tissue	Tonsil
References	<ol style="list-style-type: none"> Jones, M. <i>et al.</i> (1993) Detection of T and B cells in many animal species using cross-reactive anti-peptide antibodies. J Immunol. 150 (12): 5429-35. Shulga-Morskaya, S. <i>et al.</i> (2004) B cell-activating factor belonging to the TNF family acts through separate receptors to support B cell survival and T cell-independent antibody

- formation. [J Immunol. 173 \(4\): 2331-41.](#)
3. Kapturczak, M.H. *et al.* (2004) Heme oxygenase-1 modulates early inflammatory responses: evidence from the heme oxygenase-1-deficient mouse. [Am J Pathol. 165 \(3\): 1045-53.](#)
 4. Kothlow, S. *et al.* (2005) Characterization of duck leucocytes by monoclonal antibodies. [Dev Comp Immunol. 29 \(8\): 733-48.](#)
 5. Patole, P.S. *et al.* (2006) Expression and regulation of Toll-like receptors in lupus-like immune complex glomerulonephritis of MRL-Fas(lpr) mice. [Nephrol Dial Transplant 21 \(11\): 3062-73.](#)
 6. Foryst-Ludwig, A. *et al.* (2010) PPARgamma activation attenuates T-lymphocyte-dependent inflammation of adipose tissue and development of insulin resistance in obese mice. [Cardiovasc Diabetol. 9: 64.](#)
 7. Osorio, Y. *et al.* (2011) Identification of small molecule lead compounds for visceral leishmaniasis using a novel *ex vivo* splenic explant model system. [PLoS Negl Trop Dis. 5:e962.](#)
 8. Flatz, L. *et al.* (2011) T cell-dependence of Lassa fever pathogenesis. [PLoS Pathog. 6: e1000836.](#)
 9. Gendronneau, G. *et al.* (2010) Influence of Hoxa5 on p53 tumorigenic outcome in mice. [Am J Pathol. 176: 995-1005.](#)
 10. Herrmann, I. *et al.* (2006) *Streptococcus pneumoniae* Infection aggravates experimental autoimmune encephalomyelitis via Toll-like receptor 2. [Infect Immun. 74: 4841-8.](#)
 11. Ruf, M.T. *et al.* (2012) Chemotherapy-Associated Changes of Histopathological Features of *Mycobacterium ulcerans* Lesions in a Buruli Ulcer Mouse Model. [Antimicrob Agents Chemother. 56: 687-96.](#)
 12. Roy, M. *et al.* (2012) CXCL1 can be regulated by IL-6 and promotes granulocyte adhesion to brain capillaries during bacterial toxin exposure and encephalomyelitis. [J Neuroinflammation. 9: 18.](#)
 13. Campuzano, O. *et al.* (2012) Arrhythmogenic right ventricular cardiomyopathy: severe structural alterations are associated with inflammation. [J Clin Pathol. 65 \(12\): 1077-83.](#)
 14. Lau, Q. *et al.* (2012) Expression and *in vitro* upregulation of MHCII in koala lymphocytes. [Vet Immunol Immunopathol. 147: 35-43.](#)
 15. Beineke, A. *et al.* (2007) Phenotypical characterization of changes in thymus and spleen associated with lymphoid depletion in free-ranging harbor porpoises (*Phocoena phocoena*). [Vet Immunol Immunopathol. 117: 254-65.](#)
 16. Pusterla, N. *et al.* (2006) Multicentric T-cell lymphosarcoma in an alpaca. [Vet J. 171: 181-5.](#)
 17. Wiessner, C. *et al.* (2011) The Second-Generation Active A{beta} Immunotherapy CAD106 Reduces Amyloid Accumulation in APP Transgenic Mice While Minimizing Potential Side Effects. [J Neurosci. 31: 9323-31.](#)
 18. Singleton, C.L. *et al.* (2007) Diagnosis and treatment of chronic T-lymphocytic leukemia in a spotted hyena (*Crocuta crocuta*). [J Zoo Wildl Med. 38: 488-91.](#)
 19. Colegrove, K.M. *et al.* (2010) Polyomavirus infection in a free-ranging California sea lion (*Zalophus californianus*) with intestinal T-cell lymphoma. [J Vet Diagn Invest. 22: 628-32.](#)
 20. Steinberg, J.D. and Keating, J.H. (2008) What is your diagnosis? Cervical mass in a cat. [Vet Clin Pathol. 37: 323-7.](#)

21. Osofsky, A. *et al.* (2011) T-cell chronic lymphocytic leukemia in a double yellow-headed Amazon parrot (*Amazona ochrocephala oratrix*). [J Avian Med Surg. 25: 286-94.](#)
22. Giannitti, F. *et al.* (2014) Temporal and geographic clustering of polyomavirus-associated olfactory tumors in 10 free-ranging raccoons (*Procyon lotor*). [Vet Pathol. 51 \(4\): 832-45.](#)
23. Malka, S. *et al.* (2008) Disseminated lymphoma of presumptive T-cell origin in a great horned owl (*Bubo virginianus*). [J Avian Med Surg. 22: 226-33.](#)
24. Bricker, N.K. *et al.* (2012) Cytochemical and immunocytochemical characterization of blood cells and immunohistochemical analysis of spleen cells from 2 species of frog, *Rana (Aquarana) catesbeiana* and *Xenopus laevis*. [Vet Clin Pathol. 41: 353-61.](#)
25. de Winde, C.M. *et al.* (2015) Multispectral imaging reveals the tissue distribution of tetraspanins in human lymphoid organs. [Histochem Cell Biol. 144 \(2\): 133-46.](#)
26. Dewals B.G., *et al.* (2011) Malignant catarrhal fever induced by Alcelaphine herpesvirus 1 is characterized by an expansion of activated CD3+CD8+CD4- T cells expressing a cytotoxic phenotype in both lymphoid and non-lymphoid tissues [Vet Res. 42:95](#)
27. Muljono, A. *et al.* (2009) Primary cutaneous lymphoblastic lymphoma in children: series of eight cases with review of the literature. [Pathology. 41 \(3\): 223-8.](#)
28. Sommer, A. *et al.* (2016) Infiltrating T lymphocytes reduce myeloid phagocytosis activity in synucleinopathy model. [J Neuroinflammation 13 \(1\): 174.](#)
29. Velu, V. *et al.* (2016) Induction of Th1-Biased T Follicular Helper (Tfh) Cells in Lymphoid Tissues during Chronic Simian Immunodeficiency Virus Infection Defines Functionally Distinct Germinal Center Tfh Cells. [J Immunol. 197 \(5\): 1832-42.](#)
30. Wen, J. *et al.* (2015) TNF-like weak inducer of apoptosis promotes blood brain barrier disruption and increases neuronal cell death in MRL/lpr mice. [J Autoimmun. 60: 40-50.](#)
31. Sample, S.J. *et al.* (2017) Radiographic and magnetic resonance imaging predicts severity of cruciate ligament fiber damage and synovitis in dogs with cranial cruciate ligament rupture. [PLoS One. 12 \(6\): e0178086.](#)
32. Zhang, M.Z. *et al.* (2015) Inhibition of cyclooxygenase-2 in hematopoietic cells results in salt-sensitive hypertension. [J Clin Invest. 125 \(11\): 4281-94.](#)
33. Kallikourdis, M. *et al.* (2017) T cell costimulation blockade blunts pressure overload-induced heart failure. [Nat Commun. 8: 14680.](#)
34. Bonnefont-Rebeix, C. *et al.* (2016) Characterization of a novel canine T-cell line established from a spontaneously occurring aggressive T-cell lymphoma with large granular cell morphology. [Immunobiology. 221 \(1\): 12-22.](#)
35. Montes-Cobos, E. *et al.* (2017) Targeted delivery of glucocorticoids to macrophages in a mouse model of multiple sclerosis using inorganic-organic hybrid nanoparticles. [J Control Release. 245: 157-169.](#)
36. Bartlett SL *et al.* (2010) Intestinal lymphoma of granular lymphocytes in a fisher (*Martes pennanti*) and a Eurasian otter (*Lutra lutra*). [J Zoo Wildl Med. 41 \(2\): 309-15.](#)
37. Houser, K.V. *et al.* (2017) Enhanced inflammation in New Zealand white rabbits when MERS-CoV reinfection occurs in the absence of neutralizing antibody. [PLoS Pathog. 13 \(8\): e1006565.](#)
38. Sparger, E.E. *et al.* (2018) Investigation of immune cell markers in feline oral squamous cell carcinoma. [Vet Immunol Immunopathol. 202: 52-62.](#)
39. Palomo, J. *et al.* (2018) The severity of imiquimod-induced mouse skin inflammation is independent of endogenous IL-38 expression. [PLoS One. 13 \(3\): e0194667.](#)

40. Declue, A.E. *et al.* (2018) Identification of immunologic and clinical characteristics that predict inflammatory response to C. Novyi-NT bacteriolytic immunotherapy. [BMC Vet Res. 14 \(1\): 119.](#)
41. DaSilva, A.V.A. *et al.* (2018) Morphophysiological changes in the splenic extracellular matrix of *Leishmania infantum*-naturally infected dogs is associated with alterations in lymphoid niches and the CD4+ T cell frequency in spleens. [PLoS Negl Trop Dis. 12 \(4\): e0006445.](#)
42. Withers, S.S. *et al.* (2018) Multi-color flow cytometry for evaluating age-related changes in memory lymphocyte subsets in dogs. [Dev Comp Immunol. 87: 64-74.](#)
43. Basu, A. *et al.* (2019) Association of PD-L1, PD-L2, and Immune Response Markers in Matched Renal Clear Cell Carcinoma Primary and Metastatic Tissue Specimens. [Am J Clin Pathol. 151 \(2\): 217-25.](#)
44. Pellegrini, S. *et al.* (2019) Selective local irradiation improves islet engraftment and survival in intra-bone marrow islet transplantation. [Cytotherapy. 21 \(10\): 1025-32.](#)
45. Ricat, C.M. *et al.* (2020) Immunohistochemical Findings in Idiopathic Inflammatory Bowel Disease in Nine Cats [BioMed Research International. 2020: 1-6.](#)

Further Reading

1. Alterio de Goss, M. *et al.* (1998) Control of cytomegalovirus in bone marrow transplantation chimeras lacking the prevailing antigen-presenting molecule in recipient tissues rests primarily on recipient-derived CD8 T cells. [J Virol. 72 \(10\): 7733-44.](#)
2. Burudi, E.M. *et al.* (2002) Regulation of indoleamine 2,3-dioxygenase expression in simian immunodeficiency virus-infected monkey brains. [J Virol. 76 \(23\): 12233-41.](#)
3. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. [Vet Res. 39: 54.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR73...)	RPE

Rabbit Anti Rat IgG (STAR21...) [HRP](#)
Rabbit Anti Rat IgG (STAR16...) [DyLight@800](#)
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) [DyLight@800](#)
Goat Anti Rat IgG (STAR72...) [HRP](#)

Recommended Negative Controls

[RAT IgG1 NEGATIVE CONTROL \(MCA6004GA\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M365354:200529'

Printed on 19 Oct 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)