

## Datasheet: MCA152

<b>Description:</b>	RAT ANTI MOUSE KAPPA LIGHT CHAIN
<b>Specificity:</b>	KAPPA LIGHT CHAIN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-20
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	1 mg

## Product Details

**RRID** AB\_321880

**Applications** This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**Target Species** Mouse

**Product Form** Purified IgG - liquid

**Buffer Solution** Phosphate buffered saline

**Preservative Stabilisers** 0.09% Sodium Azide

**Carrier Free** Yes

**Approx. Protein Concentrations** IgG concentration 1 mg/ml

**Immunogen** Purified mouse IgG.

**External Database** **UniProt:**

Links [P01837](#) [Related reagents](#)

**Entrez Gene:**

[16071](#) Igk-C [Related reagents](#)

---

**Fusion Partners** Spleen cells from an immunized PVG.RT1c rat were fused with cells of the rat Y3 Ag.1.2.3 myeloma cell line.

---

**Specificity** **Rat anti Mouse kappa light chain antibody, clone OX-20** recognizes the kappa light chain of murine immunoglobulins. OX-20 does not recognize mouse lambda light chains or bind to the immunoglobulin heavy chain.

---

**Flow Cytometry** Use 10ul of the suggested working dilution to label 10<sup>6</sup> cells in 100ul.

---

**References**

1. Mason, D.W. *et al.* (1981) The rat mixed lymphocyte reaction: roles of a dendritic cell in intestinal lymph and T-cell subsets defined by monoclonal antibodies. [Immunology. 44 \(1\): 75-87.](#)
2. White, R.A. *et al.* (1984) Differentiation markers in the Drosophila ovary. [J Embryol Exp Morphol. 84: 275-86.](#)
3. Meyer, K.B. *et al.* (1990) The importance of the 3'-enhancer region in immunoglobulin kappa gene expression. [Nucleic Acids Res. 18 \(19\): 5609-15.](#)
4. Thomas, M.J. *et al.* (1999) CD8 T-lymphocyte-mediated regulation of ovalbumin-specific murine IgE responses. [Int Arch Allergy Immunol. 118 \(2-4\): 289-91.](#)
5. Wasserman, G.M. *et al.* (1993) An approach to development of specific T-lymphocyte lines by use of preprocessed antigens in *Plasmodium vinckei vinckei* murine malaria. [Infect Immun. 61 \(5\): 1958-63.](#)
6. Thomas, M.J. *et al.* (2002) CD8 T cells inhibit IgE via dendritic cell IL-12 induction that promotes Th1 T cell counter-regulation. [J Immunol. 168 \(1\): 216-23.](#)
7. Salagianni, M. *et al.* (2007) An essential role for IL-18 in CD8 T cell-mediated suppression of IgE responses. [J Immunol. 178 \(8\): 4771-8.](#)

---

**Further Reading**

1. Galfrè, G. *et al.* (1979) Rat x rat hybrid myelomas and a monoclonal anti-Fd portion of mouse IgG. [Nature. 277 \(5692\): 131-3.](#)

---

**Storage** Store at +4°C for one month or at -20°C for longer.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 18 months from date of despatch.

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory** For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...) [FITC](#)

Goat Anti Rat IgG (STAR69...) [FITC](#)

Goat Anti Rat IgG (STAR131...)	<a href="#">Alk. Phos., Biotin</a>
Goat Anti Rat IgG (STAR73...)	<a href="#">RPE</a>
Rabbit Anti Rat IgG (STAR21...)	<a href="#">HRP</a>
Goat Anti Rat IgG (STAR72...)	<a href="#">HRP</a>
Rabbit Anti Rat IgG (STAR16...)	<a href="#">DyLight®800</a>
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	<a href="#">DyLight®800</a>

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M339476:181219'

**Printed on 20 May 2019**

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)