

Datasheet: MCA1522G

Description:	MOUSE ANTI POLY(ADP-RIBOSE) POLYMERASE-1
Specificity:	POLY(ADP-RIBOSE) POLYMERASE-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	A6.4.12
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			
Immunohistology - Paraffin (1)	▪			
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting	▪			1/1000 - 1/5000
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Clone A6.4.12 requires antigen retrieval using heat treatment prior to staining of paraffin sections. Sodium citrate buffer pH 6.0 is recommended for this purpose.

Target Species	Human
Species Cross Reactivity	Reacts with: Hamster, Mouse, Drosophila, Xenopus, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Human PARP-1
External Database Links	<p>UniProt: P09874 Related reagents</p> <p>Entrez Gene: 142 PARP1 Related reagents</p>
Synonyms	ADPRT, PPOL
RRID	AB_2236751
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of mouse NS0 myeloma cell line.
Specificity	<p>Mouse anti poly (ADP-ribose) polymerase 1 antibody, clone A6.4.12 recognizes poly (ADP-ribose) polymerase 1 (PARP-1), a ~116 kDa nuclear enzyme, cleaved during apoptosis (Soldani et al. 2002).</p> <p>PARP-1, a caretaker enzyme, is involved in DNA damage repair (Langelier et al. 2013), plays roles in diabetes pathophysiology (Andreone et al. 2012) and tumour proliferation (Rosado et al 2013).</p> <p>As well as protecting cells from genomic instability, PARP-1 is involved in the development of both inflammatory and immune responses, and cell death by apoptosis and necrosis (Erdélyi et al. 2005).</p> <p>Mouse anti poly(ADP-ribose) polymerase 1 antibody, clone A6.4.12, targets PARP-1, an enzyme which represents a promising target for new developments in therapeutic treatment of immune mediated diseases (Rosado et al. 2013). PARP-1 has considerable potential for delivering selective tumour cell killing while sparing normal cells (Pinton et al. 2013).</p>
References	1. Harris, J.L. <i>et al.</i> (2009) Aprataxin, poly-ADP ribose polymerase 1 (PARP-1) and apurinic endonuclease 1 (APE1) function together to protect the genome against oxidative

- damage. [Hum Mol Genet. 18: 4102-17.](#)
2. Freire, R. *et al.* (2001) Cleavage of the Bloom's syndrome gene product during apoptosis by caspase-3 results in an impaired interaction with topoisomerase IIIalpha. [Nucleic Acids Res. 29 \(15\): 3172-80.](#)
 3. Krohn, A.J. *et al.* (1998) Staurosporine-induced apoptosis of cultured rat hippocampal neurons involves caspase-1-like proteases as upstream initiators and increased production of superoxide as a main downstream effector. [J Neurosci. 18 \(20\): 8186-97.](#)
 4. Staples, C.J. *et al.* (2010) Cross-talk between the p38alpha and JNK MAPK pathways mediated by MAP kinase phosphatase-1 determines cellular sensitivity to UV radiation. [J Biol Chem. 285 \(34\): 25928-40.](#)
 5. Alexander, B.M. *et al.* (2010) DNA repair protein biomarkers associated with time to recurrence in triple-negative breast cancer. [Clin Cancer Res. 16: 5796-804.](#)
 6. Gueven, N. *et al.* (2004) Aprataxin, a novel protein that protects against genotoxic stress. [Hum Mol Genet. 13 \(10\): 1081-93.](#)
 7. Gueven, N. *et al.* (2006) Defective p53 response and apoptosis associated with an ataxia-telangiectasia-like phenotype. [Cancer Res. 66: 2907-12.](#)
 8. Kim, J.W. *et al.* (2000) Inhibition of homodimerization of poly(ADP-ribose) polymerase by its C-terminal cleavage products produced during apoptosis. [J Biol Chem. 275: 8121-5.](#)
 9. Hanzlikova, H. *et al.* (2017) Overlapping roles for PARP1 and PARP2 in the recruitment of endogenous XRCC1 and PNKP into oxidized chromatin. [Nucleic Acids Res. 45 \(5\): 2546-2557.](#)
 10. Olaussen, K.A. *et al.* (2013) PARP1 impact on DNA repair of platinum adducts: Preclinical and clinical read-outs. [Lung Cancer. 80: 216-22.](#)
 11. Fabrice, A. *et al.* (2012) PARP and adjuvant cisplatin-based chemotherapy in non-small-cell lung cancer. US patent: [20120277110](#)
 12. Geistrikh, I. *et al.* (2011) Ca²⁺-induced PARP-1 activation and ANF expression are coupled events in cardiomyocytes. [Biochem J. 438: 337-47.](#)
 13. Mirzaa, G.M. *et al.* (2014) Mutations in CENPE define a novel kinetochore-centromeric mechanism for microcephalic primordial dwarfism. [Hum Genet. 133: 1023-39.](#)
 14. Milner, R. *et al.* (2013) Validation of the BRCA1 antibody MS110 and the utility of BRCA1 as a patient selection biomarker in immunohistochemical analysis of breast and ovarian tumours. [Virchows Arch. 462: 269-79.](#)
 15. Inbar, D. *et al.* (2012) Erythropoietin-driven signalling and cell migration mediated by polyADP-ribosylation. [Br J Cancer. 107: 1317-26.](#)
 16. Buchsbaum, S. *et al.* (2012) FAT10 is a proteasomal degradation signal that is itself regulated by ubiquitination. [Mol Biol Cell. 23: 225-32.](#)
 17. Mullane, S.A. *et al.* (2016) Expression Levels of DNA Damage Repair Proteins Are Associated With Overall Survival in Platinum-Treated Advanced Urothelial Carcinoma. [Clin Genitourin Cancer. 14 \(4\): 352-9.](#)
 18. Zeng, J. *et al.* (2016) Nucleolar PARP-1 Expression Is Decreased in Alzheimer's Disease: Consequences for Epigenetic Regulation of rDNA and Cognition. [Neural Plast. 2016: 8987928.](#)
 19. Okuda, A. *et al.* (2017) Poly(ADP-ribose) polymerase inhibitors activate the p53 signaling pathway in neural stem/progenitor cells. [BMC Neurosci. 18 \(1\): 14.](#)
 20. Kubelac, P. *et al.* (2020) Changes in DNA Damage Response Markers with Treatment in Advanced Ovarian Cancer. [Cancers \(Basel\). 12\(3\): 707.](#)
-

Further Reading	<p>1. Pinton, G. <i>et al.</i> (2013) PARP1 inhibition affects pleural mesothelioma cell viability and uncouples AKT/mTOR axis via SIRT1. J Cell Mol Med. 17: 233-41.</p> <p>2. Rosado, M. <i>et al.</i> (2013) Beyond dna repair,the immunological role of parp-1 and its siblings. Immunology. 139: 428-37.</p> <p>3. Andreone, T. <i>et al.</i> (2012) Cytokine-mediatedβ-cell damage in PARP-1-deficient islets. Am J Physiol Endocrinol Metab. 303: E172-9.</p> <p>4. Langelier, M.F. and Pascal, J.M. (2013) PARP-1 mechanism for coupling DNA damage detection to poly(ADP-ribose) synthesis. Curr Opin Struct Biol. 23: 134-43.</p>
------------------------	--

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos. , HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight@800
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight@488 , DyLight@680 , DyLight@800 , FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com

'M365382:200529'

