

# Datasheet: MCA1531 BATCH NUMBER 167684

Description:	RAT ANTI HUMAN INTERLEUKIN-10
Specificity:	IL-10
Format:	Purified
<b>Product Type:</b>	Monoclonal Antibody
Clone:	JES3-9D7
Isotype:	IgG1
Quantity:	0.5 mg

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	•			1/10 - 1/50
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA	-			
Immunoprecipitation			•	
Western Blotting	•			1/100

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code <u>BUF09</u>) is recommended for this purpose.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein of supernatant	G from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	0.09% sodium azide (NaN <sub>3</sub> )	

Carrier Free	Yes					
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml					
Immunogen	Recombinant human IL-10.					
External Database						
Links	UniProt:					
	P22301 Related reagents					
	Entrez Gene:					
	3586 IL10 Related reagents					
RRID	AB_567068					
Specificity	Rat anti Human Interleukin-10 antibody, clone JES3-9D7 recognizes human Interleukin					
	- 10 (IL-10). IL-10 is an immunoregulatory cytokine which is expressed by activated					
	CD8+ve peripheral blood T-cells, CD4+ve T-cell clones, B cell lymphomas, activated					
	monocytes and mast cells.					
	Rat anti Human Interleukin-10 antibody, clone JES3-9D7 is reported to act as a					
	neutralising antibody ( <u>Othieno <i>et al.</i> 1999</u> ).					
Flow Cytometry	Use 10μl of the suggested working dilution to label 10 <sup>6</sup> cells in 100μl					
References	<ol> <li>Abrams, J.S. <i>et al.</i> (1992) Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. <u>Immunol Rev. 127: 5-24.</u></li> <li>Cua, D.J. <i>et al.</i> (1999) Transgenic interleukin 10 prevents induction of experimental</li> </ol>					
	autoimmune encephalomyelitis. <u>J Exp Med. 189: 1005-10.</u>					
	3. Lindmark, E. & Siegbahn, A. (2002) Tissue factor regulation and cytokine expression in monocyte-endothelial cell co-cultures: effects of a statin, an ACE-inhibitor and a					
	low-molecular-weight heparin. Thromb Res. 108 (1): 77-84.					
	4. Malmassari, S.L. et al. (2007) Impact of hepatitis B virus basic core promoter mutations					
	on T cell response to an immunodominant HBx-derived epitope. <u>Hepatology. 45 (5):</u> 1199-209.					
	5. Benjamin, J.L. <i>et al.</i> (2011) Randomised, double-blind, placebo-controlled trial of fructo-					
	oligosaccharides in active Crohn's disease. <u>Gut. 60 (7): 923-9.</u>					
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at					
	-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.					
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in					
	frost-free freezers is not recommended.					
Guarantee	12 months from date of despatch					

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:

https://www.bio-rad-antibodies.com/SDS/MCA1531

10040

Regulatory

For research purposes only

# **Related Products**

## **Recommended Secondary Antibodies**

Rabbit Anti Rat IgG (STAR16...) <u>DyLight®800</u>

Rabbit Anti Rat IgG (STAR17...)

Goat Anti Rat IgG (STAR72...)

Goat Anti Rat IgG (STAR69...)

Goat Anti Rat IgG (STAR73...)

RPE

Rabbit Anti Rat IgG (STAR21...)

Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) <u>DyLight®550</u>, <u>DyLight®650</u>, <u>DyLight®800</u>

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

### **Recommended Negative Controls**

#### RAT IgG1 NEGATIVE CONTROL (MCA6004GA)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M409947:221021'

#### Printed on 06 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint