

Datasheet: MCA1538 BATCH NUMBER 081214

Description:	MOUSE ANTI HUMAN THIOREDOXIN
Specificity:	THIOREDOXIN
Other names:	TXN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	2B1
Isotype:	lgG1
Quantity:	0.2 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal						
	communications from the	e originato	ors. Pleas	e refer to references ir	ndicated for further		
	information. For general	protocol i	recommer	ndations, please visit w	/ww.bio-		
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-					
	Immunohistology - Frozen	-					
	Immunohistology - Paraffin	-					
	ELISA	-					
	Immunoprecipitation						
	Western Blotting	-			10ug/ml - 50ug/ml		
	Where this antibody has not been tested for use in a particular technique this does not						
Target Species	system using appropriate	e negative	e/positive	controls.			
Target Species	Human						
Species Cross Reactivity	Reacts with: Marmoset N.B. Antibody reactivity a reactivity is derived from personal communications further information.	testing w	ithin our l	aboratories, peer-revie	wed publications or		
Product Form	Purified IgG - liquid						
Buffer Solution	Phosphate buffered salin	e					

Preservative Stabilisers	0.09% Sodium Azide			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
Immunogen	Recombinant human thioredoxin.			
External Database Links	UniProt: <u>P10599</u> <u>Related reagents</u> Entrez Gene: <u>7295</u> TXN <u>Related reagents</u>			
Synonyms	TRDX, TRX, TRX1			
RRID	AB_2212137			
Specificity	 Mouse anti Human thioredoxin antibody, clone 2B1 recognizes human thioredoxin, also known as ATL-derived factor or Surface-associated sulphydryl protein. Thioredoxin is a 105 amino acid ~12kDa dithiol oxidoreductase with powerful protein disulphide oxidoreductase activity containing a single thioredoxin domain (UniProt: P10599). Thioredoxin is involved in a number of cellular processes, including activation of the transcription factor NF-KB (Kelleher <i>et al.</i> 2014). Thioredoxin is highly expressed in the glandular cells of the GI tract and uterine tissues, also activated B and T-cells (Rubartelli <i>et al.</i> 1992). Mouse anti Human thioredoxin antibody, clone 2B1 has been used successfully in a biotinylated format as a detection reagent in a thioredoxin sandwich ELISA as part of a study looking at thioredoxin levels in placental trophoblast cells (Di Trapani <i>et al.</i> 1998), a study which also used the antibody for immunoprecipitation of thioredoxin as well as detection of thioredoxin antibody, clone 2B1 is also useful for the demonstration of thioredoxin expression on formalin fixed, paraffin embedded material by immunohistochemistry (Lincoln <i>et al.</i> 2003). Clone 2B1 has also been used as a capture reagent in a sandwich ELISA for thioredoxin of Plana and Plana antibody. 			
References	 Perkins, A.V. <i>et al.</i> (1995) Immunocytochemical localization trophoblast and decidua. <u>Placenta. 16 (7): 635-42.</u> Lincoln, D.T. <i>et al.</i> (2003) The thioredoxin-thioredoxin reduct over-expression in human cancer. <u>Anticancer Res. 23 (3B): 24</u> Stantchev, T.S. <i>et al.</i> (2012) Cell-type specific requirements during HIV-1 entry and infection. <u>Retrovirology. 9: 97.</u> Ma, X. <i>et al.</i> (2001) Regulation of interferon and retinoic aci activation through thioredoxin reductase. <u>J Biol Chem. 276: 24</u> Ramanathan, R.K. <i>et al.</i> (2007) A Phase I pharmacokinetic 	tase system: 1 <u>25-33.</u> for thiol/disulfide exchange d-induced cell death 1843-54.		

	study of PX-12, a novel inhibitor of thioredoxin-1, in patients with advanced solid tumors.			
	Clin Cancer Res. 13: 2109-14.			
	6. Arnold, N.B. <i>et al.</i> (2004) Thioredoxin is downstream of Smad7 in a pathway that			
	promotes growth and suppresses cisplatin-induced apoptosis in pancreatic cancer. <u>Cancer</u>			
	Res. 64: 3599-606.			
	7. Kinoshita, T. <i>et al.</i> (2007) Thioredoxin prevents the development and progression of			
	elastase-induced emphysema. <u>Biochem Biophys Res Commun. 354: 712-9.</u>			
	8. Dorion, S. <i>et al.</i> (2002) Activation of the p38 signaling pathway by heat shock involves			
	the dissociation of glutathione S-transferase Mu from Ask1. <u>J Biol Chem. 277: 30792-7.</u>			
	9. Sensi, M. <i>et al.</i> (2005) Immunogenicity without immunoselection: a mutant but			
functional antioxidant enzyme retained in a human metastatic melanoma and t				
	CD8(+) T cells with a memory phenotype. <u>Cancer Res. 65: 632-40.</u>			
	10. Mamoon, N.M. <i>et al.</i> (2007) Multiple cysteine residues are implicated in Janus kinase			
	2-mediated catalysis. Biochemistry. 46: 14810-8.			
	11. Agathanggelou, A. et al. (2003) Identification of Novel Gene Expression Targets for the			
	Ras Association Domain Family 1 (RASSF1A) Tumor Suppressor Gene in Non-Small Cell			
	Lung Cancer and Neuroblastoma. <u>Cancer Res. 63: 5344-51.</u>			
	12. Di Trapani, G. et al. (1998) Production and secretion of thioredoxin from transformed			
	human trophoblast cells. <u>Mol Hum Reprod. 4: 369-75.</u>			
	13. Lopata, A. et al. (2001) Expression and localization of thioredoxin during early			
	implantation in the marmoset monkey. <u>Mol Hum Reprod. 7: 1159-65.</u>			
	14. Wang, M.Y. et al. (2011) A redox switch in C-reactive protein modulates activation of			
	endothelial cells. <u>FASEB J. 25 (9): 3186-96.</u>			
Storage	Store at +4°C or at -20°C if preferred.			
	This product should be stored undiluted.			
	Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing			
	as this may denature the antibody. Should this product contain a precipitate we			
	recommend microcentrifugation before use.			
Guarantee	12 months from date of despatch			
Hoolth And Cofety				
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1538			
internation	10040			
Regulatory	For research purposes only			

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12)	<u>RPE</u>
Goat Anti Mouse IgG IgA IgM (STAR87) <u>HRP</u>
Goat Anti Mouse IgG (STAR76)	<u>RPE</u>
Goat Anti Mouse IgG (STAR70)	<u>FITC</u>

Goat Anti Mouse IgG (H/L) (STAR117) <u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®550</u> ,							
		DyLight®650, DyLight®680, DyLight®800,					
		FIT	<u>FITC, HRP</u>				
Goat Anti Mo	ouse IgG (STAR77)	HRP					
Goat Anti Mo	ouse IgG (Fc) (STAR120) <u>FI</u>	FITC, HRP				
Rabbit Anti M	Rabbit Anti Mouse IgG (STAR13) <u>HRP</u>						
Rabbit Anti M	Rabbit Anti Mouse IgG (STAR9) <u>FITC</u>						
Recommended Negative Controls							
MOUSE IgG1 NEGATIVE CONTROL (MCA928)							
North & South Tel:	+1 800 265 7376 World	wide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21		
America Fax	x: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50		
Em	ail: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com		
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365390:200529'							
Printed on 01 May 2024							

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint