

Datasheet: MCA1539F

BATCH NUMBER 151139

Description:	MOUSE ANTI HUMAN CD95:FITC
Specificity:	CD95
Other names:	FAS
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	LOB 3/17
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Species Cross Reactivity

Reacts with: Rhesus Monkey

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on Protein A

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
Immunogen	Fusion protein comprising extracellular domain of human Fas linked to human Fc.
External Database Links	<p>UniProt: P25445 Related reagents</p> <p>Entrez Gene: 355 FAS Related reagents</p>
Synonyms	APT1, FAS1, TNFRSF6
RRID	AB_321976
Fusion Partners	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NSI myeloma cell line.
Specificity	Mouse anti Human CD95 antibody, clone LOB 3/17 recognizes the human CD95 cell surface antigen, also known as Tumor necrosis factor receptor superfamily member 6, Fas, Apo-1 antigen, Apoptosis-mediating surface antigen FAS or FASLG receptor. CD95 is a 310 amino acid ~40-50 kDa single pass type I transmembrane glycoprotein expressed by activated T and B cells, NK cells and thymocytes. Mutations in the CD95 gene, FAS can lead to the development of Autoimmune lymphoproliferative syndrome 1A (ALPS1A), an apoptotic disorder with early onset resulting in an accumulation of autoreactive lymphocytes (Peters et al. 1999).
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Mesdaghi, M. <i>et al.</i> (2010) Natural killer cells in allergic rhinitis patients and nonatopic controls. Int Arch Allergy Immunol. 153 (3): 234-8. Ximeri, M. <i>et al.</i> (2010) Effect of lenalidomide therapy on hematopoiesis of patients with myelodysplastic syndrome associated with chromosome 5q deletion. Haematologica. 95 (3): 406-14. Aref, S. <i>et al.</i> (2004) Accelerated neutrophil apoptosis in neutropenic patients with hepatosplenic schistosomiasis is induced by serum Fas ligand. Hematol J. 5 (5): 434-9. Welsh, J.P. <i>et al.</i> (2004) In vitro effects of interferon-gamma and tumor necrosis factor-alpha on CD34+ bone marrow progenitor cells from aplastic anemia patients and normal donors. Hematol J. 5 (1): 39-46. Wethkamp, N. <i>et al.</i> (2011) Daxx-beta and Daxx-gamma, two novel splice variants of the transcriptional co-repressor Daxx. J Biol Chem. 286 (22): 19576-88. Chen, J.Y. <i>et al.</i> (2003) TNF-alpha renders human peritoneal mesothelial cells sensitive to anti-Fas antibody-induced apoptosis. Nephrol Dial Transplant. 18 (9): 1741-7. Papadaki, H.A. <i>et al.</i> (2002) Bone marrow progenitor cell reserve and function and

stromal cell function are defective in rheumatoid arthritis: evidence for a tumor necrosis factor alpha-mediated effect. [Blood. 99 \(5\): 1610-9.](#)

8. Mavroudi, I. *et al.* (2011) The CD40/CD40 ligand interactions exert pleiotropic effects on bone marrow granulopoiesis. [J Leukoc Biol. 89 \(5\): 771-83.](#)

9. Pyrovolaki, K. *et al.* (2009) Increased expression of CD40 on bone marrow CD34+ hematopoietic progenitor cells in patients with systemic lupus erythematosus: contribution to Fas-mediated apoptosis. [Arthritis Rheum. 60 \(2\): 543-52.](#)

10. Boula, A. *et al.* (2006) Effect of cA2 anti-tumor necrosis factor-alpha antibody therapy on hematopoiesis of patients with myelodysplastic syndromes. [Clin Cancer Res. 12 \(10\): 3099-108.](#)

11. Papadaki, H.A. *et al.* (2005) Normal bone marrow hematopoietic stem cell reserves and normal stromal cell function support the use of autologous stem cell transplantation in patients with multiple sclerosis. [Bone Marrow Transplant. 36 \(12\): 1053-63.](#)

12. Bachsais, M. *et al.* (2016) The Interaction of CD154 with the $\alpha 5\beta 1$ Integrin Inhibits Fas-Induced T Cell Death. [PLoS One. 11 \(7\): e0158987.](#)

13. Ismail, M. *et al.* (2001) Bcl-2 and Bcl-x expression in the CD34+ cells of aplastic anaemia patients: relationship with increased apoptosis and upregulation of Fas antigen. [Br J Haematol. 113 \(3\): 706-12.](#)

14. Bachsais, M. *et al.* (2020) CD154 inhibits death of T cells via a Cis interaction with the $\alpha 5\beta 1$ integrin. [PLoS One. 15 \(8\): e0235753.](#)

15. Ismail, M.M. *et al.* (2003) Differential apoptosis and Fas expression on GPI-negative and GPI-positive stem cells: a mechanism for the evolution of paroxysmal nocturnal haemoglobinuria. [Br J Haematol. 123 \(3\): 545-51.](#)

Further Reading 1. Paulsen, M. & Janssen, O. (2011) Pro- and anti-apoptotic CD95 signaling in T cells. [Cell Commun Signal. 9: 7.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1539F>
10041

Regulatory For research purposes only

Related Products

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL:FITC \(MCA928F\)](#)

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M365392:200529'

Printed on 19 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)