

Datasheet: MCA153XZ

Description:	MOUSE ANTI RAT CD4 (DOMAIN 2):Preservative Free
Specificity:	CD4 (DOMAIN 2)
Format:	Preservative Free
Product Type:	Monoclonal Antibody
Clone:	OX-35
Isotype:	IgG2a
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Western Blotting			▪	
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	None present
Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	MLR generated rat T helper lymphocytes.
External Database	UniProt:

Links

[P05540](#) [Related reagents](#)

Entrez Gene:

[24932](#) Cd4 [Related reagents](#)

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.

Specificity

Mouse anti Rat CD4 (Domain 2) antibody, clone OX-35 recognizes the rat CD4 cell surface antigen, a ~55kDa glycoprotein expressed by helper T cells and weakly by monocytes.

Mouse anti Rat CD4 (Domain 2) antibody, clone OX-35 recognizes a different epitope on the CD4 molecule to Mouse anti Rat CD4 antibody, [clone W3/25](#).

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Wang, C.C. *et al.* (1996) Immunohistochemical study of amoeboid microglial cells in fetal rat brain. [J Anat. 189 \(Pt 3\): 567-74.](#)
2. Jefferies, W.A. *et al.* (1985) Authentic T helper CD4 (W3/25) antigen on rat peritoneal macrophages. [J Exp Med. 162 \(1\): 117-27.](#)
3. Camelo, S. *et al.* (2004) The distribution of antigen in lymphoid tissues following its injection into the anterior chamber of the rat eye. [J Immunol. 172: 5388-95.](#)
4. Elflein, K. *et al.* (2003) Rapid recovery from T lymphopenia by CD28 superagonist therapy. [Blood. 102: 1764-70.](#)
5. Scherr, M. *et al.* (2002) Efficient gene transfer into the CNS by lentiviral vectors purified by anion exchange chromatography. [Gene Ther. 9: 1708-14.](#)
6. Cho, K.S. *et al.* (2010) Mechanism analysis of long-term graft survival by monocarboxylate transporter-1 inhibition. [Transplantation. 90: 1299-306.](#)
7. Chang, C.J. *et al.* (2004) The immunization site of cytokine-secreting tumor cell vaccines influences the trafficking of tumor-specific T lymphocytes and antitumor efficacy against regional tumors. [J Immunol. 173: 6025-32.](#)
8. Basiri, M. and Doucette, R. (2010) Sensorimotor cortex aspiration: a model for studying Wallerian degeneration-induced glial reactivity along the entire length of a single CNS axonal pathway. [Brain Res Bull. 81: 43-52.](#)
9. Esquifino, A.I. *et al.* (2007) Immune response after experimental allergic encephalomyelitis in rats subjected to calorie restriction. [J Neuroinflammation. 4:6.](#)
10. Zhao, S. *et al.* (2007) Extensive FDG uptake and its modification with corticosteroid in a granuloma rat model: an experimental study for differentiating granuloma from tumors. [Eur J Nucl Med Mol Imaging 34: 2096-105.](#)
11. Yan, Y. *et al.* (2003) Pathogenesis of autoimmunity after xenogeneic thymus transplantation. [J Immunol. 170: 5936-46.](#)

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life

18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10162 available at:
10162: <https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR8...) [DyLight®800](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Human Anti Mouse IgG2a (HCA037...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight®488](#), [DyLight®549](#),
[DyLight®649](#), [DyLight®680](#), [DyLight®800](#),
[FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL:Preservative Free \(MCA1210XZ\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

'M319039:180719'

Printed on 28 Jul 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)