

Datasheet: MCA1548

BATCH NUMBER 164585

Description:	RAT ANTI MOUSE INTERFERON GAMMA
Specificity:	IFN GAMMA
Other names:	INTERFERON GAMMA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	XMG1.2
Isotype:	IgG1
Quantity:	0.25 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/10 - 1/50
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1ug/ml - 10ug/ml
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilization is required for this application. The use of Leucoperm (Product Code [BUF09](#)) is recommended for this purpose.

Target Species	Mouse
Product Form	Purified IgG - liquid
Preparation	Antibody purified from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative	0.09% sodium azide (NaN ₃)

Stabilisers

Carrier Free Yes

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen Recombinant mouse IFN gamma.

External Database Links

UniProt:
[P01580](#) [Related reagents](#)

Entrez Gene:
[15978](#) Ifng [Related reagents](#)

RRID AB_324085

Specificity **Rat anti Mouse Interferon gamma antibody, clone XMG1.2** recognizes murine IFN gamma. Rat anti Mouse Interferon gamma antibody, clone XMG1.2 may be used for the detection of intracellular IFN gamma using flow cytometric techniques and has been shown to block the functional activity of murine interferon gamma.

Flow Cytometry Use 10µl of the suggested working dilution to label 1x10⁶ cells in 100µl

References

1. Cherwinski, H.M. *et al.* (1987) Two types of mouse helper T cell clone. III. Further differences in lymphokine synthesis between Th1 and Th2 clones revealed by RNA hybridization, functionally monospecific bioassays, and monoclonal antibodies. [J Exp Med. 166 \(5\): 1229-44.](#)
2. Holz, A. *et al.* (2001) Constitutive beta cell expression of IL-12 does not perturb self-tolerance but intensifies established autoimmune diabetes. [J Clin Invest. 108 \(12\): 1749-58.](#)
3. Hsieh, B. *et al.* (1996) *In vivo* cytokine production in murine listeriosis. Evidence for immunoregulation by gamma delta+ T cells. [J Immunol. 156 \(1\): 232-7.](#)
4. Chiu, W.C. *et al.* (2009) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. [J Nutr Biochem. 20: 254-60.](#)
5. Hsu, C.S. *et al.* (2006) Dietary fish oil enhances adhesion molecule and interleukin-6 expression in mice with polymicrobial sepsis. [Br J Nutr. 96: 854-60.](#)
6. Rountree, R.B. *et al.* (2011) Exosome Targeting of Tumor Antigens Expressed by Cancer Vaccines Can Improve Antigen Immunogenicity and Therapeutic Efficacy. [Cancer Res. 71: 5235-44.](#)
7. Pletinckx K *et al.* (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3- IL-10+ regulatory T cells by engaging CD28 and CTLA-4. [Eur J Immunol. 45 \(2\): 480-91.](#)
8. Tsagozis, P. *et al.* (2003) CD8(+) T cells with parasite-specific cytotoxic activity and a Tc1 profile of cytokine and chemokine secretion develop in experimental visceral leishmaniasis. [Parasite Immunol. 25 \(11-12\): 569-79.](#)
9. ReisL, E.S. *et al.* (2017) Mixed Formulation of Conventional and Pegylated Meglumine

Antimoniate-Containing Liposomes Reduces Inflammatory Process and Parasite Burden in *Leishmania infantum*-Infected BALB/c Mice. [Antimicrob Agents Chemother. 61 \(11\)Oct 24 \[Epub ahead of print\]](#).

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1548>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR72...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight@550 , DyLight@650 , DyLight@800
Rabbit Anti Rat IgG (STAR21...)	HRP
Rabbit Anti Rat IgG (STAR16...)	DyLight@800

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M410022:221024'

Printed on 25 Mar 2023