

Datasheet: MCA1548EL

Description:	RAT ANTI MOUSE INTERFERON GAMMA:Low Endotoxin
Specificity:	IFN GAMMA
Other names:	INTERFERON GAMMA
Format:	Low Endotoxin
Product Type:	Monoclonal Antibody
Clone:	XMG1.2
Isotype:	lgG1
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	•			
Immunohistology - Frozen				1/200 - 1/2000
Immunohistology - Paraffin				
ELISA	•			
Functional Assays	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm[™] (Product Code <u>BUF09</u>) for this purpose.

Target Species	Mouse	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein C supernatant	6 from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	None present	

Carrier Free	Yes			
Endotoxin Level	<0.01EU/ug			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
Immunogen	Recombinant mouse IFN gamma.			
External Database Links	UniProt: P01580 Related reagents Entrez Gene: 15978 Ifng Related reagents			
RRID	AB_2280100			
Specificity	Rat anti Mouse Interferon gamma antibody, clone XMG1.2 recognizes murine IFN gamma. Rat anti Mouse Interferon gamma antibody, clone XMG1.2 may be used for the detection of intracellular IFN gamma using flow cytometric techniques and has been shown to block the functional activity of murine interferon gamma.			
Flow Cytometry	Use 10ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.			

References

1. Cherwinski, H.M. et al. (1987) Two types of mouse helper T cell clone. III. Further differences in lymphokine synthesis between Th1 and Th2 clones revealed by RNA hybridization, functionally monospecific bioassays, and monoclonal antibodies. J Exp Med. <u>166 (5): 1229-44.</u>

be used for the

- 2. Holz, A. et al. (2001) Constitutive beta cell expression of IL-12 does not perturb self-tolerance but intensifies established autoimmune diabetes. J Clin Invest. 108 (12): <u>1749-58.</u>
- 3. Hsieh, B. et al. (1996) In vivo cytokine production in murine listeriosis. Evidence for immunoregulation by gamma delta+ T cells. J Immunol. 156 (1): 232-7.
- 4. Chiu, W.C. et al. (2009) Effects of dietary fish oil supplementation on cellular adhesion molecule expression and tissue myeloperoxidase activity in hypercholesterolemic mice with sepsis. J Nutr Biochem. 20: 254-60.
- 5. Hsu, C.S. et al. (2006) Dietary fish oil enhances adhesion molecule and interleukin-6 expression in mice with polymicrobial sepsis. Br J Nutr. 96: 854-60.
- 6. Rountree, R.B. et al. (2011) Exosome Targeting of Tumor Antigens Expressed by Cancer Vaccines Can Improve Antigen Immunogenicity and Therapeutic Efficacy. Cancer Res. 71: 5235-44.
- 7. Pletinckx K et al. (2015) Immature dendritic cells convert anergic nonregulatory T cells into Foxp3- IL-10+ regulatory T cells by engaging CD28 and CTLA-4. Eur J Immunol. 45 (2): 480-91.
- 8. Tsagozis, P. et al. (2003) CD8(+) T cells with parasite-specific cytotoxic activity and a Tc1 profile of cytokine and chemokine secretion develop in experimental visceral leishmaniasis. Parasite Immunol. 25 (11-12): 569-79.

9. ReisL, E.S. et al. (2017) Mixed Formulation of Conventional and Pegylated Meglumine Antimoniate-Containing Liposomes Reduces Inflammatory Process and Parasite Burden in Leishmania infantum-Infected BALB/c Mice. Antimicrob Agents Chemother. 61 (11)Oct 24 [Epub ahead of print].

Storage

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

18 months from date of despatch.

Health And Safety Information

Material Safety Datasheet documentation #10162 available at:

10162: https://www.bio-rad-antibodies.com/uploads/MSDS/10162.pdf

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR17...) **FITC** Goat Anti Rat IgG (STAR69...) **FITC**

Goat Anti Rat IgG (STAR131...) Alk. Phos., Biotin

Goat Anti Rat IgG (STAR73...) **RPE** Rabbit Anti Rat IgG (STAR21...) **HRP**

Rabbit Anti Rat IgG (STAR16...) DyLight®800 Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...) DyLight®800

Goat Anti Rat IgG (STAR72...) **HRP**

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M331488:180912'

Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint