

Datasheet: MCA154GA

| | |
|----------------------|---------------------------|
| Description: | MOUSE ANTI RAT CD2 |
| Specificity: | CD2 |
| Other names: | E-ROSETTE RECEPTOR, LFA-2 |
| Format: | Purified |
| Product Type: | Monoclonal Antibody |
| Clone: | OX-34 |
| Isotype: | IgG2a |
| Quantity: | 0.1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/50 |
| Immunohistology - Frozen | ▪ | | | |
| Immunohistology - Paraffin (1) | ▪ | | | |
| ELISA | | | ▪ | |
| Immunoprecipitation | ▪ | | | |
| Western Blotting | | | ▪ | |
| Immunofluorescence | ▪ | | | |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1) **We recommend the use of [PLP](#) fixation for optimal results, see Whiteland et al.**

| | |
|------------------------|---|
| Target Species | Rat |
| Product Form | Purified IgG - liquid |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant |
| Buffer Solution | Phosphate buffered saline |

| | |
|---------------------------------------|---|
| Preservative Stabilisers | 0.09% sodium azide (NaN ₃) |
| Carrier Free | Yes |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml |
| Immunogen | Activated rat T helper cells. |
| External Database Links | <p>UniProt: P08921 Related reagents</p> <p>Entrez Gene: 497761 Cd2 Related reagents</p> |
| RRID | AB_566608 |
| Fusion Partners | Spleen cells from immunised BALB/c mice were fused with cells of the NS1 mouse myeloma cell line. |
| Specificity | Mouse anti Rat CD2 antibody, clone OX-34 recognizes a determinant on thymocytes and peripheral T-cells but it does not bind to B cells or peritoneal macrophages. The antigen recognized by this antibody is a 50-54 kDa glycoprotein, homolog of the human CD2 antigen (Williams et al. 1987). |
| Flow Cytometry | Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl |
| References | <ol style="list-style-type: none"> Williams, A.F. <i>et al.</i> (1987) Similarities in sequences and cellular expression between rat CD2 and CD4 antigens. J Exp Med. 165 (2): 368-80. Barclay, A.N. (1981) The localization of populations of lymphocytes defined by monoclonal antibodies in rat lymphoid tissues. Immunology. 42 (4): 593-600. Whiteland, J.L. <i>et al.</i> (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20. Baker, S.C. <i>et al.</i> (2011) Cellular Integration and Vascularisation Promoted by a Resorbable, Particulate-Leached, Cross-Linked Poly(ε-caprolactone) Scaffold. Macromol Biosci. 11: 618-27. Romani, P. <i>et al.</i> (2009) Cell survival and polarity of <i>Drosophila</i> follicle cells require the activity of ecdysone receptor B1 isoform. Genetics. 181: 165-75. Bastock, R. <i>et al.</i> (2003) Strabismus is asymmetrically localised and binds to Prickle and Dishevelled during <i>Drosophila</i> planar polarity patterning. Development. 130: 3007-14. Brückner, K. <i>et al.</i> (2000) Glycosyltransferase activity of Fringe modulates Notch-Delta interactions. Nature. 406: 411-5. Liversidge, J. <i>et al.</i> (2002) Nitric oxide mediates apoptosis through formation of peroxynitrite and Fas/Fas-ligand interactions in experimental autoimmune uveitis. Am J Pathol. 160: 905-16. Sarpal, R. <i>et al.</i> (2012) Mutational analysis supports a core role for <i>Drosophila</i> |

α -catenin in adherens junction function. [J Cell Sci. 125: 233-45.](#)

10. Zhang, H. *et al.* (2011) Basic residues in the T-cell receptor ζ cytoplasmic domain mediate membrane association and modulate signaling. [Proc Natl Acad Sci U S A. 108: 19323-8.](#)

11. Heck, B.W. *et al.* (2012) The transcriptional corepressor SMRTER influences both Notch and ecdysone signaling during *Drosophila* development. [Biol Open. 1 \(3\): 182-96.](#)

12. Clark, I.B. *et al.* (2011) Fibroblast growth factor signalling controls successive cell behaviours during mesoderm layer formation in *Drosophila*. [Development. 138: 2705-15.](#)

13. Domanitskaya, E. and Schüpbach, T. (2012) CoREST acts as a positive regulator of Notch signaling in the follicle cells of *Drosophila melanogaster*. [J Cell Sci. 125: 399-410.](#)

14. Dragovic, R.A. *et al.* (2015) Isolation of syncytiotrophoblast microvesicles and exosomes and their characterisation by multicolour flow cytometry and fluorescence Nanoparticle Tracking Analysis. [Methods. 87: 64-74.](#)

15. Zecca, M. & Struhl, G. (2021) A unified mechanism for the control of *Drosophila* wing growth by the morphogens Decapentaplegic and Wingless. [PLoS Biol. 19 \(3\): e3001111.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA154GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@550](#),
[DyLight@650](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)

Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA1210\)](#)

| | | | | | |
|----------------------------------|---|------------------|---|---------------|---|
| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com | Europe | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com |
|----------------------------------|---|------------------|---|---------------|---|

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M419920:230704'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)