

Datasheet: MCA1556APC

Description:	MOUSE ANTI HUMAN CD10:APC		
Specificity:	CD10		
Other names:	CALLA		
Format:	APC		
Product Type:	Monoclonal Antibody		
Clone:	SN5c		
Isotype:	IgG1		
Quantity:	100 TESTS		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human					
Product Form	Purified IgG conju	Purified IgG conjugated to allophycocyanin (APC) - lyophilized				
Reconstitution	Reconstitute with	1 ml distilled water				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)			
	APC	650	661			
Preparation	Purified IgG preparts	ared by affinity chromatog	raphy on Protein A from tissue cu	ultuı		
Buffer Solution	Phosphate buffer	ed saline				
Preservative	0.09% sodium az	ide (NaN ₃)				
Stabilisers	1% bovine serum	albumin				
	5% sucrose					

lm	mı	un	OC	ien	ì

Partially purified cell membrane antigens from fresh leukemia cells

External Database Links

UniProt:

P08473 Related reagents

Entrez Gene:

4311 MME Related reagents

Synonyms

EPN

RRID

AB_322806

Fusion Partners

Spleen cells from immunized BALB/c mice were fused with cells of the mouse PS/NS1 /1-Ag4-1 myeloma cell line

Specificity

Mouse anti Human CD10 antibody, clone SN5c recognizes human neprilysin, also known as CD10, atriopeptidase, enkephalinase, neutral endopeptidase 24.11, skin fibroblast elastase or common acute lymphocytic leukemia antigen (CALLA). CD10 is a 749 aminoacid, ~100 kDa single pass type II transmembrane glycoprotein expressed by acute lymphoblastic leukaemia cells and by peripheral blood granulocytes.

Defects in the MME gene encoding CD10 can lead to the development of the peripheral nervous system disorder, Charcot-Marie-Tooth disease 2T (CMY2T), an axonal form of Marie-Charcot-Tooth disease characterized by either dominantly inherited primary peripheral demyelinating neuropathies, designated CMT1, or primary peripheral axonal neuropathies showing axonal degeneration in the absence of any obvious myelin alteration (CMT2).

Flow Cytometry

Use 10µl of the suggested working dilution to label 10⁶ cells in 100µl

References

- 1. Biddle, W.C. *et al.* (1989) *In vitro* and *in vivo* cytotoxic activity of anti-human leukemia monoclonal antibodies SN5c and SN6 daunorubicin conjugates. <u>Leuk Res. 13 (8):</u> 699-707.
- 2. Riemann, D. *et al.* (2001) Caveolae/lipid rafts in fibroblast-like synoviocytes: ectopeptidase-rich membrane microdomains. Biochem J. 354 (Pt 1): 47-55.
- 3. Diaz-Romero, J. *et al.* (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect *bona fide* dedifferentiation rather than amplification of progenitor cells. J Cell Physiol. 214: 75-83.
- 4. Pilling, D. *et al.* (2009) Identification of markers that distinguish monocyte-derived fibrocytes from monocytes, macrophages, and fibroblasts. PLoS One. 4 (10): e7475.
- 5. Manini, I. *et al.* (2020) Heterogeneity Matters: Different Regions of Glioblastoma Are Characterized by Distinctive Tumor-Supporting Pathways. <u>Cancers (Basel). 12 (10)Oct 13 [Epub ahead of print].</u>
- 6. Glynn, E. & Fromm, J.R. (2020) Immunophenotypic Characterization and Purification of Neoplastic Cells from Lymph Nodes Involved by T-Cell/Histiocyte-rich Large B-cell Lymphoma by Flow Cytometry and Flow Cytometric Cell Sorting. Cytometry B Clin Cytom. 98 (1): 88-98.

7. Caponnetto, F. et al. (2020) Human Adipose-Derived Stem Cells in Madelung's Disease: Morphological and Functional Characterization. Cells. 10 (1): 44.

8. Kirolos, S.A. and Gomer, R.H. (2022) The extracellular sialidase NEU3 induces neutrophil priming bioRxiv. 24 Feb [Epub ahead of print].

Storage

Prior to reconstitution store at +4°C. Following reconstitution store at +4°C.

DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend

microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #20487 available at:

https://www.bio-rad-antibodies.com/SDS/MCA1556APC

20487

Regulatory

For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:APC (MCA928APC)

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

America

Fax: +1 919 878 3751

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419716:230616'

Printed on 03 Oct 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint