

Datasheet: MCA1556GA

BATCH NUMBER 165252

Description:	MOUSE ANTI HUMAN CD10
Specificity:	CD10
Other names:	CALLA
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	SN5c
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% sodium azide (NaN ₃)

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Partially purified cell membrane antigens from fresh leukemia cells
External Database Links	<p>UniProt: P08473 Related reagents</p> <p>Entrez Gene: 4311 MME Related reagents</p>
Synonyms	EPN
RRID	AB_323728
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse PS/NS1 /1-Ag4-1 myeloma cell line
Specificity	<p>Mouse anti Human CD10 antibody, clone SN5c recognizes human neprilysin, also known as CD10, atriopetidase, enkephalinase, neutral endopeptidase 24.11, skin fibroblast elastase or common acute lymphocytic leukemia antigen (CALLA). CD10 is a 749 aminoacid, ~100 kDa single pass type II transmembrane glycoprotein expressed by acute lymphoblastic leukaemia cells and by peripheral blood granulocytes.</p> <p>Defects in the MME gene encoding CD10 can lead to the development of the peripheral nervous system disorder, Charcot-Marie-Tooth disease 2T (CMY2T), an axonal form of Marie-Charcot-Tooth disease characterized by either dominantly inherited primary peripheral demyelinating neuropathies, designated CMT1, or primary peripheral axonal neuropathies showing axonal degeneration in the absence of any obvious myelin alteration (CMT2).</p>
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 ⁶ cells in 100µl
References	<ol style="list-style-type: none"> Biddle, W.C. <i>et al.</i> (1989) <i>In vitro</i> and <i>in vivo</i> cytotoxic activity of anti-human leukemia monoclonal antibodies SN5c and SN6 daunorubicin conjugates. Leuk Res. 13 (8): 699-707. Riemann, D. <i>et al.</i> (2001) Caveolae/lipid rafts in fibroblast-like synoviocytes: ectopeptidase-rich membrane microdomains. Biochem J. 354 (Pt 1): 47-55. Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect <i>bona fide</i> dedifferentiation rather than amplification of progenitor cells. J Cell Physiol. 214: 75-83. Pilling, D. <i>et al.</i> (2009) Identification of markers that distinguish monocyte-derived fibrocytes from monocytes, macrophages, and fibroblasts. PLoS One. 4 (10): e7475. Manini, I. <i>et al.</i> (2020) Heterogeneity Matters: Different Regions of Glioblastoma Are Characterized by Distinctive Tumor-Supporting Pathways. Cancers (Basel). 12 (10)Oct 13

[\[Epub ahead of print\]](#).

6. Glynn, E. & Fromm, J.R. (2020) Immunophenotypic Characterization and Purification of Neoplastic Cells from Lymph Nodes Involved by T-Cell/Histiocyte-rich Large B-cell Lymphoma by Flow Cytometry and Flow Cytometric Cell Sorting. [Cytometry B Clin Cytom. 98 \(1\): 88-98.](#)

7. Caponnetto, F. *et al.* (2020) Human Adipose-Derived Stem Cells in Madelung's Disease: Morphological and Functional Characterization. [Cells. 10 \(1\): 44.](#)

8. Kirolos, S.A. and Gomer, R.H. (2022) The extracellular sialidase NEU3 induces neutrophil priming [bioRxiv. 24 Feb \[Epub ahead of print\]](#).

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/MCA1556GA>
10040

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...) [RPE](#)
Goat Anti Mouse IgG IgA IgM (STAR87...) [HRP](#)
Goat Anti Mouse IgG (STAR76...) [RPE](#)
Goat Anti Mouse IgG (STAR70...) [FITC](#)
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@550](#),
[DyLight@650](#), [DyLight@680](#), [DyLight@800](#),
[FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)
Goat Anti Mouse IgG (STAR77...) [HRP](#)
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M410062:221024'

