

Datasheet: MCA1556SBB580

Description:	MOUSE ANTI HUMAN CD10:StarBright Blue 580
Specificity:	CD10
Other names:	CALLA
Format:	StarBright Blue 580
Product Type:	Monoclonal Antibody
Clone:	SN5c
Isotype:	lgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugate	ed to StarBright Blue 5	580 - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	StarBright Blue 580	475	582
Preparation	Purified IgG prepared supernatant	by affinity chromatog	raphy on Protein A
Buffer Solution	Phosphate buffered sa	aline	
Preservative	0.09% sodium azide (NaN ₃)	
Stabilisers	1% bovine serum albu	ımin	
	0.1% Pluronic F68		
	0.1% Pluronic F68 0.1% PEG 3350		

Immunogen	Partially purified cell membrane antigens from fresh leukemia cells			
External Database Links	UniProt: P08473 Related reagents Entrez Gene: 4311 MME Related reagents			
Synonyms	EPN			
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse PS/NS1 /1-Ag4-1 myeloma cell line			
Specificity	Mouse anti Human CD10 antibody, clone SN5c recognizes human neprilysin, also known as CD10, atriopeptidase, enkephalinase, neutral endopeptidase 24.11, skin fibroblast elastase or common acute lymphocytic leukemia antigen (CALLA). CD10 is a 749 aminoacid, ~100 kDa single pass type II transmembrane glycoprotein expressed by acute lymphoblastic leukaemia cells and by peripheral blood granulocytes. Defects in the MME gene encoding CD10 can lead to the development of the peripheral nervous system disorder, Charcot-Marie-Tooth disease 2T (CMY2T), an axonal form of Marie-Charcot-Tooth disease characterized by either dominantly inherited primary peripheral demyelinating neuropathies, designated CMT1, or primary peripheral axonal neuropathies showing axonal degeneration in the absence of any obvious myelin alteration (CMT2).			
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.			
References	 Biddle, W.C. <i>et al.</i> (1989) <i>In vitro</i> and <i>in vivo</i> cytotoxic activity of anti-human leukemia monoclonal antibodies SN5c and SN6 daunorubicin conjugates. <u>Leuk Res. 13 (8): 699-707.</u> Riemann, D. <i>et al.</i> (2001) Caveolae/lipid rafts in fibroblast-like synoviocytes: ectopeptidase-rich membrane microdomains. <u>Biochem J. 354 (Pt 1): 47-55.</u> Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect <i>bona fide</i> dedifferentiation rather than amplification of progenitor cells. <u>J Cell Physiol. 214: 75-83.</u> Pilling, D. <i>et al.</i> (2009) Identification of markers that distinguish monocyte-derived fibrocytes from monocytes, macrophages, and fibroblasts. <u>PLoS One. 4 (10): e7475.</u> Manini, I. <i>et al.</i> (2020) Heterogeneity Matters: Different Regions of Glioblastoma Are Characterized by Distinctive Tumor-Supporting Pathways. <u>Cancers (Basel). 12 (10)Oct 13 [Epub ahead of print].</u> Glynn, E. & Fromm, J.R. (2020) Immunophenotypic Characterization and Purification of Neoplastic Cells from Lymph Nodes Involved by T-Cell/Histiocyte-rich Large B-cell Lymphoma by Flow Cytometry and Flow Cytometric Cell Sorting. <u>Cytometry B Clin Cytom.</u> 			

7. Caponnetto, F. et al. (2020) Human Adipose-Derived Stem Cells in Madelung's

98 (1): 88-98.

8. Kirolos, S.A. and Gomer, R.H. (2022) The extracellular sialidase NEU3 induces neutrophil priming bioRxiv. 24 Feb [Epub ahead of print]. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Guarantee 12 months from date of despatch **Acknowledgements** This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts **Health And Safety** Material Safety Datasheet documentation #20471 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA1556SBB580 20471 Regulatory For research purposes only

Disease: Morphological and Functional Characterization. Cells. 10 (1): 44.

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M410097:221024'

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com