

Datasheet: MCA1556SBV670

Description:	MOUSE ANTI HUMAN CD10:StarBright Violet 670
Specificity:	CD10
Other names:	CALLA
Format:	StarBright Violet 670
Product Type:	Monoclonal Antibody
Clone:	SN5c
Isotype:	IgG1
Quantity:	100 TESTS/0.5ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human		
Product Form	Purified IgG conjugate	ed to StarBright Violet	670 - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm
	StarBright Violet 670	401	667
Preparation	Purified IgG prepared supernatant	l by affinity chromatogi	raphy on Protein A
·	•		raphy on Protein A
Buffer Solution	supernatant	aline	raphy on Protein A t
Buffer Solution Preservative Stabilisers	supernatant Phosphate buffered s	aline (NaN ₃)	raphy on Protein A t
Buffer Solution Preservative	Phosphate buffered s 0.09% Sodium Azide	aline (NaN ₃)	raphy on Protein A
Buffer Solution Preservative	Phosphate buffered s 0.09% Sodium Azide 1% Bovine Serum Alk	aline (NaN ₃)	raphy on Protein A t

Approx. Protein Concentrations

For information on the concentration of our StarBright Dye conjugated reagents please visit our FAQ page.

Immunogen

Partially purified cell membrane antigens from fresh leukemia cells

External Database

Links

UniProt:

P08473 Related reagents

Entrez Gene:

4311 MME Related reagents

Synonyms

EPN

Fusion Partners

Spleen cells from immunized BALB/c mice were fused with cells of the mouse PS/NS1 /1-Ag4-1 myeloma cell line

Specificity

Mouse anti Human CD10 antibody, clone SN5c recognizes human neprilysin, also known as CD10, atriopeptidase, enkephalinase, neutral endopeptidase 24.11, skin fibroblast elastase or common acute lymphocytic leukemia antigen (CALLA). CD10 is a 749 aminoacid, ~100 kDa single pass type II transmembrane glycoprotein expressed by acute lymphoblastic leukaemia cells and by peripheral blood granulocytes.

Defects in the MME gene encoding CD10 can lead to the development of the peripheral nervous system disorder, Charcot-Marie-Tooth disease 2T (CMY2T), an axonal form of Marie-Charcot-Tooth disease characterized by either dominantly inherited primary peripheral demyelinating neuropathies, designated CMT1, or primary peripheral axonal neuropathies showing axonal degeneration in the absence of any obvious myelin alteration (CMT2).

Flow Cytometry

Use 5μ I of the suggested working dilution to label 10^6 cells in 100μ I. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.

References

- 1. Biddle, W.C. *et al.* (1989) *In vitro* and *in vivo* cytotoxic activity of anti-human leukemia monoclonal antibodies SN5c and SN6 daunorubicin conjugates. <u>Leuk Res. 13 (8):</u> 699-707.
- 2. Riemann, D. *et al.* (2001) Caveolae/lipid rafts in fibroblast-like synoviocytes: ectopeptidase-rich membrane microdomains. Biochem J. 354 (Pt 1): 47-55.
- 3. Diaz-Romero, J. *et al.* (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect *bona fide* dedifferentiation rather than amplification of progenitor cells. J Cell Physiol. 214: 75-83.
- 4. Pilling, D. *et al.* (2009) Identification of markers that distinguish monocyte-derived fibrocytes from monocytes, macrophages, and fibroblasts. <u>PLoS One. 4 (10): e7475.</u>
- 5. Manini, I. *et al.* (2020) Heterogeneity Matters: Different Regions of Glioblastoma Are Characterized by Distinctive Tumor-Supporting Pathways. <u>Cancers (Basel). 12 (10)Oct 13 [Epub ahead of print].</u>
- 6. Glynn, E. & Fromm, J.R. (2020) Immunophenotypic Characterization and Purification of Neoplastic Cells from Lymph Nodes Involved by T-Cell/Histiocyte-rich Large B-cell

Lymphoma by Flow Cytometry and Flow Cytometric Cell Sorting. Cytometry B Clin Cytom. 98 (1): 88-98. 7. Caponnetto, F. et al. (2020) Human Adipose-Derived Stem Cells in Madelung's Disease: Morphological and Functional Characterization. Cells. 10 (1): 44. 8. Kirolos, S.A. and Gomer, R.H. (2022) The extracellular sialidase NEU3 induces neutrophil priming bioRxiv. 24 Feb [Epub ahead of print]. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Guarantee 12 months from date of despatch **Acknowledgements** This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts **Health And Safety** Material Safety Datasheet documentation #20471 available at: Information https://www.bio-rad-antibodies.com/SDS/MCA1556SBV670

Related Products

Regulatory

America

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

Fax: +1 919 878 3751

North & South Tel: +1 800 265 7376

Worldwide

For research purposes only

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21

Email: antibody sales us@bio-rad.com

20471

Email: antibody_sales_uk@bio-rad.com

Fax: +49 (0) 89 8090 95 50
Email: antibody sales de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M435027:250224'

Printed on 15 Apr 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint