

Datasheet: MCA1556SBY575

Description:	MOUSE ANTI HUMAN CD10:StarBright Yellow 575			
Specificity:	CD10			
Other names:	CALLA			
Format:	StarBright Yellow 575			
Product Type:	Monoclonal Antibody			
Clone:	SN5c			
Isotype:	lgG1			
Quantity:	100 TESTS/0.5ml			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			Neat	
	Where this product has necessarily exclude its a guide only. It is recon system using appropria	use in such pr nmended that t	ocedur the use	es. Suggested workin titrates the product	ng dilutions are given as	
Target Species	Human					
Product Form	Purified IgG conjugated to StarBright Yellow 575 - liquid					
Max Ex/Em		Excitation Max	(nm)	Emission Max (nm)		
	StarBright Yellow 575	548		579		
Preparation	Purified IgG prepared b supernatant	by affinity chror	matogra	aphy on Protein A froi	m tissue culture	
Buffer Solution	Phosphate buffered saline					
Preservative	0.09% Sodium Azide (NaN ₃)					
Stabilisers	1% Bovine Serum Albumin					
	0.1% Pluronic F68					
	0.1% PEG 3350					
	0.05% Tween 20					

Immunogen	Partially purified cell membrane antigens from fresh leukemia cells
External Database Links	UniProt: P08473 Related reagents Entrez Gene: <u>4311</u> MME Related reagents
Synonyms	EPN
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse PS/NS1 /1-Ag4-1 myeloma cell line
Specificity	Mouse anti Human CD10 antibody, clone SN5c recognizes human neprilysin, also known as CD10, atriopeptidase, enkephalinase, neutral endopeptidase 24.11, skin fibroblast elastase or common acute lymphocytic leukemia antigen (CALLA). CD10 is a 749 aminoacid, ~100 kDa single pass type II transmembrane glycoprotein expressed by acute lymphoblastic leukaemia cells and by peripheral blood granulocytes. Defects in the MME gene encoding CD10 can lead to the development of the peripheral nervous system disorder, Charcot-Marie-Tooth disease 2T (CMY2T), an axonal form of Marie-Charcot-Tooth disease characterized by either dominantly inherited primary peripheral demyelinating neuropathies, designated CMT1, or primary peripheral axonal neuropathies showing axonal degeneration in the absence of any obvious myelin alteration (CMT2).
Flow Cytometry	Use 5µl of the suggested working dilution to label 10 ⁶ cells in 100µl. Best practices suggest a 5 minutes centrifugation at 6,000g prior to sample application.
References	 Biddle, W.C. <i>et al.</i> (1989) <i>In vitro</i> and <i>in vivo</i> cytotoxic activity of anti-human leukemia monoclonal antibodies SN5c and SN6 daunorubicin conjugates. Leuk Res. 13 (8): 699-707. Riemann, D. <i>et al.</i> (2001) Caveolae/lipid rafts in fibroblast-like synoviocytes: ectopeptidase-rich membrane microdomains. <u>Biochem J. 354 (Pt 1): 47-55.</u> Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect <i>bona fide</i> dedifferentiation rather than amplification of progenitor cells. <u>J Cell Physiol. 214: 75-83.</u> Pilling, D. <i>et al.</i> (2009) Identification of markers that distinguish monocyte-derived fibrocytes from monocytes, macrophages, and fibroblasts. <u>PLoS One. 4 (10): e7475.</u> Manini, I. <i>et al.</i> (2020) Heterogeneity Matters: Different Regions of Glioblastoma Are Characterized by Distinctive Tumor-Supporting Pathways. <u>Cancers (Basel). 12 (10)Oct 13</u> [Epub ahead of print]. Glynn, E. & Fromm, J.R. (2020) Immunophenotypic Characterization and Purification of Neoplastic Cells from Lymph Nodes Involved by T-Cell/Histiocyte-rich Large B-cell Lymphoma by Flow Cytometry and Flow Cytometric Cell Sorting. <u>Cytometry B Clin Cytom. 98 (1): 88-98.</u> Caponnetto, F. <i>et al.</i> (2020) Human Adipose-Derived Stem Cells in Madelung's

	Disease: Morphological and Functional Characterization. <u>Cells. 10 (1): 44.</u> 8. Kirolos, S.A. and Gomer, R.H. (2022) The extracellular sialidase NEU3 induces neutrophil priming <u>bioRxiv. 24 Feb [Epub ahead of print].</u>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted.
Guarantee	12 months from date of despatch
Acknowledgements	This product is covered by U.S. Patent No. 10,150,841 and related U.S. and foreign counterparts
Health And Safety Information	Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/MCA1556SBY575 20471
Regulatory	For research purposes only

Related Products

Recommended Useful Reagents

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-r	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M413707:221125'

Printed on 08 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint