

Datasheet: MCA1556XZ

Description:	MOUSE ANTI HUMAN CD10:Preservative Free		
Specificity:	CD10		
Other names:	CALLA		
Format:	Preservative Free		
Product Type:	Monoclonal Antibody		
Clone:	SN5c		
Isotype:	lgG1		
Quantity:	1 mg		

## **Product Details**

# **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			
Immunohistology - Frozen	•			
Immunohistology - Paraffin				
ELISA				
Immunoprecipitation	•			
Western Blotting	-			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein A supernatant	A from tissue culture
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	None present	

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Partially purified cell membrane antigens from fresh leukemia cells
External Database Links	UniProt: P08473 Related reagents  Entrez Gene: 4311 MME Related reagents
Synonyms	EPN
RRID	AB_324601
Fusion Partners	Spleen cells from immunized BALB/c mice were fused with cells of the mouse PS/NS1 /1-Ag4-1 myeloma cell line
Specificity	Mouse anti Human CD10 antibody, clone SN5c recognizes human neprilysin, also known as CD10, atriopeptidase, enkephalinase, neutral endopeptidase 24.11, skin fibroblast elastase or common acute lymphocytic leukemia antigen (CALLA). CD10 is a 749 aminoacid, ~100 kDa single pass type II transmembrane glycoprotein expressed by acute lymphoblastic leukaemia cells and by peripheral blood granulocytes.  Defects in the MME gene encoding CD10 can lead to the development of the peripheral nervous system disorder, Charcot-Marie-Tooth disease 2T (CMY2T), an axonal form of Marie-Charcot-Tooth disease characterized by either dominantly inherited primary peripheral demyelinating neuropathies, designated CMT1, or primary peripheral axonal neuropathies showing axonal degeneration in the absence of any obvious myelin alteration (CMT2).
Flow Cytometry	Use 10µl of the suggested working dilution to label 10 <sup>6</sup> cells in 100µl
References	<ol> <li>Biddle, W.C. <i>et al.</i> (1989) <i>In vitro</i> and <i>in vivo</i> cytotoxic activity of anti-human leukemia monoclonal antibodies SN5c and SN6 daunorubicin conjugates. <u>Leuk Res. 13 (8): 699-707.</u></li> <li>Riemann, D. <i>et al.</i> (2001) Caveolae/lipid rafts in fibroblast-like synoviocytes: ectopeptidase-rich membrane microdomains. <u>Biochem J. 354 (Pt 1): 47-55.</u></li> <li>Diaz-Romero, J. <i>et al.</i> (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect <i>bona fide</i> dedifferentiation rather than amplification of progenitor cells. <u>J Cell Physiol. 214: 75-83.</u></li> <li>Pilling, D. <i>et al.</i> (2009) Identification of markers that distinguish monocyte-derived fibrocytes from monocytes, macrophages, and fibroblasts. <u>PLoS One. 4 (10): e7475.</u></li> <li>Manini, I. <i>et al.</i> (2020) Heterogeneity Matters: Different Regions of Glioblastoma Are Characterized by Distinctive Tumor-Supporting Pathways. <u>Cancers (Basel). 12 (10)Oct 13</u></li> </ol>

#### [Epub ahead of print].

- 6. Glynn, E. & Fromm, J.R. (2020) Immunophenotypic Characterization and Purification of Neoplastic Cells from Lymph Nodes Involved by T-Cell/Histiocyte-rich Large B-cell Lymphoma by Flow Cytometry and Flow Cytometric Cell Sorting. Cytometry B Clin Cytom. 98 (1): 88-98.
- 7. Caponnetto, F. *et al.* (2020) Human Adipose-Derived Stem Cells in Madelung's Disease: Morphological and Functional Characterization. Cells. 10 (1): 44.
- 8. Kirolos, S.A. and Gomer, R.H. (2022) The extracellular sialidase NEU3 induces neutrophil priming bioRxiv. 24 Feb [Epub ahead of print].

Storage Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety
Information

Material Safety Datasheet documentation #10162 available at:
<a href="https://www.bio-rad-antibodies.com/SDS/MCA1556XZ">https://www.bio-rad-antibodies.com/SDS/MCA1556XZ</a>
10162

Regulatory

For research purposes only

## Related Products

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (STAR77...) HRP

Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP
Goat Anti Mouse IgG (STAR70...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Rabbit Anti Mouse IgG (STAR9...) FITC

#### **Recommended Negative Controls**

#### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M410083:221024'

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint