

Datasheet: MCA1557T BATCH NUMBER 151041

Description:	MOUSE ANTI HUMAN CD105	
Specificity:	CD105	
Other names:	ENDOGLIN	
Format:	Purified	
Product Type:	Monoclonal Antibody	
Clone:	SN6	
Isotype:	lgG1	
Quantity:	25 μg	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	-			1/10 - 1/50
Immunohistology - Frozen (1)	-			
Immunohistology - Paraffin				
ELISA			•	
Immunoprecipitation	-			
Western Blotting	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1) The epitope recognised by this antibody is reported to be sensitive to formal dehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

Target Species	Human
Species Cross Reactivity	Reacts with: Horse, Cynomolgus monkey, Rhesus Monkey Based on sequence similarity, is expected to react with:Primate

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for

further information. **Product Form** Purified IgG - liquid Preparation Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant **Buffer Solution**

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide

Carrier Free

Yes

Approx. Protein Concentrations

IgG concentration 1.0 mg/ml

Immunogen

Partially purified cell membrane antigens from fresh leukemia cells

External Database Links

UniProt:

P17813 Related reagents

Entrez Gene:

2022 ENG Related reagents

Synonyms

END

RRID

AB 1100559

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the mouse P3/NS1 /1-Ag4-1 myeloma cell line

Specificity

Mouse anti Human CD105 antibody, clone SN6 recognizes the human endoglin, also known as CD105. CD105 is a glycoprotein homodimer of ~95 kDa subunits expressed by endothelial cells, activated monocytes and some leukemia cells.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

- 1. Hauser, P.V. et al. (2010) Stem cells derived from human amniotic fluid contribute to acute kidney injury recovery. Am J Pathol. 177: 2011-21.
- 2. Jin, H.J. et al. (2010) GD2 expression is closely associated with neuronal differentiation of human umbilical cord blood-derived mesenchymal stem cells. Cell Mol Life Sci. 67 (11):
- 3. Nagano, M. et al. (2007) Identification of functional endothelial progenitor cells suitable for the treatment of ischemic tissue using human umbilical cord blood. Blood 110 (1):
- 4. Braun, J. et al. (2010) Evaluation of the osteogenic and chondrogenic differentiation capacities of equine adipose tissue-derived mesenchymal stem cells. Am J Vet Res. 71

(10): 1228-36.

- 5. Diaz-Romero, J. *et al.* (2008) Immunophenotypic changes of human articular chondrocytes during monolayer culture reflect bona fide dedifferentiation rather than amplification of progenitor cells. J Cell Physiol. 214: 75-83.
- 6. Agha-Hosseini, F. *et al.* (2010) *In vitro* isolation of stem cells derived from human dental pulp. <u>Clin Transplant. 24: E23-8.</u>
- 7. Arufe, M.C. *et al.* (2010) Chondrogenic potential of subpopulations of cells expressing mesenchymal stem cell markers derived from human synovial membranes. <u>J Cell Biochem. 111: 834-45.</u>
- 8. Balmayor, E.R. *et al.* (2011) Synthesis and functionalization of superparamagnetic poly-ε-caprolactone microparticles for the selective isolation of subpopulations of human adipose-derived stem cells. <u>J R Soc Interface 8: 896-908.</u>
- 9. Benetti, A. *et al.* (2008) Transforming growth factor-beta1 and CD105 promote the migration of hepatocellular carcinoma-derived endothelium. Cancer Res. 68: 8626-34.
- 10. Ciccocioppo, R. *et al.* (2011) Autologous bone marrow-derived mesenchymal stromal cells in the treatment of fistulising Crohn's disease. Gut 60: 788-98.
- 11. Cox, G. *et al.* (2011) The use of the reamer-irrigator-aspirator to harvest mesenchymal stem cells. <u>J Bone Joint Surg Br. 93: 517-24.</u>
- 12. Ferro, F. *et al.* (2010) Biochemical and biophysical analyses of tissue-engineered bone obtained from three-dimensional culture of a subset of bone marrow mesenchymal stem cells. Tissue Eng Part A 16: 3657-67.
- 13. Lozanoska-Ochser, B. *et al.* (2008) Expression of CD86 on human islet endothelial cells facilitates T cell adhesion and migration. J Immunol. 181: 6109-16.
- 14. Sallustio, F. *et al.* (2010) TLR2 plays a role in the activation of human resident renal stem/progenitor cells. FASEB J. 24: 514-25.
- 15. Tso, C. *et al.* (2012) Phenotypic and functional changes in blood monocytes following adherence to endothelium. PLoS One 7: e37091.
- 16. Hu, N. *et al.* (2013) Long-term outcome of the repair of 50 mm long median nerve defects in rhesus monkeys with marrow mesenchymal stem cells-containing, chitosan-based tissue engineered nerve grafts. <u>Biomaterials 34: 100-11.</u>
- 17. Cho, H.J. *et al.* (2013) Generation of human secondary cardiospheres as a potent cell processing strategy for cell-based cardiac repair. Biomaterials 34: 651-61.
- 18. Kang, S.D. *et al.* (2013) Isolation of Functional Human Endothelial Cells from Small Volumes of Umbilical Cord Blood. <u>Ann Biomed Eng. 41: 2181-92.</u>
- 19. Mehrkens, A. *et al.* (2013) Non-adherent mesenchymal progenitors from adipose tissue stromal vascular fraction. <u>Tissue Eng Part A 20: 1081-8.</u>
- 20. De Schauwer, C. *et al.* (2012) In search for cross-reactivity to immunophenotype equine mesenchymal stromal cells by multicolor flow cytometry. Cytometry A 81: 312-23.
- 21. Zhang, J. *et al.* (2016) Bone mesenchymal stem cells differentiate into myofibroblasts in the tumor microenvironment. <u>Oncol Lett. 12 (1): 644-50.</u>
- 22. Morsing, M. *et al.* (2016) Evidence of two distinct functionally specialized fibroblast lineages in breast stroma. <u>Breast Cancer Res. 18 (1): 108.</u>
- 23. Williamson, K.A. *et al.* (2015) Restricted differentiation potential of progenitor cell populations obtained from the equine superficial digital flexor tendon (SDFT). <u>J Orthop</u> Res. 33 (6): 849-58.
- 24. Lützkendorf, J. *et al.* (2017) Resistance for Genotoxic Damage in Mesenchymal Stromal Cells Is Increased by Hypoxia but Not Generally Dependent on p53-Regulated

Cell Cycle Arrest. PLoS One. 12 (1): e0169921.

- 25. Lee, H.J. *et al.* (2017) ICOSL expression in human bone marrow-derived mesenchymal stem cells promotes induction of regulatory T cells. <u>Sci Rep. 7: 44486.</u>
- 26. Yi, T. *et al.* (2015) Manufacture of Clinical-Grade Human Clonal Mesenchymal Stem Cell Products from Single Colony Forming Unit-Derived Colonies Based on the Subfractionation Culturing Method. Tissue Eng Part C Methods. 21 (12): 1251-62.
- 27. Boccardo, S. *et al.* (2016) Engineered mesenchymal cell-based patches as controlled VEGF delivery systems to induce extrinsic angiogenesis. <u>Acta Biomater. 42: 127-35.</u>
- 28. Mumaw, J.L. *et al.* (2015) Feline mesenchymal stem cells and supernatant inhibit reactive oxygen species production in cultured feline neutrophils. Res Vet Sci. 103: 60-9.
- 29. Bertolo, A. *et al.* (2017) Oxidative status predicts quality in human mesenchymal stem cells. <u>Stem Cell Res Ther. 8 (1): 3.</u>
- 30. GarikipatiV, N.S. *et al.* (2018) Isolation and characterization of mesenchymal stem cells from human fetus heart. PLoS One. 13 (2): e0192244.
- 31. Olimpio, R.M.C. *et al.* (2018) Cell viability assessed in a reproducible model of human osteoblasts derived from human adipose-derived stem cells. PLoS One. 13 (4): e0194847.
- 32. Lotfi, R. *et al.* (2018) ATP promotes immunosuppressive capacities of mesenchymal stromal cells by enhancing the expression of indoleamine dioxygenase. <u>Immun Inflamm Dis. Oct 10 [Epub ahead of print].</u>
- 33. Santos, V.H.D. *et al.* (2019) Evaluation of alginate hydrogel encapsulated mesenchymal stem cell migration in horses. Res Vet Sci. 124: 38-45.
- 34. Noda, S. *et al.* (2019) Effect of cell culture density on dental pulp-derived mesenchymal stem cells with reference to osteogenic differentiation. Sci Rep. 9 (1): 5430.
- 35. Manini, I. *et al.* (2020) Heterogeneity Matters: Different Regions of Glioblastoma Are Characterized by Distinctive Tumor-Supporting Pathways. <u>Cancers (Basel). 12 (10) Oct 13 [Epub ahead of print].</u>
- 36. Cargnoni, A. *et al.* (2020) Amniotic MSCs reduce pulmonary fibrosis by hampering lung B-cell recruitment, retention, and maturation. <u>Stem Cells Transl Med. 9 (9): 1023-35.</u>
- 37. Huang, Q. *et al.* (2021) Human Umbilical Cord Mesenchymal Stem Cells-Derived Exosomal MicroRNA-18b-3p Inhibits the Occurrence of Preeclampsia by Targeting LEP. Nanoscale Res Lett. 16 (1): 27.
- 38. Piñeiro-Ramil, M. *et al.* (2020) Immortalizing Mesenchymal Stromal Cells from Aged Donors While Keeping Their Essential Features. <u>Stem Cells Int. 2020: 5726947.</u>
- 39. Kim, M. *et al.* (2020) A Small-Sized Population of Human Umbilical Cord Blood-Derived Mesenchymal Stem Cells Shows High Stemness Properties and Therapeutic Benefit. Stem Cells Int. 2020: 5924983.
- 40. Niu, C.C. *et al.* (2014) Identification of mesenchymal stem cells and osteogenic factors in bone marrow aspirate and peripheral blood for spinal fusion by flow cytometry and proteomic analysis. <u>J Orthop Surg Res. 9: 32.</u>
- 41. Supokawej, A. *et al.* (2013) Cardiogenic and myogenic gene expression in mesenchymal stem cells after 5-azacytidine treatment. <u>Turk J Haematol. 30 (2): 115-21.</u>

Further Reading

- 1. Burk, J. *et al.* (2013) Equine cellular therapy--from stall to bench to bedside? <u>Cytometry</u> A 83 (1): 103-13.
- 2. Carrade, D.D. *et al.* (2012) Comparative Analysis of the Immunomodulatory Properties of Equine Adult-Derived Mesenchymal Stem Cells. <u>Cell Med. 4: 1-11.</u>

Storage	Store at +4°C or at -20°C if preferred.
	This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1557T
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP Goat Anti Mouse IgG (STAR70...) **FITC** Rabbit Anti Mouse IgG (STAR13...) HRP Rabbit Anti Mouse IgG (STAR9...) **FITC** Goat Anti Mouse IgG (STAR77...) **HRP** Goat Anti Mouse IgG (STAR76...) **RPE** Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550, DyLight®650, DyLight®680, DyLight®800, FITC, HRP Rabbit Anti Mouse IgG (STAR12...) **RPE**

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365440:200529'

Printed on 27 Nov 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint