

## Datasheet: MCA1558

**BATCH NUMBER 160557**

<b>Description:</b>	MOUSE ANTI PCNA
<b>Specificity:</b>	PCNA
<b>Other names:</b>	PROLIFERATING CELL NUCLEAR ANTIGEN
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	PC10
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry (1)	▪			1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			1/50 - 1/100
ELISA			▪	
Immunoprecipitation	▪			
Western Blotting	▪			1/1000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**(1) Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm™ (Product Code [BUF09](#)) for this purpose.**

<b>Target Species</b>	Rat
<b>Species Cross Reactivity</b>	<p>Reacts with: Ferret, Chicken, Rabbit, Xenopus, Mouse, Horse, Sheep, Dog, Cat, Cynomolgus monkey, Rhesus Monkey, Hamster, Atlantic Salmon, Human, Bearded Dragon, Corn Snake, Nile Crocodile</p> <p>Based on sequence similarity, is expected to react with: Vertebrates, Invertebrates</p> <p><b>N.B.</b> Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for</p>

further information.

<b>Product Form</b>	Purified IgG - liquid
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Rat PCNA made in the protein A expression vector pR1T2T
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">P04961</a> <a href="#">Related reagents</a>  <b>Entrez Gene:</b> <a href="#">25737</a> Pcna <a href="#">Related reagents</a>
<b>RRID</b>	AB_324955
<b>Specificity</b>	<p><b>Mouse anti PCNA antibody, clone PC10</b> recognizes the proliferating cell nuclear antigen, also known as PCNA or cyclin. PCNA is a 261 amino acid ~28 kDa nuclear protein vital for cellular DNA synthesis at the replication fork (<a href="#">Li et al. 1995</a>) through its interaction with <a href="#">FEN1</a> (<a href="#">Wu et al. 1996</a>). PCNA is the auxilliary protein for DNA polymerase <math>\delta</math> (<a href="#">Bravo et al. 1987</a>).</p> <p>PCNA is highly conserved between mammalian species and other vertebrates. Mouse anti PCNA antibody, clone PC10 has been used for the detection of PCNA in a number of species including human, rat, mouse (<a href="#">Park et al. 2008</a>), chicken (<a href="#">Franz-Odendaal 2008</a>) and abalone (<a href="#">Harris et al. 2006</a>).</p>
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul.
<b>Immunohistology</b>	This product does not require protein digestion pre-treatment of paraffin sections. This product does not require antigen retrieval using heat pre-treatment prior to staining of paraffin sections.
<b>Histology Positive Control Tissue</b>	Tonsil
<b>References</b>	<ol style="list-style-type: none"><li>Mathews, M.B. <i>et al.</i> (1984) Identity of the proliferating cell nuclear antigen and cyclin. <a href="#">Nature. 309 (5966): 374-6.</a></li><li>Ogata, K. <i>et al.</i> (1985) Purification and N-terminal amino acid sequence of proliferating cell nuclear antigen (PCNA)/cyclin and development of ELISA for anti-PCNA antibodies. <a href="#">J</a></li></ol>

[Immunol. 135 \(4\): 2623-7.](#)

3. Jenkins, H. *et al.* (1993) Nuclei that lack a lamina accumulate karyophilic proteins and assemble a nuclear matrix. [J Cell Sci. 106: 275-85.](#)
4. Landberg, G. *et al.* (1990) Flow cytometric multiparameter analysis of proliferating cell nuclear antigen/cyclin and Ki-67 antigen: a new view of the cell cycle. [Exp Cell Res. 187 \(1\): 111-8.](#)
5. Wilson, G.D. *et al.* (1992) Flow cytometric characterisation of proliferating cell nuclear antigen using the monoclonal antibody PC10. [Eur J Cancer. 28A \(12\): 2010-7.](#)
6. Prosperi, E. *et al.* (1993) Proliferating cell nuclear antigen complex formation induced by ultraviolet irradiation in human quiescent fibroblasts as detected by immunostaining and flow cytometry. [Exp Cell Res. 205 \(2\): 320-5.](#)
7. Elsässer, H.P. *et al.* (1994) Growth of rat pancreatic acinar cells quantitated with a monoclonal antibody against the proliferating cell nuclear antigen. [Cell Tissue Res. 276 \(3\): 603-9.](#)
8. Harris, L. *et al.* (2006) Characterisation of cell types in abalone (*Haliotis* spp.) tissues using immunohistochemical techniques [Aquaculture 261: 1413-21](#)
9. Buggins, A.G. *et al.* (2001) Microenvironment produced by acute myeloid leukemia cells prevents T cell activation and proliferation by inhibition of NF-kappaB, c-Myc, and pRb pathways. [J Immunol. 167: 6021-30.](#)
10. Kapitonova, M.Y. *et al.* (2010) Immunohistochemical characteristics of the hypophysis in normal conditions and chronic stress. [Neurosci Behav Physiol. 40: 97-102.](#)
11. Franz-Odenaal, T.A. (2008) Toward understanding the development of scleral ossicles in the chicken, *Gallus gallus*. [Dev Dyn. 237: 3240-51.](#)
12. Hashimoto, Y. *et al.* (2010) Rad51 protects nascent DNA from Mre11-dependent degradation and promotes continuous DNA synthesis. [Nat Struct Mol Biol. 17: 1305-11.](#)
13. Fenton, M. *et al.* (2001) Cellular senescence after single and repeated balloon catheter denudations of rabbit carotid arteries. [Arterioscler Thromb Vasc Biol. 21: 220-6.](#)
14. Park, J.H. *et al.* (2008) Gastric lesions and immune responses caused by long-term infection with *Helicobacter heilmannii* in C57BL/6 mice. [J Comp Pathol. 139: 208-17.](#)
15. Izhak, L. *et al.* (2012) Dissecting the autocrine and paracrine roles of the CCR2-CCL2 axis in tumor survival and angiogenesis. [PLoS One. 7: e28305.](#)
16. Di-poï, N. & Milinkovitch, M.C. (2016) The anatomical placode in reptile scale morphogenesis indicates shared ancestry among skin appendages in amniotes. [Sci Adv. 2 \(6\): e1600708.](#)
17. Guzera, M. *et al.* (2016) *In Vitro* Influence of Mycophenolic Acid on Selected Parameters of Stimulated Peripheral Canine Lymphocytes. [PLoS One. 11 \(5\): e0154429.](#)
18. Khlebnikov, V. V. *et al.* (2015) Developmental Characteristics of the Hypothalamo-Hypophyseal-Adrenal System in Chronic Heterotypical Stress [Neuroscience and Behavioral Physiology. 46 \(1\): 100-5.](#)
19. Nakatsuka, M. & Kumabe, S (2018) Histological Alterations from Condyle Repositioning with Functional Appliances in Rats. [J Clin Pediatr Dent. 42 \(5\): 391-7.](#)
20. Falbo, L. *et al.* (2020) SSRP1-mediated histone H1 eviction promotes replication origin assembly and accelerated development. [Nat Commun. 11 \(1\): 1345.](#)

---

#### Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1558">https://www.bio-rad-antibodies.com/SDS/MCA1558</a> 10040
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	<a href="#">RPE</a>
Goat Anti Mouse IgG IgA IgM (STAR87...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (STAR76...)	<a href="#">RPE</a>
Goat Anti Mouse IgG (STAR70...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (H/L) (STAR117...)	<a href="#">Alk. Phos.</a> , <a href="#">DyLight®488</a> , <a href="#">DyLight®550</a> , <a href="#">DyLight®650</a> , <a href="#">DyLight®680</a> , <a href="#">DyLight®800</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR9...)	<a href="#">FITC</a>
Goat Anti Mouse IgG (STAR77...)	<a href="#">HRP</a>
Goat Anti Mouse IgG (Fc) (STAR120...)	<a href="#">FITC</a> , <a href="#">HRP</a>
Rabbit Anti Mouse IgG (STAR13...)	<a href="#">HRP</a>

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA1210\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M386206:210519'

Printed on 29 Feb 2024