

Datasheet: MCA1568EL

#### **BATCH NUMBER 158942**

Description:	MOUSE ANTI HUMAN CD14:Low Endotoxin		
Specificity:	CD14		
Format:	Low Endotoxin		
Product Type:	Monoclonal Antibody		
Clone:	TÜK4		
Isotype:	lgG2a		
Quantity:	0.5 mg		

# **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/100 - 1/200
Immunohistology - Frozen			•	
Immunohistology - Paraffin			•	
ELISA				
Immunoprecipitation			•	
Western Blotting	•			
Functional Assays				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Dog, Goat, Cat, Rabbit, Mink, Bovine, Pig, Sheep, Cynomolgus monkey, Llama  N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	None present				
Carrier Free	Yes				
Endotoxin Level	< 0.01 EU/ug				
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml				
External Database Links	UniProt: P08571 Related reagents				
	Entrez Gene:				
	929 CD14 Related reagents				
RRID	AB_566516				
Specificity	<b>Mouse anti human CD14 antibody, clone TÜK4</b> recognizes the human CD14 cell surface antigen. CD14 is a ~55 kDa glycoprotein that contains multiple leucine-rich repeats. It is anchored to the cell membrane via a glycosylphosphatidylinositol (GPI) linkage (Simmons et al. 1989), a soluble form of CD14 also exists (Bazil et al. 1986).				
	CD14 is strongly expressed on the surface of monocytes and macrophages but has also been shown to be expressed on the surface of non-myeloid cells ( <u>Jersmann 2005</u> ). CD14 functions as a pattern recognition receptor ( <u>Pugin et al. 1994</u> , <u>Dziarski et al. 1998</u> ) in innate immunity for a variety of ligands, in particular for the LPS (endotoxin) of Gram-negative bacteria.				
	Mouse anti human CD14 antibody, clone TÜK4 has been shown to block SDF-induced chemotaxis of U937 cells in a dose –dependent manner ( <u>Yang et al. 2003</u> ). Use of the <u>anti-human CD14 antibody, Low Endotoxin format</u> is recommended for this purpose.				
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood.				
References	<ol> <li>Weiss, D.J. (2001) Evaluation of proliferative disorders in canine bone marrow by use of flow cytometric scatter plots and monoclonal antibodies. <u>Vet Pathol. 38: 512-8.</u></li> <li>Gupta, V.K. <i>et al.</i> (1996) Identification of the sheep homologue of the monocyte cell surface moleculeCD14. <u>Vet Immunol Immunopathol. 51 (1-2): 89-99.</u></li> <li>Sopp, P. &amp; Howard, C.J. (1997) Cross-reactivity of monoclonal antibodies to defined human leucocyte differentiation antigens with bovine cells. <u>Vet Immunol Immunopathol. 56 (1-2): 11-25.</u></li> <li>Xiong, W. <i>et al.</i> (2010) Human Flt3L generates dendritic cells from canine peripheral</li> </ol>				

- blood precursors: implications for a dog glioma clinical trial. PLoS One. 5: e11074.
- 5. Werling, D. *et al.* (1998) Analysis of the phenotype and phagocytic activity of monocytes/macrophages from cattle infected with the bovine leukaemia virus. <u>Vet Immunol Immunopathol.</u> 62 (3): 185-95.
- 6. Yang, H. *et al.* (2003) Antibody to CD14 like CXCR4-specific antibody 12G5 could inhibit CXCR4-dependent chemotaxis and HIV Env-mediated cell fusion. <u>Immunol Lett. 88</u> (1): 27-30.
- 7. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. <u>Exp Anim. 49 (2): 97-110.</u>
- 8. Jacobsen, C.N. *et al.* (1993) Reactivities of 20 anti-human monoclonal antibodies with leucocytes from ten different animal species. <u>Vet Immunol Immunopathol.</u> 39 (4): 461-6.
- 9. Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. Vet Immunol Immunopathol. 132:109-15.
- 10. Dalli J *et al.* (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. Blood. 112 (6): 2512-9.
- 11. Lybeck, K.R. *et al.* (2009) Neutralization of interleukin-10 from CD14(+) monocytes enhances gamma interferon production in peripheral blood mononuclear cells from *Mycobacterium avium* subsp. *paratuberculosis*-infected goats. <u>Clin Vaccine Immunol. 16</u> (7): 1003-11.
- 12. Ferret-Bernard, S. *et al.* (2010) Cellular and molecular mechanisms underlying the strong neonatal IL-12 response of lamb mesenteric lymph node cells to R-848. <u>PLoS One.</u> 5: e13705.
- 13. Fulton, B.E. Jr. *et al.* (2006) Dissemination of bovine leukemia virus-infected cells from a newly infected sheep lymph node. J Virol. 80: 7873-84.
- 14. Willett, B.J. *et al.* (2007) Probing the interaction between feline immunodeficiency virus and CD134 by using the novel monoclonal antibody 7D6 and the CD134 (Ox40) ligand. <u>J Virol. 81: 9665-79.</u>
- 15. Kallapur, S.G. *et al.* (2011) Pulmonary and systemic inflammatory responses to intraamniotic IL-1α in fetal sheep. Am J Physiol Lung Cell Mol Physiol. 301 (3): L285-95.
- 16. Lund, H. *et al.* (2016) Transient Migration of Large Numbers of CD14(++) CD16(+) Monocytes to the Draining Lymph Node after Onset of Inflammation. <u>Front Immunol. 7:</u> 322.
- 17. Krueger, L.A. *et al.* (2016) Gamma delta T cells are early responders to Mycobacterium avium ssp. paratuberculosis in colostrum-replete Holstein calves. <u>J Dairy</u> Sci. 99 (11): 9040-50.
- 18. Gelain, M.E. *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. <u>Vet Immunol Immunopathol. 159 (1-2): 91-6.</u>
- 19. Schaut, R.G. *et al.* (2015) Bovine viral diarrhea virus type 2 *in vivo* infection modulates TLR4 responsiveness in differentiated myeloid cells which is associated with decreased MyD88 expression. <u>Virus Res. 208: 44-55.</u>
- 20. Westover, A.J. *et al.* (2016) An Immunomodulatory Device Improves Insulin Resistance in Obese Porcine Model of Metabolic Syndrome. <u>J Diabetes Res. 2016:</u> 3486727.
- 21. Pomeroy, B. *et al.* (2017) Counts of bovine monocyte subsets prior to calving are predictive for postpartum occurrence of mastitis and metritis. <u>Vet Res. 48 (1): 13.</u>
- 22. Gibson, A.J. et al. (2016) Differential macrophage function in Brown Swiss and

- Holstein Friesian cattle. Vet Immunol Immunopathol. 181: 15-23.
- 23. Martini, V. *et al.* (2018) Flow cytometry for feline lymphoma: a retrospective study regarding pre-analytical factors possibly affecting the quality of samples. <u>J Feline Med Surg.</u> 20 (6): 494-501.
- 24. Novacco, M. *et al.* (2016) Prognostic factors in canine acute leukaemias: a retrospective study. <u>Vet Comp Oncol. 14 (4): 409-16.</u>
- 25. Feng, P.H. *et al.* (2018) S100A9<sup>+</sup> MDSC and TAM-mediated EGFR-TKI resistance in lung adenocarcinoma: the role of *RELB*. Oncotarget. 9 (7): 7631-43.
- 26. Higgins, J.L. *et al.* (2018) Cell mediated immune response in goats after experimental challenge with the virulent Brucella melitensis strain 16M and the reduced virulence strain Rev. 1. Vet Immunol Immunopathol. 202: 74-84.
- 27. Penadés, M. *et al.* (2020) Early deviations in performance, metabolic and immunological indicators affect stayability in rabbit females. <u>Animal. 14 (4): 780-9.</u>
- 28. Schwarz, E.R. *et al.* (2020) Experimental Infection of Mid-Gestation Pregnant Female and Intact Male Sheep with Zika Virus. <u>Viruses</u>. 12 (3)Mar 07 [Epub ahead of print].
- 29. Mas, A. *et al.* (2020) A further investigation of the leishmaniosis outbreak in Madrid (Spain): low-infectivity phenotype of the *Leishmania infantum* BOS1FL1 isolate to establish infection in canine cells. Vet Immunol Immunopathol. 230: 110148.
- 30. Tuohy, J.L. *et al.* (2020) Immune dysregulation and osteosarcoma: *Staphylococcus aureus*. downregulates TGF- $\beta$  and heightens the inflammatory signature in human and canine macrophages suppressed by osteosarcoma. <u>Vet Comp Oncol. 18 (1): 64-75.</u>
- 31. Sipka, A.S. *et al.* (2020) The effect of *ex vivo*. lipopolysaccharide stimulation and nutrient availability on transition cow innate immune cell AKT/mTOR pathway responsiveness. <u>J Dairy Sci. 103 (2): 1956-1968.</u>
- 32. Lessard, M. *et al.* (2018) Piglet weight gain during the first two weeks of lactation influences the immune system development. <u>Vet Immunol Immunopathol.</u> 206: 25-34.
- 33. Moncada-Saucedo, N.K. *et al.* (2019) A Bioactive Cartilage Graft of IGF1-Transduced Adipose Mesenchymal Stem Cells Embedded in an Alginate/Bovine Cartilage Matrix Tridimensional Scaffold. <u>Stem Cells Int. 2019</u>: 9792369.
- 34. Muñoz-Silvestre, A. *et al.* (2020) Pathogenesis of Intradermal Staphylococcal Infections: Rabbit Experimental Approach to Natural *Staphylococcus aureus* Skin Infections. <u>Am J Pathol. 190 (6): 1188-210.</u>
- 35. Park, D.S. *et al.* (2021) Dynamic changes in blood immune cell composition and function in Holstein and Jersey steers in response to heat stress. <u>Cell Stress Chaperones</u>. <u>26 (4): 705-20.</u>
- 36. Grudzien, M. *et al.* (2021) A newly established canine NK-type cell line and its cytotoxic properties. <u>Vet Comp Oncol. 19 (3): 567-77.</u>
- 37. Risalde, M.A. *et al.* (2020) BVDV permissiveness and lack of expression of co-stimulatory molecules on PBMCs from calves pre-infected with BVDV. <u>Comp Immunol Microbiol Infect Dis. 68: 101388.</u>
- 38. Kolar, Q.K. *et al.* (2020) Anatomical distribution of respiratory tract leukocyte cell subsets in neonatal calves. <u>Vet Immunol Immunopathol.</u> 227: 110090.
- 39. Riccardo, F. *et al.* (2022) Antigen mimicry as an effective strategy to induce CSPG4-targeted immunity in dogs with oral melanoma: a veterinary trial. <u>J Immunother Cancer.</u> 10(5):e004007. [Epub ahead of print].
- 40. Shiue, S.J. *et al.* (2022) Arthrospira Enhances Seroclearance in Patients with Chronic Hepatitis B Receiving Nucleos(t)ide Analogue through Modulation of TNF-α/IFN-γ Profile.

#### Nutrients. 14 (14)Jul 06 [Epub ahead of print].

41. Jaensch, S. *et al.* (2022) Clinicopathologic and immunophenotypic features in dogs with presumptive large granular lymphocyte leukaemia <u>Australian Veterinary Journal. 12</u> Aug [Epub ahead of print].

#### **Further Reading**

- 1. Simmons, D. L. *et al.* (1989) Monocyte antigen CD14 is a phospholipid anchored membrane protein. <u>Blood. 73:284-9.</u>
- 2. Bazil, V. *et al.* (1986) Biochemical characterization of a soluble form of the 53-kDa monocyte surface antigen. <u>Eur J Immunol. 16:1583-9.</u>
- 3. Jersmann, H.P. (2005) Time to abandon dogma: CD14 is expressed by non-myeloid lineage cells. <u>Immunol Cell Biol. 83:462-7.</u>
- 4. Pugin, J. et al. (1994) CD14 is a pattern recognition receptor. <u>Immunity.1:509-16.</u>
- 5. Dziarski, R. *et al.* (1998) Binding of bacterial peptidoglycan to CD14. <u>J Biol Chem.</u> 273:8680-90.
- 6. Piriou-Guzylack, L. (2008) Membrane markers of the immune cells in swine: an update. Vet Res. 39: 54.

#### **Storage**

Store at -20°C only.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

# Health And Safety

Guarantee

12 months from date of despatch

Material Safety Datasheet documentation #10162 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1568EL">https://www.bio-rad-antibodies.com/SDS/MCA1568EL</a>

10162

#### Regulatory

Information

For research purposes only

## Related Products

### **Recommended Secondary Antibodies**

Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (STAR70...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Goat Anti Mouse IgG (STAR77...) HRP

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP
Rabbit Anti Mouse IgG (STAR9...) FITC

# **Recommended Negative Controls**

# MOUSE IgG2a NEGATIVE CONTROL:Low Endotoxin (MCA929EL)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M365468:200529'

## Printed on 04 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint