

# Datasheet: MCA1568F BATCH NUMBER 162004

Description:	cription: MOUSE ANTI HUMAN CD14:FITC		
Specificity:	CD14		
Format:	FITC		
Product Type:	Monoclonal Antibody		
Clone:	TÜK4		
lsotype:	lgG2a		
Quantity:	0.1 mg		

## **Product Details**

Applications	derived from testing w communications from	has been reported to work in the following applications. This information is testing within our laboratories, peer-reviewed publications or personal tions from the originators. Please refer to references indicated for further				
	information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.					
			No	Not Determined	Suggested Dilution	
	Flow Cytometry	-			Neat - 1/2	
	Where this antibody has not been tested for use in a particular technique this does not					
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.					
Target Species	Human					
Species Cross Reactivity	Reacts with: Dog, Goat, Cat, Rabbit, Mink, Bovine, Pig, Sheep, Cynomolgus monkey, Llama					
	reactivity is derived from	om testing within	n our la	ons may vary between boratories, peer-revie rs. Please refer to refe	wed publications or	
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid					
Max Ex/Em	Fluorophore	Excitation Max	(nm)	Emission Max (nm)		
	FITC	490		525		
Preparation	Purified IgG prepared supernatant	by affinity chror	matogr	aphy on Protein A fron	n tissue culture	

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 0.1 mg/ml
External Database Links	UniProt: <u>P08571</u> <u>Related reagents</u> Entrez Gene: <u>929</u> CD14 <u>Related reagents</u>
RRID	AB_323712
Specificity	Mouse anti human CD14 antibody, clone TÜK4 recognizes the human CD14 cell surface antigen. CD14 is a ~55 kDa glycoprotein that contains multiple leucine-rich repeats. It is anchored to the cell membrane via a glycosylphosphatidylinositol (GPI) linkage (Simmons <i>et al.</i> 1989), a soluble form of CD14 also exists (Bazil <i>et al.</i> 1986). CD14 is strongly expressed on the surface of monocytes and macrophages but has also been shown to be expressed on the surface of non-myeloid cells (Jersmann 2005). CD14 functions as a pattern recognition receptor (Pugin <i>et al.</i> 1994, Dziarski <i>et al.</i> 1998) in innate immunity for a variety of ligands, in particular for the LPS (endotoxin) of Gram-negative bacteria.
	Mouse anti human CD14 antibody, clone TÜK4 has been shown to block SDF-induced chemotaxis of U937 cells in a dose –dependent manner ( <u>Yang <i>et al.</i> 2003</u> ). Use of the <u>anti-human CD14 antibody, Low Endotoxin format</u> is recommended for this purpose.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells or 100ul whole blood.
References	<ol> <li>Weiss, D.J. (2001) Evaluation of proliferative disorders in canine bone marrow by use of flow cytometric scatter plots and monoclonal antibodies. <u>Vet Pathol. 38: 512-8.</u></li> <li>Gupta, V.K. <i>et al.</i> (1996) Identification of the sheep homologue of the monocyte cell surface moleculeCD14. <u>Vet Immunol Immunopathol. 51 (1-2): 89-99.</u></li> <li>Sopp, P. &amp; Howard, C.J. (1997) Cross-reactivity of monoclonal antibodies to defined human leucocyte differentiation antigens with bovine cells. <u>Vet Immunol Immunopathol. 56 (1-2): 11-25.</u></li> <li>Xiong, W. <i>et al.</i> (2010) Human Flt3L generates dendritic cells from canine peripheral blood precursors: implications for a dog glioma clinical trial. <u>PLoS One. 5: e11074.</u></li> <li>Werling, D. <i>et al.</i> (1998) Analysis of the phenotype and phagocytic activity of monocytes/macrophages from cattle infected with the bovine leukaemia virus. <u>Vet Immunol Immunopathol. 62 (3): 185-95.</u></li> <li>Yang, H. <i>et al.</i> (2003) Antibody to CD14 like CXCR4-specific antibody 12G5 could inhibit CXCR4-dependent chemotaxis and HIV Env-mediated cell fusion. <u>Immunol Lett. 88</u></li> </ol>

#### (1): 27-30.

7. Yoshino, N. *et al.* (2000) Upgrading of flow cytometric analysis for absolute counts, cytokines and other antigenic molecules of cynomolgus monkeys (*Macaca fascicularis*) by using anti-human cross-reactive antibodies. <u>Exp Anim. 49 (2): 97-110.</u>

 Jacobsen, C.N. *et al.* (1993) Reactivities of 20 anti-human monoclonal antibodies with leucocytes from ten different animal species. <u>Vet Immunol Immunopathol. 39 (4): 461-6.</u>
 Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret

immunoglobulins, cytokines and CD markers. <u>Vet Immunol Immunopathol. 132:109-15.</u> 10. Dalli J *et al.* (2008) Annexin 1 mediates the rapid anti-inflammatory effects of neutrophil-derived microparticles. <u>Blood. 112 (6): 2512-9.</u>

11. Lybeck, K.R. *et al.* (2009) Neutralization of interleukin-10 from CD14(+) monocytes enhances gamma interferon production in peripheral blood mononuclear cells from *Mycobacterium avium* subsp. *paratuberculosis*-infected goats. <u>Clin Vaccine Immunol. 16</u> (7): 1003-11.

12. Ferret-Bernard, S. *et al.* (2010) Cellular and molecular mechanisms underlying the strong neonatal IL-12 response of lamb mesenteric lymph node cells to R-848. <u>PLoS One.</u> <u>5: e13705.</u>

13. Fulton, B.E. Jr. *et al.* (2006) Dissemination of bovine leukemia virus-infected cells from a newly infected sheep lymph node. <u>J Virol. 80: 7873-84.</u>

14. Willett, B.J. *et al.* (2007) Probing the interaction between feline immunodeficiency virus and CD134 by using the novel monoclonal antibody 7D6 and the CD134 (Ox40) ligand.  $\underline{J}$  <u>Virol. 81: 9665-79</u>.

15. Kallapur, S.G. *et al.* (2011) Pulmonary and systemic inflammatory responses to intraamniotic IL-1α in fetal sheep. <u>Am J Physiol Lung Cell Mol Physiol. 301 (3): L285-95.</u>
16. Lund, H. *et al.* (2016) Transient Migration of Large Numbers of CD14(++) CD16(+)

Monocytes to the Draining Lymph Node after Onset of Inflammation. <u>Front Immunol. 7:</u> <u>322.</u>

17. Krueger, L.A. *et al.* (2016) Gamma delta T cells are early responders to Mycobacterium avium ssp. paratuberculosis in colostrum-replete Holstein calves. <u>J Dairy</u> <u>Sci. 99 (11): 9040-50.</u>

18. Gelain, M.E. *et al.* (2014) CD44 in canine leukemia: analysis of mRNA and protein expression in peripheral blood. <u>Vet Immunol Immunopathol. 159 (1-2): 91-6.</u>

19. Schaut, R.G. *et al.* (2015) Bovine viral diarrhea virus type 2 *in vivo* infection modulates TLR4 responsiveness in differentiated myeloid cells which is associated with decreased MyD88 expression. <u>Virus Res. 208: 44-55.</u>

20. Westover, A.J. *et al.* (2016) An Immunomodulatory Device Improves Insulin Resistance in Obese Porcine Model of Metabolic Syndrome. <u>J Diabetes Res. 2016:</u> <u>3486727.</u>

Pomeroy, B. *et al.* (2017) Counts of bovine monocyte subsets prior to calving are predictive for postpartum occurrence of mastitis and metritis. <u>Vet Res. 48 (1): 13.</u>
 Gibson, A.J. *et al.* (2016) Differential macrophage function in Brown Swiss and Holstein Friesian cattle. <u>Vet Immunol Immunopathol. 181: 15-23.</u>

23. Martini, V. *et al.* (2018) Flow cytometry for feline lymphoma: a retrospective study regarding pre-analytical factors possibly affecting the quality of samples. <u>J Feline Med</u> <u>Surg. 20 (6): 494-501.</u>

24. Novacco, M. *et al.* (2016) Prognostic factors in canine acute leukaemias: a retrospective study. <u>Vet Comp Oncol. 14 (4): 409-16.</u>

25. Feng, P.H. *et al.* (2018) S100A9<sup>+</sup> MDSC and TAM-mediated EGFR-TKI resistance in lung adenocarcinoma: the role of *RELB*. <u>Oncotarget. 9 (7): 7631-43.</u>

26. Higgins, J.L. *et al.* (2018) Cell mediated immune response in goats after experimental challenge with the virulent Brucella melitensis strain 16M and the reduced virulence strain Rev. 1. <u>Vet Immunol Immunopathol. 202: 74-84.</u>

27. Penadés, M. *et al.* (2020) Early deviations in performance, metabolic and immunological indicators affect stayability in rabbit females. <u>Animal. 14 (4): 780-9.</u>
28. Schwarz, E.R. *et al.* (2020) Experimental Infection of Mid-Gestation Pregnant Female and Intact Male Sheep with Zika Virus. <u>Viruses. 12 (3)Mar 07 [Epub ahead of print].</u>
29. Mas, A. *et al.* (2020) A further investigation of the leishmaniosis outbreak in Madrid (Spain): low-infectivity phenotype of the *Leishmania infantum* BOS1FL1 isolate to establish infection in canine cells. <u>Vet Immunol Immunopathol. 230: 110148.</u>

30. Tuohy, J.L. *et al.* (2020) Immune dysregulation and osteosarcoma: *Staphylococcus aureus*. downregulates TGF- $\beta$  and heightens the inflammatory signature in human and canine macrophages suppressed by osteosarcoma. <u>Vet Comp Oncol. 18 (1): 64-75.</u> 31. Sipka, A.S. *et al.* (2020) The effect of *ex vivo*. lipopolysaccharide stimulation and nutrient availability on transition cow innate immune cell AKT/mTOR pathway responsiveness. <u>J Dairy Sci. 103 (2): 1956-1968.</u>

Lessard, M. *et al.* (2018) Piglet weight gain during the first two weeks of lactation influences the immune system development. <u>Vet Immunol Immunopathol. 206: 25-34.</u>
 Moncada-Saucedo, N.K. *et al.* (2019) A Bioactive Cartilage Graft of IGF1-Transduced Adipose Mesenchymal Stem Cells Embedded in an Alginate/Bovine Cartilage Matrix Tridimensional Scaffold. <u>Stem Cells Int. 2019: 9792369.</u>

34. Muñoz-Silvestre, A. *et al.* (2020) Pathogenesis of Intradermal Staphylococcal Infections: Rabbit Experimental Approach to Natural *Staphylococcus aureus* Skin Infections. <u>Am J Pathol. 190 (6): 1188-210.</u>

35. Park, D.S. *et al.* (2021) Dynamic changes in blood immune cell composition and function in Holstein and Jersey steers in response to heat stress. <u>Cell Stress Chaperones.</u> <u>26 (4): 705-20.</u>

36. Grudzien, M. *et al.* (2021) A newly established canine NK-type cell line and its cytotoxic properties. <u>Vet Comp Oncol. 19 (3): 567-77.</u>

37. Risalde, M.A. *et al.* (2020) BVDV permissiveness and lack of expression of co-stimulatory molecules on PBMCs from calves pre-infected with BVDV. <u>Comp Immunol Microbiol Infect Dis. 68: 101388.</u>

38. Kolar, Q.K. *et al.* (2020) Anatomical distribution of respiratory tract leukocyte cell subsets in neonatal calves. <u>Vet Immunol Immunopathol. 227: 110090.</u>

39. Riccardo, F. *et al.* (2022) Antigen mimicry as an effective strategy to induce CSPG4targeted immunity in dogs with oral melanoma: a veterinary trial. <u>J Immunother Cancer.</u> <u>10(5):e004007. [Epub ahead of print].</u>

40. Shiue, S.J. *et al.* (2022) Arthrospira Enhances Seroclearance in Patients with Chronic Hepatitis B Receiving Nucleos(t)ide Analogue through Modulation of TNF-α/IFN-γ Profile. <u>Nutrients. 14 (14)Jul 06 [Epub ahead of print].</u>

41. Jaensch, S. *et al.* (2022) Clinicopathologic and immunophenotypic features in dogs with presumptive large granular lymphocyte leukaemia <u>Australian Veterinary Journal. 12</u> <u>Aug [Epub ahead of print].</u>

Further Reading 1. Simmons, D. L. et al. (1989) Monocyte antigen CD14 is a phospholipid anchored

	<ul> <li>membrane protein. <u>Blood. 73:284-9.</u></li> <li>2. Bazil, V. <i>et al.</i> (1986) Biochemical characterization of a soluble for monocyte surface antigen. <u>Eur J Immunol. 16:1583-9.</u></li> <li>3. Jersmann, H.P. (2005) Time to abandon dogma: CD14 is express lineage cells. <u>Immunol Cell Biol. 83:462-7.</u></li> <li>4. Pugin, J. <i>et al.</i> (1994) CD14 is a pattern recognition receptor. <u>Imm</u></li> <li>5. Dziarski, R. <i>et al.</i> (1998) Binding of bacterial peptidoglycan to CD <u>273:8680-90.</u></li> <li>6. Piriou-Guzylack, L. (2008) Membrane markers of the immune cell <u>Vet Res. 39: 54.</u></li> </ul>	ed by non-myeloid <u>nunity.1:509-16.</u> 14. <u>J Biol Chem.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.	
Guarantee	12 months from date of despatch	
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/MCA1568F 10041	
Regulatory	For research purposes only	

## **Related Products**

### **Recommended Negative Controls**

MOUSE IgG2a NEGATIVE CONTROL:FITC (MCA929F)

### **Recommended Useful Reagents**

HUMAN SEROBLOCK (BUF070A) HUMAN SEROBLOCK (BUF070B)

North & South	Tel: +1 800 265 7376
America	Fax: +1 919 878 3751
	Email: antibody_sales_us@bio-

6 Worldwide 51 es\_us@bio-rad.com Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody\_sales\_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385471:210513'

### Printed on 04 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint