

## Datasheet: MCA156GA

<b>Description:</b>	MOUSE ANTI RAT MHC CLASS I RT1Ac
<b>Specificity:</b>	MHC CLASS I RT1Ac
<b>Format:</b>	Purified
<b>Product Type:</b>	Monoclonal Antibody
<b>Clone:</b>	OX-27
<b>Isotype:</b>	IgG2a
<b>Quantity:</b>	0.1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Rat
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Carrier Free</b>	Yes

<b>Approx. Protein Concentrations</b>	Isotype concentration 1.0 mg/ml
<b>Immunogen</b>	PHA activated rat lymphocytes.
<b>RRID</b>	AB_567197
<b>Fusion Partners</b>	Spleen cells from immunised BALB/c mice were fused with cells of the mouse NS1 myeloma cell line.
<b>Specificity</b>	<b>Mouse anti Rat MHC Class I RT1Ac antibody, clone OX-27</b> recognizes a polymorphic determinant (c haplotype) of rat Class I MHC Antigen (RT-1A).
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Hikita, N. <i>et al.</i> (1997) Use of topical FK506 in a corneal graft rejection model in Lewis rats. <a href="#">Invest Ophthalmol Vis Sci. 38 (5): 901-9.</a></li> <li>Sharland, A. <i>et al.</i> (1999) Evidence that apoptosis of activated T cells occurs in spontaneous tolerance of liver allografts and is blocked by manipulations which break tolerance. <a href="#">Transplantation. 68:1736-45.</a></li> <li>Huang, W.C. <i>et al.</i> (2010) Vascularized bone grafts within composite tissue allotransplants can autogenerate tolerance through mixed chimerism with partial myeloablative conditioning: an experimental study in rats. <a href="#">Plast Reconstr Surg. 125 (4): 1095-103.</a></li> <li>Wang Y <i>et al.</i> (2012) Role of donor-specific regulatory T cells in long-term acceptance of rat hind limb allograft. <a href="#">PLoS One. 7 (8): e43825.</a></li> <li>Liu, Q. <i>et al.</i> (2013) Heart allograft tolerance induced and maintained by vascularized hind-limb transplant in rats. <a href="#">Clin Dev Immunol. 2013: 483856.</a></li> <li>Zhu, H. <i>et al.</i> (2015) Rat model of heterotopic toe allotransplantation. <a href="#">J Surg Res. 199 (2): 707-17.</a></li> <li>Gu, C. <i>et al.</i> (2016) Triptolide Reduces the Required Dose of Tacrolimus by Attenuating Inflammation, Enhancing Immunosuppression, and Increasing Donor Chimerism in a Heterotopic Hindlimb Transplantation Model. <a href="#">Plast Reconstr Surg. 138 (6): 1243-1253.</a></li> <li>von Websky, M.W. <i>et al.</i> (2016) Recombinant HLA-G as Tolerogenic Immunomodulant in Experimental Small Bowel Transplantation. <a href="#">PLoS One. 11 (7): e0158907.</a></li> <li>Schweizer, R. <i>et al.</i> (2020) Adipose-derived stromal cell therapy combined with a short course nonmyeloablative conditioning promotes long-term graft tolerance in vascularized composite allotransplantation. <a href="#">Am J Transplant. 20 (5): 1272-84.</a></li> </ol>
<b>Storage</b>	<p>This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.</p> <p>Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.</p>
<b>Guarantee</b>	12 months from date of despatch

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at:  
<https://www.bio-rad-antibodies.com/SDS/MCA156GA>  
10040

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) [HRP](#)  
Rabbit Anti Mouse IgG (STAR12...) [RPE](#)  
Goat Anti Mouse IgG IgA IgM (STAR87...) [Alk. Phos.](#), [HRP](#)  
Goat Anti Mouse IgG (STAR76...) [RPE](#)  
Goat Anti Mouse IgG (Fc) (STAR120...) [FITC](#), [HRP](#)  
Rabbit Anti Mouse IgG (STAR13...) [HRP](#)  
Rabbit Anti Mouse IgG (STAR9...) [FITC](#)  
Goat Anti Mouse IgG (H/L) (STAR117...) [Alk. Phos.](#), [DyLight@488](#), [DyLight@550](#),  
[DyLight@650](#), [DyLight@680](#), [DyLight@800](#),  
[FITC](#), [HRP](#)  
Goat Anti Mouse IgG (STAR70...) [FITC](#)

### Recommended Negative Controls

[MOUSE IgG2a NEGATIVE CONTROL \(MCA1210\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M405388:220916'

Printed on 12 Aug 2023

---

© 2023 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)