

Datasheet: MCA1576F

Specificity:	CD8
•	
Format:	FITC
Product Type:	Monoclonal Antibody
Clone:	12.C7
Isotype:	lgG1
Quantity:	0.1 mg

# **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat - 1/10

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Rabbit				
Product Form	Purified IgG conjugat	niocyanate Isomer 1 (FITC) - liqu			
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)		
	FITC	490	525		
Preparation	Purified IgG prepared by affinity chromatography on Protein A from tissue cultu supernatant				
Buffer Solution	Phosphate buffered saline				
Preservative	0.09% Sodium Azide (NaN <sub>3</sub> )				
Stabilisers	1% Bovine Serum Albumin				
Approx. Protein Concentrations	IgG concentration 0.1	l mg/ml			
RRID	AB_566891				

#### Specificity

**Mouse anti Rabbit CD8 antibody, clone 12.C7** recognizes the rabbit CD8 cell surface antigen, expressed by a subset of T lymphocytes with cytotoxic/suppressor activity.

### **Flow Cytometry**

Use 10µl of the suggested working dilution to label 106 cells in 100µl

#### References

- 1. De Smet, W. *et al.* (1983) Rabbit leukocyte surface antigens defined by monoclonal antibodies. Eur J Immunol. 13: 919-28.
- 2. Wilkinson, J.M. *et al.* (1992) A cytotoxic rabbit T-cell line infected with a gamma-herpes virus which expresses CD8 and class II antigens. <a href="mailto:lmmunology.77">lmmunology.77</a>: 106-8.
- 3. Schock, A. and Reid, H.W. (1996) Characterisation of the lymphoproliferation in rabbits experimentally affected with malignant catarrhal fever. <u>Vet Microbiol. 53: 111-9.</u>
- 4. Dewals, B. *et al.* (2008) Malignant catarrhal fever induced by alcelaphine herpesvirus 1 is associated with proliferation of CD8+ T cells supporting a latent infection. <u>PLos ONE 3:</u> e1627.
- 5. Hanson, N.B. & Lanning, D.K. (2008) Microbial induction of B and T cell areas in rabbit appendix. <u>Dev Comp Immunol.</u> 32 (8): 980-91.
- 6. Anderson, I.E. *et al.* (2008) Production and utilization of interleukin-15 in malignant catarrhal fever. <u>J Comp Pathol. 138 (2-3): 131-44.</u>
- 7. Pakandl, M. *et al.* (2008) Dependence of the immune response to coccidiosis on the age of rabbit suckling. <u>Parasitol Res. 103 (6): 1265-71.</u>
- 8. Waclavicek, M. *et al.* (2009) Analysis of the early response to TSST-1 reveals Vbeta-unrestricted extravasation, compartmentalization of the response, and unresponsiveness but not anergy to TSST-1. <u>J Leukoc Biol. 85 (1): 44-54.</u>
- 9. Stich N *et al.* (2010) Staphylococcal superantigen (TSST-1) mutant analysis reveals that t cell activation is required for biological effects in the rabbit including the cytokine storm. <u>Toxins (Basel)</u>. 2 (9): 2272-88.
- 10. Dewals, B. *et al.* (2011) *Ex vivo* bioluminescence detection of alcelaphine herpesvirus 1 infection during malignant catarrhal fever. <u>J Virol. 85 (14): 6941-54.</u>
- 11. Zhao, L. *et al.* (2011) Evaluation of immunocompatibility of tissue-engineered periosteum. <u>Biomed Mater.6:015005.</u>
- 12. Dewals, B.G. & Vanderplasschen, A. (2011) Malignant catarrhal fever induced by Alcelaphine herpesvirus 1 is characterized by an expansion of activated CD3+CD8+CD4-T cells expressing a cytotoxic phenotype in both lymphoid and non-lymphoid tissues. <u>Vet Res. 42 (1): 95.</u>
- 13. Marques, R.M. *et al.* (2012) Early inflammatory response of young rabbits attending natural resistance to calicivirus (RHDV) infection. <u>Vet Immunol Immunopathol. 150: 181-8.</u>
- 14. Srivastava, R. *et al.* (2015) A Herpes Simplex Virus Type 1 Human Asymptomatic CD8+ T-Cell Epitopes-Based Vaccine Protects Against Ocular Herpes in a "Humanized" HLA Transgenic Rabbit Model. <u>Invest Ophthalmol Vis Sci. 56 (6): 4013-28.</u>
- 15. Myster, F. *et al.* (2015) Viral semaphorin inhibits dendritic cell phagocytosis and migration but is not essential for gammaherpesvirus-induced lymphoproliferation in malignant catarrhal fever. J Virol. 89 (7): 3630-47.
- 16. Khan AA *et al.* (2015) Therapeutic immunization with a mixture of herpes simplex virus 1 glycoprotein D-derived "asymptomatic" human CD8+ T-cell epitopes decreases spontaneous ocular shedding in latently infected HLA transgenic rabbits: association with low frequency of local PD-1+ TIM-3+ CD8+ exhausted T cells. <u>J Virol. 89 (13): 6619-32.</u>
  17. Srivastava, R. *et al.* (2016) The Herpes Simplex Virus Latency-Associated Transcript

Gene Is Associated with a Broader Repertoire of Virus-Specific Exhausted CD8+ T Cells Retained within the Trigeminal Ganglia of Latently Infected HLA Transgenic Rabbits. <u>J Virol. 90 (8): 3913-28.</u>

- 18. Khan, A.A. *et al.* (2018) Human Asymptomatic Epitope Peptide/CXCL10-Based Prime/Pull Vaccine Induces Herpes Simplex Virus-Specific Gamma Interferon-Positive CD107<sup>+</sup> CD8<sup>+</sup> T Cells That Infiltrate the Corneas and Trigeminal Ganglia of Humanized HLA Transgenic Rabbits and Protect against Ocular Herpes Challenge. <u>J Virol. 92 (16)</u>: e00535-18.
- 19. Gates, K.V. & Griffiths, L.G. (2018) Chronic graft-specific cell-mediated immune response toward candidate xenogeneic biomaterial. <a href="mailto:lmmunol-Res. 66">lmmunol-Res. 66</a> (2): 288-98.
- 20. Prakash, S. *et al.* (2020) Unique molecular signatures of antiviral memory CD8<sup>+</sup> T cells associated with asymptomatic recurrent ocular herpes. Sci Rep. 10 (1): 13843.
- 21. Jeklova, E. *et al.* (2020) Characterization of humoral and cell-mediated immunity in rabbits orally infected with *Encephalitozoon cuniculi*.. Vet Res. 51 (1): 79.
- 22. Niedźwiedzka-Rystwej, P. *et al.* (2020) B and T lymphocytes in rabbits change according to the sex and throughout the year. Pol J Vet Sci. 23 (1): 37-42.
- 23. Myster, F. *et al.* (2020) Alcelaphine herpesvirus 1 genes A7 and A8 regulate viral spread and are essential for malignant catarrhal fever. PLoS Pathog. 16 (3): e1008405.
- 24. Niedźwiedzka-Rystwej, P. *et al.* (2022) Reactivity of selected markers of innate and adaptive immunity in rabbits experimentally infected with antigenic variants of RHD (Lagovirus europaeus/GI.1a). Vet Res Commun. 46 (1): 233-42.
- 25. Niedźwiedzka-Rystwej, P. & Deptuła, W. (2023) Crosstalk between Apoptosis and Cytotoxic Lymphocytes (CTLs) in the Course of Lagovirus Europaeus GI.1a Infection in Rabbits. J Vet Res. 67 (1): 41-47.
- 26. Tansiri, Y. *et al.* (2021) New potent epitopes from Leptospira borgpetersenii for the stimulation of humoral and cell-mediated immune responses: Experimental and theoretical studies <u>Informatics in Medicine Unlocked. 25: 100649.</u>

# Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/MCA1576F">https://www.bio-rad-antibodies.com/SDS/MCA1576F</a> 10041
Regulatory	For research purposes only

## Related Products

## **Recommended Negative Controls**

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 America

Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.comd a Email: antibody\_sales\_us@bio-rad.com Email: antibody\_sales\_uk@bio-rad.com

То

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M421907:230809'

### Printed on 07 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint