

Datasheet: MCA1576GA

BATCH NUMBER 156447

Description:	MOUSE ANTI RABBIT CD8
Specificity:	CD8
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	12.C7
Isotype:	IgG1
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/100 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rabbit
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on protein A from tissue culture supernatant.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Carrier Free	Yes

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Specificity	Mouse anti Rabbit CD8 antibody, clone 12.C7 recognizes the rabbit CD8 cell surface antigen, expressed by a subset of T lymphocytes with cytotoxic/suppressor activity.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> 1. De Smet, W. <i>et al.</i> (1983) Rabbit leukocyte surface antigens defined by monoclonal antibodies. Eur J Immunol. 13: 919-28. 2. Dewals, B. <i>et al.</i> (2008) Malignant catarrhal fever induced by alcelaphine herpesvirus 1 is associated with proliferation of CD8+ T cells supporting a latent infection. PLoS ONE 3: e1627. 3. Zhao, L. <i>et al.</i> (2011) Evaluation of immunocompatibility of tissue-engineered periosteum. Biomed Mater.6:015005. 4. Wilkinson, J.M. <i>et al.</i> (1992) A cytotoxic rabbit T-cell line infected with a gamma-herpes virus which expresses CD8 and class II antigens. Immunology. 77: 106-8. 5. Marques, R.M. <i>et al.</i> (2012) Early inflammatory response of young rabbits attending natural resistance to calicivirus (RHDV) infection. Vet Immunol Immunopathol. 150: 181-8. 6. Schock, A. and Reid, H.W. (1996) Characterisation of the lymphoproliferation in rabbits experimentally affected with malignant catarrhal fever. Vet Microbiol. 53: 111-9. 7. Beghelli, D. <i>et al.</i> (2012) Phytoderivates in Rabbit Diet and Immune responses. Proceedings 10th World Rabbit Congress: 1019-23 8. Khan AA <i>et al.</i> (2015) Therapeutic immunization with a mixture of herpes simplex virus 1 glycoprotein D-derived "asymptomatic" human CD8+ T-cell epitopes decreases spontaneous ocular shedding in latently infected HLA transgenic rabbits: association with low frequency of local PD-1+ TIM-3+ CD8+ exhausted T cells. J Virol. 89 (13): 6619-32. 9. Srivastava, R. <i>et al.</i> (2015) A Herpes Simplex Virus Type 1 Human Asymptomatic CD8+ T-Cell Epitopes-Based Vaccine Protects Against Ocular Herpes in a "Humanized" HLA Transgenic Rabbit Model. Invest Ophthalmol Vis Sci. 56 (6): 4013-28. 10. Hanson, N.B. & Lanning, D.K. (2008) Microbial induction of B and T cell areas in rabbit appendix. Dev Comp Immunol. 32 (8): 980-91. 11. Waclavicek, M. <i>et al.</i> (2009) Analysis of the early response to TSST-1 reveals Vbeta-unrestricted extravasation, compartmentalization of the response, and unresponsiveness but not anergy to TSST-1. J Leukoc Biol. 85 (1): 44-54. 12. Dewals, B. <i>et al.</i> (2011) <i>Ex vivo</i> bioluminescence detection of alcelaphine herpesvirus 1 infection during malignant catarrhal fever. J Virol. 85 (14): 6941-54. 13. Anderson, I.E. <i>et al.</i> (2008) Production and utilization of interleukin-15 in malignant catarrhal fever. J Comp Pathol. 138 (2-3): 131-44. 14. Stich N <i>et al.</i> (2010) Staphylococcal superantigen (TSST-1) mutant analysis reveals that t cell activation is required for biological effects in the rabbit including the cytokine storm. Toxins (Basel). 2 (9): 2272-88. 15. Myster, F. <i>et al.</i> (2015) Viral semaphorin inhibits dendritic cell phagocytosis and migration but is not essential for gammaherpesvirus-induced lymphoproliferation in malignant catarrhal fever. J Virol. 89 (7): 3630-47. 16. Srivastava, R. <i>et al.</i> (2016) The Herpes Simplex Virus Latency-Associated Transcript Gene Is Associated with a Broader Repertoire of Virus-Specific Exhausted CD8+ T Cells

Retained within the Trigeminal Ganglia of Latently Infected HLA Transgenic Rabbits. [J Virol. 90 \(8\): 3913-28.](#)

17. Gates, K.V. & Griffiths, L.G. (2018) Chronic graft-specific cell-mediated immune response toward candidate xenogeneic biomaterial. [Immunol Res. 66 \(2\): 288-98.](#)

18. Prakash, S. *et al.* (2020) Unique molecular signatures of antiviral memory CD8⁺ T cells associated with asymptomatic recurrent ocular herpes. [Sci Rep. 10 \(1\): 13843.](#)

19. Jeklova, E. *et al.* (2020) Characterization of humoral and cell-mediated immunity in rabbits orally infected with *Encephalitozoon cuniculi*. [Vet Res. 51 \(1\): 79.](#)

Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/MCA1576GA 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Mouse IgG (STAR12...)	RPE
Goat Anti Mouse IgG IgA IgM (STAR87...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC , HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (STAR77...)	HRP
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos. , DyLight®488 , DyLight®550 , DyLight®650 , DyLight®680 , DyLight®800 , FITC , HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M365493:200529'

Printed on 07 May 2025